

LOCAL HÖLDER CONTINUITY FOR DOUBLY NONLINEAR PARABOLIC EQUATIONS

TUOMO KUUSI, JUHANA SILJANDER AND JOSÉ MIGUEL URBANO

ABSTRACT. We give a proof of the Hölder continuity of weak solutions of certain degenerate doubly nonlinear parabolic equations in measure spaces. We only assume the measure to be a doubling non-trivial Borel measure which supports a Poincaré inequality. The proof discriminates between large scales, for which a Harnack inequality is used, and small scales, that require intrinsic scaling methods.

1. INTRODUCTION

We consider the regularity issue for nonnegative weak solutions of the doubly nonlinear parabolic equation

$$\frac{\partial(u^{p-1})}{\partial t} - \nabla \cdot (|\nabla u|^{p-2} \nabla u) = 0, \quad 2 \leq p < \infty. \quad (1.1)$$

This equation is a prototype of a parabolic equation of p -Laplacian type. The solutions of this equation can be scaled by nonnegative factors, but in general we cannot add a constant to a solution so that the resulting function would be a solution to the same equation.

The purpose of this paper is to obtain a clear and transparent proof for the local Hölder continuity of nonnegative weak solutions of (1.1). Our work is a continuation to [17], where Harnack's inequality for the same equation is proved. See also [21], [11], [10] and [24]. However, since we cannot add constants to solutions, the Harnack estimates do not directly imply the local Hölder continuity. To show that our proof is based on a general principle, we consider the case where the Lebesgue measure is replaced by a more general Borel measure which is merely assumed to be doubling and to support a Poincaré inequality. In the weighted case, parabolic equations have earlier been studied in [1], [2] and [20]. See also [8].

This kind of doubly nonlinear equations have been considered by Vespri [23], Porzio and Vespri [19], and Ivanov [14], [15]. The known regularity proofs are based on the method of intrinsic scaling, originally introduced by DiBenedetto, and they seem to depend highly on the particular form of the equation. However, the passage from one equation to another is not completely clear. For other parabolic equations, the problem has been studied at length, see [4], [3], [7] and [22], and the references therein.

The difficulty with equation (1.1) is that there is a certain kind of dichotomy in its behavior. Correspondingly, the proof has been divided in

2000 *Mathematics Subject Classification.* Primary 35B65. Secondary 35K65, 35D10.

Key words and phrases. Hölder continuity, Caccioppoli inequality, intrinsic scaling, Harnack's inequality.

two complementary cases:

$$\text{Case I : } 0 \leq \text{ess inf } u \ll \text{ess osc } u$$

and

$$\text{Case II : } u^{p-2}u_t \approx Cu_t.$$

In large scales, i.e., in Case I, the scaling property of the equation dominates and, consequently, the reduction of the oscillation follows immediately from Harnack's inequality. In small scales, on the other hand, the oscillation is already very small and, thus, the solution itself is between two constants, the infimum and the supremum, whose difference is negligible. Correspondingly, the nonlinear time derivative term, which formally looks like $u^{p-2}u_t$, behaves like u_t and we end up with a p -parabolic type behavior. However, also in this case, we still need to modify the known arguments. In particular, the energy estimates are not available in the usual form and we need to use modified versions as in [5], [15] and [25].

Our argument also applies to doubly nonlinear equations of p -Laplacian type that are of the form

$$\frac{\partial(u^{p-1})}{\partial t} - \nabla \cdot \mathcal{A}(x, t, u, \nabla u) = 0.$$

For expository purposes, we only consider (1.1).

Very recently, a direct geometric method to obtain local Hölder continuity for parabolic equations has been developed in [6] and [9]. Despite the effort, the general picture remains unclear.

2. PRELIMINARIES

Let μ be a Borel measure and Ω be an open set in \mathbb{R}^d . The Sobolev space $H^{1,p}(\Omega; \mu)$ is defined to be the completion of $C^\infty(\Omega)$ with respect to the Sobolev norm

$$\|u\|_{1,p,\Omega} = \left(\int_{\Omega} (|u|^p + |\nabla u|^p) d\mu \right)^{1/p}.$$

A function u belongs to the local Sobolev space $H_{loc}^{1,p}(\Omega; \mu)$ if it belongs to $H^{1,p}(\Omega'; \mu)$ for every $\Omega' \Subset \Omega$. Moreover, the Sobolev space with zero boundary values $H_0^{1,p}(\Omega; \mu)$ is defined as the completion of $C_0^\infty(\Omega)$ with respect to the Sobolev norm. For more properties of Sobolev spaces, see e.g. [13].

Let $t_1 < t_2$. The parabolic Sobolev space $L^p(t_1, t_2; H^{1,p}(\Omega; \mu))$ is the space of functions $u(x, t)$ such that, for almost every t , with $t_1 < t < t_2$, the function $u(\cdot, t)$ belongs to $H^{1,p}(\Omega; \mu)$ and

$$\int_{t_1}^{t_2} \int_{\Omega} (|u|^p + |\nabla u|^p) d\nu < \infty,$$

where we denote $d\nu = d\mu dt$.

The definition of the space $L_{loc}^p(t_1, t_2; H_{loc}^{1,p}(\Omega; \mu))$ is analogous.

Definition 2.1. A function $u \in L_{loc}^p(t_1, t_2; H_{loc}^{1,p}(\Omega; \mu))$ is a weak solution of equation (1.1) in $\Omega \times (t_1, t_2)$ if it satisfies the integral equality

$$\int_{t_1}^{t_2} \int_{\Omega} \left(|\nabla u|^{p-2} \nabla u \cdot \nabla \phi - u^{p-1} \frac{\partial \phi}{\partial t} \right) d\nu = 0 \quad (2.2)$$

for every $\phi \in C_0^\infty(\Omega \times (t_1, t_2))$.

Next, we recall a few definitions and results from analysis on metric measure spaces. The measure μ is doubling if there is a universal constant $D_0 \geq 1$ such that

$$\mu(B(x, 2r)) \leq D_0 \mu(B(x, r)),$$

for every $B(x, 2r) \subset \Omega$. Here

$$B(x, r) = \{y \in \mathbb{R}^d : |y - x| < r\}$$

denotes the standard open ball in \mathbb{R}^d . Let $0 < r < R < \infty$. A simple iteration of the doubling condition implies that

$$\frac{\mu(B(x, R))}{\mu(B(x, r))} \leq C \left(\frac{R}{r} \right)^{d_\mu},$$

where $d_\mu = \log_2 D_0$. A doubling measure in Ω also satisfies the following annular decay property. For every $0 < \alpha < 1$, there exists a constant $c \geq 1$ such that

$$\mu(B(x, r) \setminus B(x, (1 - \delta)r)) \leq c\delta^\alpha \mu(B(x, r)), \quad (2.3)$$

for all $B(x, r) \subset \Omega$ and $0 < \delta < 1$.

The measure is said to support a weak $(1, p)$ -Poincaré inequality if there exist constants $P_0 > 0$ and $\tau \geq 1$ such that

$$\int_{B(x, r)} |u - u_{B(x, r)}| d\mu \leq P_0 r \left(\int_{B(x, \tau r)} |\nabla u|^p d\mu \right)^{1/p},$$

for every $u \in H_{loc}^{1,p}(\Omega; \mu)$ and $B(x, \tau r) \subset \Omega$. Here, we denote

$$u_{B(x, r)} = \int_{B(x, r)} u d\mu = \frac{1}{\mu(B(x, r))} \int_{B(x, r)} u d\mu.$$

The word weak refers to the constant τ , that may be strictly greater than one. In \mathbb{R}^d with a doubling measure, the weak $(1, p)$ -Poincaré inequality with some $\tau \geq 1$ implies the $(1, p)$ -Poincaré inequality with $\tau = 1$, see Theorem 3.4 in [12]. Hence, we may assume that $\tau = 1$.

On the other hand, the weak $(1, p)$ -Poincaré inequality and the doubling condition imply a weak (κ, p) -Sobolev-Poincaré inequality with

$$\kappa = \begin{cases} \frac{d_\mu p}{d_\mu - p}, & 1 < p < d_\mu, \\ 2p, & p \geq d_\mu, \end{cases} \quad (2.4)$$

where d_μ is as above. In other words, Poincaré and doubling imply the Sobolev inequalities. More precisely, there are constants $C > 0$ and $\tau' \geq 1$ such that

$$\left(\int_{B(x, r)} |u - u_{B(x, r)}|^\kappa d\mu \right)^{1/\kappa} \leq Cr \left(\int_{B(x, \tau' r)} |\nabla u|^p d\mu \right)^{1/p}, \quad (2.5)$$

for every $B(x, \tau' r) \in \Omega$. The constant C depends only on p , D_0 and P_0 . For the proof, we refer to [12]. Again, by Theorem 3.4 in [12] we may take $\tau' = 1$ in (2.5).

For Sobolev functions with the zero boundary values, we have the following version of Sobolev's inequality. Suppose that $u \in H_0^{1,p}(B(x, r); \mu)$. Then

$$\left(\int_{B(x, r)} |u|^\kappa d\mu \right)^{1/\kappa} \leq Cr \left(\int_{B(x, r)} |\nabla u|^p d\mu \right)^{1/p}. \quad (2.6)$$

For the proof we refer, for example, to [18].

Moreover, by a recent result in [16], the weak $(1, p)$ -Poincaré inequality and the doubling condition also imply the $(1, q)$ -Poincaré inequality for some $q < p$, that is

$$\int_{B(x, r)} |u - u_{B(x, r)}| d\mu \leq Cr \left(\int_{B(x, r)} |\nabla u|^q d\mu \right)^{1/q}. \quad (2.7)$$

Consequently, also (2.5) holds with p replaced by q . We also obtain the (q, q) -Poincaré inequality for some $q < p$.

In the sequel, we shall refer to *data* as the set of a priori constants p , d , D_0 , and P_0 .

Our main result is the following theorem.

Theorem 2.8. *Let $2 \leq p < \infty$ and assume that the measure is doubling, supports a weak $(1, p)$ -Poincaré inequality and is non-trivial in the sense that the measure of every non-empty open set is strictly positive and the measure of every bounded set is finite. Moreover, let $u \geq 0$ be a weak solution of equation (1.1) in \mathbb{R}^d . Then u is locally Hölder continuous.*

We will use the following notation for balls and cylinders, respectively:

$$B(r) = B(0, r)$$

and

$$Q_t(s, r) = B(r) \times (t - s, t).$$

For simplicity, we will also denote

$$Q(s, r) = Q_0(s, r) = B(r) \times (-s, 0).$$

Recall Harnack's inequality from [17].

Theorem 2.9. *Let $1 < p < \infty$ and suppose that the measure μ is doubling and supports a weak $(1, p)$ -Poincaré inequality. Moreover, let $u \geq 0$ be a weak solution to (1.1) in \mathbb{R}^d . Then there exists a constant $H_0 = H_0(p, d, D_0, P_0, (t - (s - r^p))/r^p) \geq 2$ such that*

$$\operatorname{ess\,sup}_{Q_t(r^p, r)} u \leq H_0 \operatorname{ess\,inf}_{Q_s(r^p, r)} u$$

where $s > t + r^p$.

Proof. See [17]. □

In addition, in [17] it is also proved that all solutions of equation (1.1) are locally bounded. In the sequel, we will assume this knowledge without further comments.

3. CONSTRUCTING THE SETTING

Our proof is based on the known classical argument of reducing the oscillation, see [4], [7] and [22]. However, the equation under study has some intrinsic properties which are not present, for instance, in the case of the p -parabolic equation. In large scales, the scaling property dominates and the oscillation reduction follows easily from Harnack's inequality. In this case, the equation resembles the usual heat equation.

In small scales, in turn, the equation changes its behavior to look more like the evolution p -Laplace equation. Indeed, when we zoom in by reducing the oscillation, the infimum and the supremum get closer and closer to each other. Consequently, the weight u^{p-2} in the time derivative term starts to behave like a constant coefficient and we end up with a p -parabolic type behavior. Resembling this divide between large and small scales, the proof has to be divided in two cases.

We study the (local) Hölder continuity in a compact set K and we choose the following numbers accordingly. Let

$$\mu_0^- \leq \operatorname{ess\,inf}_K u \quad \text{and} \quad \mu_0^+ \geq \operatorname{ess\,sup}_K u,$$

and define

$$\omega_0 = \mu_0^+ - \mu_0^-.$$

Furthermore, choose μ_0^- small enough so that

$$(2H_0 + 1)\mu_0^- \leq \omega_0 \tag{3.1}$$

holds. We will construct an increasing sequence $\{\mu_i^-\}$ and a decreasing sequence $\{\mu_i^+\}$ such that

$$\mu_i^+ - \mu_i^- = \omega_i = \sigma^i \omega_0$$

for some $0 < \sigma < 1$. Moreover, these sequences can be chosen so that

$$\operatorname{ess\,sup}_{Q^i} u \leq \mu_i^+$$

and

$$\operatorname{ess\,inf}_{Q^i} u \geq \mu_i^-,$$

for some suitable sequence $\{Q^i\}$ of cubes. Consequently,

$$\operatorname{ess\,osc}_{Q^i} u \leq \omega_i.$$

The cubes here can, and will be, chosen so that the size of the cubes decreases in a controllable way so that we can deduce the Hölder continuity. Observe also that if

$$(2H_0 + 1)\mu_{j_0}^- \leq \omega_{j_0}$$

fails for some j_0 , the above sequences have been chosen so that

$$\frac{\mu_j^+}{\mu_j^-} < 2H_0 + 2 \tag{3.2}$$

for all $j \geq j_0$.

We are studying the local Hölder continuity in a compact set K . Our aim is to show that the oscillation around any point in K reduces whenever we suitably decrease the size of the set where the oscillation is studied.

The next step is to iterate this reduction process. We end up with a sequence of cylinders Q^i . For all purposes, in the sequel, it is enough to study the cylinder $Q^0 := Q(\eta r^p, r)$ instead of the set K . Indeed, for any point in K we can build the sequence of suitable cylinders, but since we can always translate the equation, we can, without loss of generality, restrict the study to the origin.

The equation (1.1) has its own time geometry too, that we need to respect in the arguments. This is important, especially in small scales, when the equation resembles the evolution p -Laplace equation. We will use a scaling factor $\eta = 2^{\lambda_1(p-2)+1}$ in the time direction, where $\lambda_1 \geq 1$ is an a priori constant to be determined later.

4. FUNDAMENTAL ESTIMATES

We start the proof of Theorem 2.8 by proving the usual energy estimate in a slightly modified setting, which overcomes the problem that we cannot add constants to solutions, see [5], [15] and [25]. We introduce the auxiliary function

$$\begin{aligned} \mathcal{J}((u-k)_\pm) &= \pm \int_{k^{p-1}}^{u^{p-1}} \left(\xi^{1/(p-1)} - k \right)_\pm d\xi \\ &= \pm (p-1) \int_k^u (\xi-k)_\pm \xi^{p-2} d\xi \\ &= (p-1) \int_0^{(u-k)_\pm} (k \pm \xi)^{p-2} \xi d\xi. \end{aligned}$$

Hence, we have

$$\frac{\partial}{\partial t} \mathcal{J}((u-k)_\pm) = \pm \frac{\partial(u^{p-1})}{\partial t} (u-k)_\pm. \quad (4.1)$$

In the sequel, we will also need the following estimates. Clearly,

$$\begin{aligned} \mathcal{J}((u-k)_+) &= (p-1) \int_0^{(u-k)_+} (k + \xi)^{p-2} \xi d\xi \\ &\leq \frac{p-1}{2} (k + (u-k)_+)^{p-2} (u-k)_+^2 \\ &\leq \frac{p-1}{2} u^{p-2} (u-k)_+^2 \end{aligned} \quad (4.2)$$

and

$$\begin{aligned} \mathcal{J}((u-k)_+) &\geq (p-1) k^{p-2} \int_0^{(u-k)_+} \xi d\xi \\ &\geq (p-1) k^{p-2} \frac{(u-k)_+^2}{2}. \end{aligned} \quad (4.3)$$

Observe, that the assumption $p \geq 2$ is used here.

On the other hand,

$$\begin{aligned} \mathcal{J}((u-k)_-) &= (p-1) \int_0^{(u-k)_-} (k-\xi)^{p-2} \xi \, d\xi \\ &\geq \frac{(p-1)}{2} u^{p-2} (u-k)_-^2. \end{aligned} \quad (4.4)$$

Moreover,

$$\begin{aligned} \mathcal{J}((u-k)_-) &= (p-1) \int_0^{(u-k)_-} (k-\xi)^{p-2} \xi \, d\xi \\ &\leq (p-1) k^{p-2} \int_0^{(u-k)_-} \xi \, d\xi \\ &= (p-1) k^{p-2} \frac{(u-k)_-^2}{2}. \end{aligned} \quad (4.5)$$

Now we are ready for the fundamental energy estimate.

Lemma 4.6. *Let $u \geq 0$ be a weak solution of (1.1) and let $k \geq 0$. Then there exists a constant $C = C(p) > 0$ such that*

$$\begin{aligned} &\operatorname{ess\,sup}_{t_1 < t < t_2} \int_{\Omega} \mathcal{J}((u-k)_{\pm}) \varphi^p \, d\mu + \int_{t_1}^{t_2} \int_{\Omega} |\nabla(u-k)_{\pm} \varphi|^p \, d\nu \\ &\leq C \int_{t_1}^{t_2} \int_{\Omega} (u-k)_{\pm}^p |\nabla \varphi|^p \, d\nu + C \int_{t_1}^{t_2} \int_{\Omega} \mathcal{J}((u-k)_{\pm}) \varphi^{p-1} \left(\frac{\partial \varphi}{\partial t} \right)_+ \, d\nu, \end{aligned}$$

for every nonnegative $\varphi \in C_0^\infty(\Omega \times (t_1, t_2))$.

Proof. Let $t_1 < \tau_1 < \tau_2 < t_2$. We formally substitute the test function $\phi = \pm(u-k)_{\pm} \varphi^p$ in the equation and obtain

$$\begin{aligned} 0 &= \int_{\tau_1}^{\tau_2} \int_{\Omega} \left(|\nabla u|^{p-2} \nabla u \cdot \nabla \phi + \frac{\partial(u^{p-1})}{\partial t} \phi \right) \, d\nu \\ &= \int_{\tau_1}^{\tau_2} \int_{\Omega} |\nabla(u-k)|^{p-2} (\pm \nabla(u-k)_{\pm}) \cdot \nabla(\pm(u-k)_{\pm} \varphi^p) \, d\nu \\ &\quad \pm \int_{\tau_1}^{\tau_2} \int_{\Omega} \frac{\partial(u^{p-1})}{\partial t} (u-k)_{\pm} \varphi^p \, d\nu. \end{aligned} \quad (4.7)$$

Now the first term on the right-hand side can be estimated pointwise from below as

$$\begin{aligned} &|\nabla(u-k)|^{p-2} (\pm \nabla(u-k)_{\pm}) \cdot \nabla(\pm(u-k)_{\pm} \varphi^p) \\ &\geq |\nabla(u-k)_{\pm}|^p \varphi^p - p |\nabla(u-k)_{\pm}|^{p-1} \varphi^{p-1} (u-k)_{\pm} |D\varphi|, \end{aligned}$$

and the last term is estimated further by Young's inequality as

$$\begin{aligned} &-p |\nabla(u-k)_{\pm}|^{p-1} \varphi^{p-1} (u-k)_{\pm} |D\varphi| \\ &\geq -\frac{1}{2} |\nabla(u-k)_{\pm}|^p \varphi^p - C (u-k)_{\pm}^p |D\varphi|^p. \end{aligned}$$

Thus we have

$$\begin{aligned} & \frac{1}{2} \int_{\tau_1}^{\tau_2} \int_{\Omega} |\nabla(u-k)_{\pm} \varphi|^p d\nu \pm \int_{\tau_1}^{\tau_2} \int_{\Omega} \frac{\partial(u^{p-1})}{\partial t} (u-k)_{\pm} \varphi^p d\nu \\ & \leq C \int_{\tau_1}^{\tau_2} \int_{\Omega} (u-k)_{\pm}^p |\nabla \varphi|^p d\nu. \end{aligned} \quad (4.8)$$

Using (4.1) and integrating by parts,

$$\begin{aligned} & \pm \int_{\tau_1}^{\tau_2} \int_{\Omega} \frac{\partial(u^{p-1})}{\partial t} (u-k)_{\pm} \varphi^p d\nu = \int_{\tau_1}^{\tau_2} \int_{\Omega} \frac{\partial}{\partial t} \mathcal{J}((u-k)_{\pm}) \varphi^p d\nu \\ & = \left[\int_{\Omega} \mathcal{J}((u(x,t)-k)_{\pm}) \varphi^p(x,t) d\mu \right]_{t=\tau_1}^{\tau_2} \\ & \quad - p \int_{\tau_1}^{\tau_2} \int_{\Omega} \mathcal{J}((u-k)_{\pm}) \varphi^{p-1} \frac{\partial \varphi}{\partial t} d\nu. \end{aligned}$$

So we obtain

$$\begin{aligned} & \int_{\Omega} \mathcal{J}((u(x,\tau_2)-k)_{\pm}) \varphi^p(x,\tau_2) d\mu + \int_{\tau_1}^{\tau_2} \int_{\Omega} |\nabla(u-k)_{\pm} \varphi|^p d\nu \\ & \leq C \int_{\Omega} \mathcal{J}((u(x,\tau_1)-k)_{\pm}) \varphi^p(x,\tau_1) d\mu + C \int_{\tau_1}^{\tau_2} \int_{\Omega} (u-k)_{\pm}^p |\nabla \varphi|^p d\nu \quad (4.9) \\ & \quad + C \int_{\tau_1}^{\tau_2} \int_{\Omega} \mathcal{J}((u-k)_{\pm}) \varphi^{p-1} \left(\frac{\partial \varphi}{\partial t} \right)_{\pm} d\nu. \end{aligned}$$

Now we can drop the second term from the left hand side, let $\tau_1 \rightarrow t_1$, choose τ_2 such that

$$\int_{\Omega} \mathcal{J}((u(x,\tau_2)-k)_{\pm}) \varphi^p(x,\tau_2) d\mu \geq \frac{1}{2} \operatorname{ess\,sup}_{t_1 < t < t_2} \int_{\Omega} \mathcal{J}((u-k)_{\pm}) \varphi^p d\mu \quad (4.10)$$

and estimate the limits of integration on the right hand side of (4.9). On the other hand, we can also drop the first term on the left hand side of (4.9) and let $\tau_1 \rightarrow t_1$ and $\tau_2 \rightarrow t_2$. Summing the estimates for both terms gives the claim. \square

Let us denote

$$\psi_{\pm}(u) := \Psi(H_k^{\pm}, (u-k)_{\pm}, c) = \left(\ln \left(\frac{H_k^{\pm}}{c + H_k^{\pm} - (u-k)_{\pm}} \right) \right)_{\pm}.$$

The following logarithmic lemma is used in forwarding information in time.

Lemma 4.11. *Let $u \geq 0$ be a weak solution of equation (1.1). Then there exists a constant $C = C(p) > 0$ such that*

$$\begin{aligned} \operatorname{ess\,sup}_{t_1 < t < t_2} \int_{\Omega} u^{p-2} \psi_-^2(u) \varphi^p d\mu & \leq \int_{\Omega} k^{p-2} \psi_-^2(u)(x, t_1) \varphi^p(x) d\mu \\ & \quad + C \int_{t_1}^{t_2} \int_{\Omega} \psi_- |(\psi_-)'|^{2-p} |\nabla \varphi|^p d\nu \end{aligned}$$

and

$$\begin{aligned} \operatorname{ess\,sup}_{t_1 < t < t_2} \int_{\Omega} k^{p-2} \psi_+^2(u) \varphi^p \, d\mu &\leq \int_{\Omega} u^{p-2} \psi_+^2(u)(x, t_1) \varphi^p(x) \, d\mu \\ &\quad + C \int_{t_1}^{t_2} \int_{\Omega} \psi_+ |(\psi_+)'|^{2-p} |\nabla \varphi|^p \, d\nu. \end{aligned}$$

Above, $\varphi \in C_0^\infty(\Omega)$ is any nonnegative time-independent test function.

Proof. Choose

$$\phi_{\pm}(u) = \frac{\partial}{\partial u} (\psi_{\pm}^2(u)) \varphi^p$$

in the definition of weak solution and observe that

$$(\psi_{\pm}^2)'' = (1 + \psi_{\pm})(\psi'_{\pm})^2. \quad (4.12)$$

The parabolic term will take the form

$$\begin{aligned} \int_{t_1}^{t_2} \int_{\Omega} \frac{\partial}{\partial t} u^{p-1} \phi_{\pm}(u) \, d\nu &= \int_{t_1}^{t_2} \int_{\Omega} \frac{\partial}{\partial t} \int_{k^{p-1}}^{u^{p-1}} \phi_{\pm}(s^{1/(p-1)}) \, ds \, d\nu \\ &= \left[\int_{\Omega} \int_{k^{p-1}}^{u^{p-1}} \phi_{\pm}(s^{1/(p-1)}) \, ds \, d\mu \right]_{t_1}^{t_2} \\ &= \left[(p-1) \int_{\Omega} \int_k^u \phi_{\pm}(r) r^{p-2} \, dr \, d\mu \right]_{t_1}^{t_2}. \end{aligned}$$

Now an integration by parts gives

$$\begin{aligned} \int_k^u \phi_{\pm}(r) r^{p-2} \, dr &= \int_k^u (\psi_{\pm}^2(r))' r^{p-2} \, dr \varphi^p \\ &= \varphi^p [\psi_{\pm}^2(r) r^{p-2}]_k^u - (p-2) \int_k^u \psi_{\pm}^2(r) r^{p-3} \, dr \varphi^p \\ &= \psi_{\pm}^2(u) u^{p-2} \varphi^p - (p-2) \int_k^u \psi_{\pm}^2(r) r^{p-3} \, dr \varphi^p. \end{aligned}$$

In the plus case, we have

$$\begin{aligned} \int_k^u \phi_+(r) r^{p-2} \, dr &\geq \psi_+^2(u) u^{p-2} \varphi^p - \psi_+^2(u) (u^{p-2} - k^{p-2}) \varphi^p \\ &= (p-1) \psi_+^2(u) k^{p-2} \varphi^p \end{aligned}$$

and trivially

$$\int_k^u \phi_+(r) r^{p-2} \, dr \leq \psi_+^2(u) u^{p-2} \varphi^p,$$

since $p \geq 2$. Similar estimates are true also for the minus case.

On the other hand, by using (4.12) together with Young's inequality, we obtain

$$\begin{aligned} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi_{\pm} &= |\nabla u|^{p-2} \nabla u \cdot \nabla ((\psi_{\pm}^2(u))' \varphi^p) \\ &\geq |\nabla u|^p (1 + \psi_{\pm})(\psi'_{\pm})^2 \varphi^p - 2p |\nabla u|^{p-1} \psi_{\pm} \psi'_{\pm} \varphi^{p-1} |\nabla \varphi| \\ &\geq \frac{1}{2} |\nabla u|^p (1 + \psi_{\pm})(\psi'_{\pm})^2 \varphi^p - C \psi_{\pm} (\psi'_{\pm})^{2-p} |\nabla \varphi|^p, \end{aligned}$$

almost everywhere, from which the claim follows. \square

We will need the following notations in the next lemma, which is the most crucial part of the argument. Let

$$r_n = \frac{r}{2} + \frac{r}{2^{n+1}}, \quad Q_n^\pm = B_n \times T_n^\pm = B(r_n) \times (t^* - \gamma^\pm r_n^p, t^*)$$

and

$$A_n^\pm = \{(x, t) \in Q_n^\pm : \pm u(x, t) > \pm k_n^\pm\},$$

for $n = 0, 1, 2, \dots$.

Recall the definitions $\mu_i^+ - \mu_i^- = \omega_i = \sigma^i \omega_0$, where $\mu_i^+ \geq \text{ess sup}_{Q_i} u$ and $\mu_i^- \leq \text{ess inf}_{Q_i} u$ for $i \geq 1$. Observe, however, that we have to choose $\mu_0^+ \leq \text{ess sup}_K u$ and $\mu_0^- \geq \text{ess inf}_K u$ where the infimum and the supremum are taken over K instead of Q^0 . This is because we need the argument to be independent of the initial cylinder Q^0 . Now we are ready to prove the fundamental lemma everything lies on.

Lemma 4.13. *Let $0 < \varepsilon_\pm \leq 1$, $(k_n^+)_n$ be an increasing sequence and $(k_n^-)_n$ a decreasing sequence, both of nonnegative real numbers. Suppose $u \geq 0$ is a weak solution of equation (1.1),*

$$(u - k_n^\pm)_\pm \leq \varepsilon_\pm \omega_i \quad \text{and} \quad |k_{n+1}^\pm - k_n^\pm| \geq \frac{\varepsilon_\pm \omega_i}{2^{n+2}}.$$

In addition, assume further

$$u \geq \frac{1}{C_0} k_n^- \tag{4.14}$$

and

$$\mu_i^+ \leq 2k_n^+, \quad n = 1, 2, \dots \tag{4.15}$$

for the minus and plus cases, respectively. Then there exist constants $C_- = C(D_0, P_0, C_0, p) > 0$ and $C_+ = C(D_0, P_0, p) > 0$ such that

$$\frac{\nu(A_{n+1}^\pm)}{\nu(Q_{n+1}^\pm)} \leq C_\pm^{m+1} \Gamma_\pm \left(\frac{\nu(A_n^\pm)}{\nu(Q_n^\pm)} \right)^{2-p/\kappa} \tag{4.16}$$

for every $n = 0, 1, 2, \dots$. Here κ is the Sobolev exponent as in (2.4) and

$$\Gamma_\pm = \frac{1}{\gamma^\pm} \left(\frac{k_n^\pm}{\varepsilon_\pm \omega_i} \right)^{p-2} \left(\gamma^\pm \left(\frac{\varepsilon_\pm \omega_i}{k_n^\pm} \right)^{p-2} + 1 \right)^{2-p/\kappa}.$$

Proof. Choose the cutoff functions $\varphi_n^\pm \in C_0^\infty(Q_n^\pm)$ so that $0 \leq \varphi_n^\pm \leq 1$, $\varphi_n^\pm = 1$ in Q_{n+1}^\pm and

$$|\nabla \varphi_n^\pm| \leq \frac{C2^{n+1}}{r} \quad \text{and} \quad \left| \frac{\partial \varphi_n^\pm}{\partial t} \right| \leq \frac{C2^{p(n+1)}}{\gamma^\pm r^p}. \tag{4.17}$$

Denote in short

$$v_n = (u - k_n)_\pm, \quad k_n = k_n^\pm, \quad \varepsilon = \varepsilon_\pm$$

and

$$Q_n = B_n \times T_n = Q_n^\pm, \quad A_n = A_n^\pm, \quad \gamma = \gamma^\pm, \quad \varphi_n = \varphi_n^\pm.$$

By Hölder's inequality, together with the Sobolev inequality (2.6), we obtain

$$\begin{aligned}
& \int_{Q_{n+1}} v_n^{2(1-p/\kappa)+p} d\nu \\
& \leq \frac{\nu(Q_n)}{\nu(Q_{n+1})} \int_{Q_n} v_n^{2(1-p/\kappa)+p} \varphi_n^{p(1-p/\kappa)+p} d\nu \\
& \leq C \int_{T_n} \left(\int_{B_n} v_n^2 \varphi_n^p d\mu \right)^{1-p/\kappa} \left(\int_{B_n} (v_n \varphi_n)^\kappa d\mu \right)^{p/\kappa} dt \\
& \leq C r^p \left(\operatorname{ess\,sup}_{T_n} \int_{B_n} v_n^2 \varphi_n^p d\mu \right)^{1-p/\kappa} \int_{Q_n} |\nabla(v_n \varphi_n)|^p d\nu.
\end{aligned} \tag{4.18}$$

Here, we applied the doubling property of the measure ν giving

$$\frac{\nu(Q_n)}{\nu(Q_{n+1})} \leq C.$$

We continue by studying the term involving the essential supremum. By the assumption

$$u \geq \frac{1}{C_0} k_n^-$$

and (4.4), we obtain

$$(u - k_n^-)_-^2 \leq \frac{2}{p-1} u^{2-p} \mathcal{J}((u - k_n^-)_-) \leq C (k_n^-)^{2-p} \mathcal{J}((u - k_n^-)_-).$$

On the other hand, the lower bound (4.3) gives immediately

$$(u - k_n^+)_+^2 \leq C (k_n^+)^{2-p} \mathcal{J}((u - k_n^+)_+).$$

Using these estimates together with the energy estimate, Lemma 4.6, yields

$$\begin{aligned}
\operatorname{ess\,sup}_{T_n} \int_{B_n} v_n^2 \varphi_n^p d\mu & \leq C (k_n)^{2-p} \operatorname{ess\,sup}_{T_n} \int_{B_n} \mathcal{J}(v_n) \varphi_n^p d\mu \\
& \leq C (k_n)^{2-p} \gamma r_n^p \int_{Q_n} \left(v_n^p |\nabla \varphi_n|^p + \mathcal{J}(v_n) \varphi_n^{p-1} \left(\frac{\partial \varphi_n}{\partial t} \right)_+ \right) d\nu.
\end{aligned}$$

Furthermore, the estimates (4.2) and (4.5) imply

$$\mathcal{J}((u - k_n)_+) \leq C (k_n^+)^{p-2} (u - k_n)_+^2, \quad \mathcal{J}((u - k_n)_-) \leq C (k_n^-)^{p-2} (u - k_n)_-^2.$$

For the plus case, we used (4.15). Next, using (4.17), we arrive at

$$\begin{aligned}
& r_n^p \int_{Q_n} \left(v_n^p |\nabla \varphi_n|^p + \mathcal{J}(v_n) \varphi_n^{p-1} \left(\frac{\partial \varphi_n}{\partial t} \right)_+ \right) d\nu \\
& \leq C 2^{np} \int_{Q_n} \left(v_n^p + \frac{(k_n)^{p-2}}{\gamma} v_n^2 \right) d\nu \\
& \leq C 2^{np} (\varepsilon \omega_i)^p \left(1 + \frac{1}{\gamma} \left(\frac{\varepsilon \omega_i}{k_n} \right)^{2-p} \right) \frac{\nu(A_n)}{\nu(Q_n)},
\end{aligned} \tag{4.19}$$

where the last inequality follows from the fact that $(u - k_n)_\pm \leq \varepsilon_\pm \omega_i$. Thus, we conclude

$$\operatorname{ess\,sup}_{T_n} \int_{B_n} v_n^2 \varphi_n^p d\mu \leq C 2^{np} (\varepsilon \omega_i)^2 \left(\gamma \left(\frac{\varepsilon \omega_i}{k_n} \right)^{p-2} + 1 \right) \frac{\nu(A_n)}{\nu(Q_n)}. \tag{4.20}$$

Furthermore, since

$$\int_{Q_n} |\nabla(v_n \varphi_n)|^p d\nu \leq C \int_{Q_n} |\nabla v_n|^p \varphi_n^p d\nu + C \int_{Q_n} v_n^p |\nabla \varphi_n|^p d\nu,$$

applying again the energy estimate and (4.19) leads to

$$\int_{Q_n} |\nabla(v_n \varphi_n)|^p d\nu \leq C 2^{np} (\varepsilon \omega_i)^p \left(1 + \frac{1}{\gamma} \left(\frac{\varepsilon \omega_i}{k_n}\right)^{2-p}\right) \frac{\nu(A_n)}{\nu(Q_n)}. \quad (4.21)$$

To finish the proof, note first that

$$\begin{aligned} (u - k_n^\pm)_\pm \chi_{\{(u - k_n^\pm)_\pm > 0\}} &\geq (u - k_n^\pm)_\pm \chi_{\{(u - k_{n+1}^\pm)_\pm > 0\}} \\ &\geq |k_{n+1}^\pm - k_n^\pm| \\ &\geq 2^{-(n+2)} \varepsilon_\pm \omega_i. \end{aligned}$$

It then follows that

$$\int_{Q_{n+1}} v_n^{2(1-p/\kappa)+p} d\nu \geq \left(2^{-(n+2)} \varepsilon \omega_i\right)^{2(1-p/\kappa)+p} \frac{\nu(A_{n+1})}{\nu(Q_{n+1})}. \quad (4.22)$$

Inserting estimates (4.20), (4.21), and (4.22) into (4.18) concludes the proof. \square

Remark 4.23. If we have the extra knowledge that $(u - k_n^-)_- = 0$ almost everywhere in $B(r)$ at a given time level, we can choose the test functions independent of time and the cylinder Q_n^- so that the length in the time direction stays constant and the bottom of the cylinder stays at the given time level. In this case, by choosing a time independent test function, the right hand side of the energy estimate simplifies so that we can get rid of the term $+1$ in the formulation and

$$\Gamma_\pm = \left(\gamma^\pm \left(\frac{\varepsilon_\pm \omega_i}{k_n^\pm}\right)^{p-2}\right)^{1-p/\kappa}$$

in (4.16). This will get us the required extra room in the end of the first alternative of Case II.

Furthermore, in the previous lemma, we chose the radii of the cylinder as

$$r_n = \frac{r}{2} + \frac{r}{2^{n+1}}.$$

However, the factor 2 in the denominator can naturally be replaced by any greater number.

We start the proof by considering Case I. There, we use the previous lemma only in the plus case. Consequently, the first case does not depend on the constant C_0 .

We recall a lemma of ‘‘fast geometric convergence’’ from [4].

Lemma 4.24. *Let $(Y_n)_n$ be a sequence of positive numbers, satisfying*

$$Y_{n+1} \leq C b^n Y_n^{1+\alpha} \quad (4.25)$$

where $C, b > 1$ and $\alpha > 0$. Then $(Y_n)_n$ converges to zero as $n \rightarrow \infty$ provided

$$Y_0 \leq C^{-1/\alpha} b^{1-\alpha^2}. \quad (4.26)$$

On several occasions in the sequel, we use this lemma, together with the fundamental estimate Lemma 4.13, to conclude that a ratio of the form

$$Y_n = \frac{\nu(A_n^\pm)}{\nu(Q_n^\pm)}$$

converges to zero and consequently that $\nu(A_n^\pm) \rightarrow 0$ as $n \rightarrow \infty$. This will ultimately lead to a reduction of the oscillation which is our final goal.

Once a recursive inequality of type (4.16) has been established, the convergence to zero of $\nu(A_n^\pm)$ follows from the condition

$$\frac{\nu(A_0^\pm)}{\nu(Q_0^\pm)} \leq \alpha_0^\pm$$

with

$$\alpha_0^\pm = \Gamma_\pm^{-1/(1-p/\kappa)} C_\pm^{-1/(1-p/\kappa)+1-(1-p/\kappa)^2}, \quad (4.27)$$

where the constants C_\pm and Γ_\pm are the constants from the previous lemma. Note that an explicit value of α_0^\pm only follows after fixing C_\pm and Γ_\pm .

5. THE CASE I

Now we assume that (3.1) holds. Our aim is to show that the measures of certain distribution sets tend to zero and that the local Hölder continuity follows from this.

We start by studying the subcylinder $Q(r^p, r) \subset Q(\eta r^p, r)$. Let $\gamma^\pm = 1$, $\varepsilon_\pm = 2^{-1}$ and

$$k_n^+ = \mu_0^+ - \frac{\omega_0}{4} - \frac{\omega_0}{2^{n+2}}.$$

Observe that after fixing these quantities, the constant α_0^+ can be fixed, as well.

We will study two different alternatives which are considered in the following two lemmata, respectively.

Lemma 5.1. *Let $\lambda_2 > 1$ be sufficiently large and let $u \geq 0$ be a weak solution of equation (1.1). Furthermore, assume*

$$\nu \left(\left\{ (x, t) \in B(r) \times (-r^p, -\frac{r^p}{\lambda_2}) : u(x, t) \geq \mu_0^- + \frac{\omega_0}{2} \right\} \right) = 0. \quad (5.2)$$

Then there exists a constant $\sigma \in (0, 1)$ such that

$$\operatorname{ess\,osc}_{Q((\frac{r}{2})^p, \frac{r}{2})} u \leq \sigma \omega_0.$$

Proof. By the choices preceding the statement of this lemma, we have

$$(u - k_n^+)_+ \leq \varepsilon_+ \omega_0.$$

The assumption (3.1) implies

$$\mu_0^+ = \mu_0^- + \omega_0 \leq 2\omega_0.$$

Thus

$$1 \leq \frac{k_n^+}{\varepsilon_+ \omega_0} \leq 4.$$

Plug these in Lemma 4.13 to deduce

$$\frac{\nu(A_{n+1}^+)}{\nu(Q_{n+1}^+)} \leq C^{n+1} \left(\frac{\nu(A_n^+)}{\nu(Q_n^+)} \right)^{2-p/\kappa}.$$

On the other hand, by (5.2) we have the trivial estimate

$$\frac{\nu(A_0^+)}{\nu(Q_0^+)} \leq \frac{1}{\lambda_2} \leq \alpha_0^+,$$

choosing $\lambda_2 > 1$ sufficiently large. By Lemma 4.24 we conclude that

$$\frac{\nu(A_n^+)}{\nu(Q_n^+)} \rightarrow 0$$

as $n \rightarrow \infty$. This implies

$$\operatorname{ess\,sup}_{Q((\frac{r}{2})^p, \frac{r}{2})} u \leq \mu_0^+ - \frac{\omega_0}{4}.$$

So, if this alternative occurs, we choose

$$\mu_1^+ = \mu_0^+ - \frac{\omega_0}{4}$$

and

$$\mu_1^- = \mu_0^-.$$

These choices yield

$$\operatorname{ess\,osc}_{Q((\frac{r}{2})^p, \frac{r}{2})} u \leq \left(1 - \frac{1}{4}\right) \omega_0$$

as required, with

$$\sigma = \frac{3}{4}.$$

□

For the second possibility, we have the following lemma.

Lemma 5.3. *Let $u \geq 0$ be a weak solution of equation (1.1) and suppose*

$$\nu \left(\left\{ (x, t) \in B(r) \times (-r^p, -\frac{r^p}{\lambda_2}) : u(x, t) \geq \mu_0^- + \frac{\omega_0}{2} \right\} \right) > 0. \quad (5.4)$$

Then there exists a constant $\sigma = \sigma(H_0) \in (0, 1)$ such that

$$\operatorname{ess\,osc}_{Q((\frac{r}{2\lambda_2})^p, \frac{r}{2\lambda_2})} u \leq \sigma \omega_0.$$

Proof. By assumption (5.4), we have

$$\operatorname{ess\,sup}_{B(r) \times (-r^p, -\frac{r^p}{\lambda_2})} u \geq \mu_0^- + \frac{\omega_0}{2}.$$

Now we can use Harnack's inequality (Theorem 2.9), together with the Case I assumption (3.1), to deduce

$$\begin{aligned}
\operatorname{ess\,inf}_{Q\left(\left(\frac{r}{2\lambda_2}\right)^p, \frac{r}{2\lambda_2}\right)} u &\geq \frac{1}{H_0} \operatorname{ess\,sup}_{B(r)\times(-r^p, -\frac{r^p}{\lambda_2})} u \\
&\geq \frac{\mu_0^-}{H_0} + \frac{\omega_0}{2H_0} \\
&\geq \mu_0^- + \frac{\mu_0^-}{H_0} - \frac{\omega_0}{2H_0 + 1} + \frac{\omega_0}{2H_0} \\
&\geq \mu_0^- + \frac{\omega_0}{2H_0(2H_0 + 1)}.
\end{aligned}$$

Observe that the constant H_0 depends on λ_2 , but this does not matter since λ_2 depends only on the data.

Now, if we end up in this alternative, we choose

$$\mu_1^- = \mu_0^- + \frac{\omega_0}{2H_0(2H_0 + 1)}$$

and

$$\mu_1^+ = \mu_0^+.$$

We also obtain

$$\operatorname{ess\,osc}_{Q\left(\left(\frac{r}{2\lambda_2}\right)^p, \frac{r}{2\lambda_2}\right)} u \leq \omega_0 - \frac{\omega_0}{2H_0(2H_0 + 1)} = \sigma\omega_0,$$

with

$$\sigma = 1 - \frac{1}{2H_0(2H_0 + 1)},$$

as required. \square

6. THE CASE II

In Case II the equation looks like the evolution p -Laplace equation. In this case, we need to use the scaling factor η in the time geometry of our cylinders. The difficulty is now that we cannot use the Harnack principle anymore, as the lower bound it gives might be trivial. Indeed, the infimum can be bigger than the lower bound Harnack's inequality gives. On the other hand, we have the following kind of elliptic Harnack's inequality.

Suppose that j_0 is the first index for which assumption (3.1) does not hold. Then we have

$$\omega_{j_0} \leq \mu_{j_0}^+ \leq (2H_0 + 2)\mu_{j_0}^-. \quad (6.1)$$

Clearly, this Harnack's inequality is valid also for every subset of the initial cylinder $Q_{j_0} = Q(\eta r^p, r)$ and, consequently, for every $j \geq j_0$.

Recall, that $\omega_{j_0} = \sigma\omega_{j_0-1}$ and

$$\begin{aligned}
\frac{\omega_{j_0}}{(2H_0 + 2)} &\leq \mu_{j_0}^- \leq \mu_{j_0-1}^- + (1 - \sigma)\omega_{j_0-1} \\
&\leq (2 - \sigma)\omega_{j_0-1} \leq \frac{2 - \sigma}{\sigma}\omega_{j_0}.
\end{aligned} \quad (6.2)$$

Thus, we obtain

$$\frac{\sigma}{(2H_0 + 2)(2 - \sigma)} \leq \frac{\sigma}{2 - \sigma} \frac{\mu_{j_0}^-}{\omega_{j_0}} \leq 1$$

and, consequently,

$$Q\left(\frac{C_1}{(2H_0 + 2)^{p-2}} \eta r^p, r\right) \subset Q\left(C_1 \left(\frac{\mu_{j_0}^-}{\omega_{j_0}}\right)^{p-2} \eta r^p, r\right) \subset Q(\eta r^p, r),$$

where $C_1 = \sigma^{p-2}/(2 - \sigma)^{p-2}$. We will consider the cylinder

$$Q := Q\left(C_1 \left(\frac{\mu_{j_0}^-}{\omega_{j_0}}\right)^{p-2} \eta r^p, r\right).$$

By the above calculation, we have shrunk the cylinder by a factor which is controllable by the data.

In the sequel, we will denote

$$\theta = C_1 \left(\frac{\mu_{j_0}^-}{\omega_{j_0}}\right)^{p-2}. \quad (6.3)$$

Recall the definitions

$$Q_n^\pm = Q_{t^*}(\gamma^\pm r_n^p, r_n) = B_n \times T_n = B(r_n) \times (t^* - \gamma^\pm r_n^p, t^*)$$

and

$$A_n^\pm = \{(x, t) \in Q_n^\pm : \pm u > \pm k_n^\pm\}.$$

Now the proof will follow the classical argument of DiBenedetto, see [4] and [22], and is divided in two alternatives. In the first one, we assume that there is a suitable cylinder for which the set where u is close to its infimum is very small. In the second alternative we assume that this is not true.

6.1. The First Alternative. We first suppose that there exists a constant $\alpha_0 \in (0, 1)$ (to be determined in the course of the next lemma) such that

$$\nu\left(\{(x, t) \in Q_0^- : u < \mu_{j_0}^- + \frac{\omega_{j_0}}{2}\}\right) \leq \alpha_0 \nu(Q_0^-)$$

for a cylinder

$$Q_0^- = Q_{t^*}(\theta r^p, r) \subset Q(2^{\lambda_1(p-2)+1} \theta r^p, r).$$

Our aim is to use Lemma 4.13 to conclude for the reduction of the oscillation.

Lemma 6.4. *For every $s > 3$*

$$\begin{aligned} & \nu\left(\left\{(x, t) \in Q\left(\theta \left(\frac{r}{4}\right)^p, \frac{r}{4}\right) : u(x, t) < \mu_{j_0}^- + \frac{\omega_{j_0}}{2^s}\right\}\right) \\ & \leq C 2^{\lambda_1(p-2)} \frac{s-2}{(s-3)^2} \nu\left(Q\left(\theta \left(\frac{r}{4}\right)^p, \frac{r}{4}\right)\right), \end{aligned}$$

where θ is as in (6.3).

Proof. We start by using Lemma 4.13, with the choices

$$r_n = \frac{r}{2} + \frac{r}{2^{n+1}}, \quad k_n^- = \mu_{j_0}^- + \frac{\omega_{j_0}}{4} + \frac{\omega_{j_0}}{2^{n+2}},$$

$\varepsilon_- = 1/2$ and $\gamma^- = \theta$. We also need the assumption (6.1) to deduce that

$$u \geq \frac{1}{C_0}(\mu_{j_0}^- + 2(2H_0 + 2)\mu_{j_0}^-) \geq \frac{1}{C_0}(\mu_{j_0}^- + 2\omega_{j_0}) \geq \frac{1}{C_0}k_n^-, \quad (6.5)$$

with $C_0 = 3(2H_0 + 2)$. This knowledge is needed in Lemma 4.13. Now, after fixing ε_- , γ^- , k_n^- and C_0 we can fix α_0^- , see (4.27).

We also obtain, using (6.1) and (6.3), the bounds

$$\frac{2^{p-2}}{C_1} \leq \frac{1}{\gamma^-} \left(\frac{k_n^-}{\varepsilon_- \omega_{j_0}} \right)^{p-2} \leq \frac{2^{p-2}}{C_1} (2H_0 + 2)^{p-2}$$

and thus we can conclude

$$\frac{\nu(A_{n+1}^-)}{\nu(Q_{n+1}^-)} \leq C^{n+1} \left(\frac{\nu(A_n^-)}{\nu(Q_n^-)} \right)^{2-p/\kappa}.$$

By the assumption of this alternative, together with the lemma of fast geometric convergence (Lemma 4.24), we have $u > k$ almost everywhere in $Q_{t^*}(\theta(r/2)^p, r/2)$. Thus

$$(u-k)_- = 0$$

and consequently

$$\psi(u) := \left(\ln \left(\frac{H_k^-}{c + H_k^- - (u-k)_-} \right) \right)_+ = 0$$

almost everywhere in $Q_{t^*}(\theta(r/2)^p, r/2)$. Let

$$t_0 \leq -\theta \left(\frac{r}{4} \right)^p \quad (6.6)$$

be a time level such that this is true for almost every $x \in B(r/2)$.

Now our goal is to apply Lemma 4.11 with

$$k = \mu_{j_0}^- + \frac{\omega_{j_0}}{4}, \quad c = \frac{\omega_{j_0}}{2^s}$$

and

$$H_k^- = \operatorname{ess\,sup}_Q (u - k)_-$$

where $Q = Q(\eta\theta r^p, r)$. Choose $\varphi \in C_0^\infty(B(r/2))$ independent of time such that $0 \leq \varphi \leq 1$, $\varphi = 1$ in $B(r/4)$ and

$$|\nabla \varphi| \leq \frac{C}{r}.$$

In the set $\{u < \mu_{j_0}^- + \frac{\omega_{j_0}}{2^s}\}$, we have

$$\psi^2 \geq (s-3)^2 \ln^2 2,$$

and, on the other hand,

$$\psi \leq (s-2) \ln 2 \quad \text{and} \quad |\psi'|^{2-p} \leq \left(\frac{\omega_{j_0}}{2} \right)^{p-2}.$$

The use of these estimates in Lemma 4.11 gives

$$\begin{aligned}
& (\mu_{j_0}^-)^{p-2} (s-3)^2 \ln^2 2 \cdot \mu(\{x \in B(r/4) : u(x, t) < \mu_{j_0}^- + \frac{\omega_{j_0}}{2^s}\}) \\
& \leq \operatorname{ess\,sup}_{t_0 < t < 0} \int_{B(r/2)} u^{p-2} \psi^2(u)(x, t) \varphi^p(x) \, d\mu \\
& \leq \int_{B(r/2)} k^{p-2} \psi^2(u)(x, t_0) \varphi^p(x) \, d\mu \\
& + C \int_{t_0}^0 \int_{B(r/2)} \psi |\psi'|^{2-p} |\nabla \varphi|^p \, d\mu \, dt \\
& \leq C(s-2) \ln 2 \left(\frac{2^{\lambda_1} \omega_{j_0}}{2} \right)^{p-2} \theta \mu(B(r/4)) \\
& \leq C(s-2) \ln 2 \left(\frac{2^{\lambda_1} \mu_{j_0}^-}{2} \right)^{p-2} \mu(B(r/4)),
\end{aligned}$$

for almost every $t \in (t_0, 0)$. Observe that, in the third inequality, we plugged in $\eta = 2^{\lambda_1(p-2)+1}$. The claim follows by integrating this estimate over $(-\theta(r/4)^p, 0)$. \square

We conclude this alternative with the following two lemmata.

Lemma 6.7. *Let $u \geq 0$ be a weak solution of equation (1.1) and assume (6.1) holds. Then*

$$u \geq \mu_{j_0}^- + \frac{\omega_{j_0}}{2^{s+1}} \quad \text{a.e. in } Q\left(\theta \left(\frac{r}{8}\right)^p, \frac{r}{8}\right),$$

where s depends only upon the data, and θ is as in (6.3).

Proof. Let

$$r_n = \frac{r}{8} + \frac{r}{2^{n+3}},$$

$$Q_n^- = B_n \times T = B(r_n) \times (t_0, 0),$$

where t_0 is, as in the previous lemma, such that

$$(u-k)_-(x, t_0) = 0, \quad t_0 \leq -\theta \left(\frac{r}{4}\right)^p,$$

for a.e. $x \in B(r/2)$. Moreover, define

$$k_n^- = \mu_{j_0}^- + \frac{\omega_{j_0}}{2^{s+1}} + \frac{\omega_{j_0}}{2^{s+n+1}}.$$

In this case, we obtain

$$(u - k_n^-)_- \leq \varepsilon_- \omega_{j_0}, \quad \text{where } \varepsilon_- = \frac{1}{2^s}.$$

Observe also that $\gamma^- = -t_0/r^p \leq \eta\theta = 2^{\lambda_1(p-2)+1}\theta$.

We will substitute these in Lemma 4.13 and, taking into account Remark 4.23 and estimate (6.5), we conclude as before that

$$\frac{\nu(A_{n+1}^-)}{\nu(Q_{n+1}^-)} \leq C^{n+1} \left(\frac{2^{\lambda_1(p-2)}}{2^{s(p-2)}} \right)^{1-p/\kappa} \left(\frac{\nu(A_n^-)}{\nu(Q_n^-)} \right)^{2-p/\kappa}.$$

Now choose $s > \lambda_1$. Then, by Lemma 4.24, we have $\nu(A_n^-)/\nu(Q_n^-) \rightarrow 0$ as $n \rightarrow \infty$, provided $\nu(A_0^-)/\nu(Q_0^-)$ is small enough. On the other hand, by choosing s large enough, Lemma 6.4 guarantees that $\nu(A_0^-)/\nu(Q_0^-)$ can be chosen to be as small as we please.

This gives

$$u \geq \mu_{j_0}^- + \frac{\omega_{j_0}}{2^{s+1}} \quad \text{a.e. in } Q\left(|t_0|, \frac{r}{8}\right)$$

and hence the lemma is proved. \square

Lemma 6.8. *There exists $0 < \sigma < 1$, depending only upon the data, such that*

$$\operatorname{ess\,osc}_{Q(\theta(\frac{r}{8})^p, \frac{r}{8})} u \leq \sigma \omega_{j_0}.$$

Proof. By the previous lemma,

$$\operatorname{ess\,inf}_{Q(\theta(\frac{r}{8})^p, \frac{r}{8})} u \geq \mu_{j_0}^- + \frac{\omega_{j_0}}{2^{s+1}},$$

for some $s > 1$, which depends only upon the data and λ_1 . Observe that here we used the knowledge

$$t_0 \leq -\theta \left(\frac{r}{8}\right)^p.$$

If this alternative occurs, we again choose

$$\mu_{j_0+1}^- := \mu_{j_0}^- + \frac{\omega_{j_0}}{2^{s+1}}$$

and

$$\mu_{j_0+1}^+ = \mu_{j_0}^+.$$

Finally, we get

$$\operatorname{ess\,osc}_{Q(\theta(\frac{r}{8})^p, \frac{r}{8})} u \leq \left(1 - \frac{1}{2^{s+1}}\right) \omega_{j_0}$$

as required, with

$$\sigma = 1 - \frac{1}{2^{s+1}}.$$

\square

Remark 6.9. Here the choice of s is possible only after λ_1 has been determined in the second alternative. Nevertheless, both of them are a priori constants which can be assigned explicit values depending only upon the data.

6.2. The Second Alternative. In the second alternative, the assumption of the first alternative is not true. In this case, for every cylinder $Q_{t^*}(\theta r^p, r) \subset Q(\eta \theta r^p, r)$, we have

$$\frac{\nu\left(\{(x, t) \in Q_{t^*}(\theta r^p, r) : u(x, t) \geq \mu_{j_0}^- + \frac{\omega_{j_0}}{2}\}\right)}{\nu(Q_{t^*}(\theta r^p, r))} < (1 - \alpha_0), \quad (6.10)$$

where $\alpha_0 := \alpha_0^-$ is the same constant as in the first alternative. This implies that, for every $t^* \in (-(\eta - 1)\theta r^p, 0)$, there exists a time level t_0 with

$$t^* - \theta r^p \leq t_0 \leq t^* - \frac{\theta \alpha_0}{2} r^p$$

for which

$$\mu(\{x \in B(r) : u(x, t_0) > k_0^-\}) \leq \frac{1 - \alpha_0}{1 - \frac{\alpha_0}{2}} \mu(B(r)). \quad (6.11)$$

Indeed, otherwise we would have

$$\begin{aligned} & \nu(\{(x, t) \in Q_{t^*}(\theta r^p, r) : u(x, t) > k_0^-\}) \\ & \geq \int_{t^* - \theta r^p}^{t^* - \frac{\theta \alpha_0}{2} r^p} \mu(\{x \in B(r) : u(x, t) > k_0^-\}) dt \\ & > (1 - \alpha_0) \nu(Q_{t^*}(\theta r^p, r)), \end{aligned}$$

which contradicts (6.10).

This alternative is also based on Lemma 4.13. We choose λ_1 in the definition of k_n^+ large enough so that we can force $\nu(A_0^+)$ to be small compared to $\nu(Q_0^+)$.

We start with forwarding the information of (6.11) in time.

Lemma 6.12. *There exists $s^* > 0$, depending only upon the data, such that*

$$\mu\left(\left\{x \in B(r) : u(x, t) > \mu_{j_0}^+ - \frac{\omega_{j_0}}{2^{s^*}}\right\}\right) \leq \frac{1 - \frac{3\alpha_0}{4}}{1 - \frac{\alpha_0}{2}} \mu(B(r)).$$

for almost all $t \in (t_0, 0)$.

Proof. Let

$$c = \frac{\omega_{j_0}}{2^{s+n}}, \quad k = \mu_{j_0}^+ - \frac{\omega_{j_0}}{2^s}$$

and

$$H_k^+ = \operatorname{ess\,sup}_Q (u - k)_+,$$

where s and n will be chosen later and $Q := Q(\theta \eta r^p, r)$. Our aim is again to use Lemma 4.11 to forward the information in time. We will need some estimates for doing this.

Recall the definition

$$\psi_+(u) = \Psi(H_k^+, (u - k)_+, c) = \ln^+ \left(\frac{H_k^+}{c + H_k^+ - (u - k)_+} \right).$$

Trivially, we have

$$\psi_+(u) \leq \ln \left(\frac{\frac{\omega_{j_0}}{2^s}}{\frac{\omega_{j_0}}{2^{s+n}}} \right) = n \ln 2$$

and, on the other hand, in the set

$$\{u > \mu_{j_0}^+ - \frac{\omega_{j_0}}{2^{s+n}}\},$$

we get

$$\psi_+(u) \geq \ln \left(\frac{\frac{\omega_{j_0}}{2^s}}{\frac{\omega_{j_0}}{2^{s+n}} + \frac{\omega_{j_0}}{2^{s+n}}} \right) = (n - 1) \ln 2$$

The last estimate we need is

$$|(\psi_+)'|^{2-p} \leq \left(\frac{1}{c + H_k^+} \right)^{2-p} \leq 2^{p-2} \left(\frac{\omega_{j_0}}{2^s} \right)^{p-2}.$$

Let now $\varphi \in C_0^\infty(B(r))$ be a cutoff function which is independent of time and has the properties $0 \leq \varphi \leq 1$, $\varphi = 1$ in $B((1 - \delta)r)$ and

$$|\nabla\varphi| \leq \frac{1}{\delta r},$$

where $0 < \delta < 1$ is to be determined later.

Apply Lemma 4.11 with these choices to conclude

$$\begin{aligned} & (n-1)^2 \ln^2 2 \cdot \mu(\{x \in B((1-\delta)r) : u(x,t) > l\}) \\ & \leq \operatorname{ess\,sup}_{t_0 < t < t^*} \int_{B(r)} \psi_+^2(u)(x,t) \varphi^p(x) \, d\mu \\ & \leq \int_{B(r)} \left(\frac{u}{k}\right)^{p-2} \psi_+^2(u)(x,t_0) \varphi^p(x) \, d\mu \\ & + Ck^{2-p} \int_{t_0}^{t^*} \int_{B(r)} \psi_+ |(\psi_+)'|^{2-p} |\nabla\varphi|^p \, d\mu \, dt \\ & \leq n^2 \ln^2(2) \left(\frac{\mu_{j_0}^+}{\mu_{j_0}^+ - \frac{\mu_{j_0}^+}{2^s}}\right)^{p-2} \frac{1 - \alpha_0}{1 - \frac{\alpha_0}{2}} \mu(B(r) \times \{t_0\}) \\ & + C \frac{n \ln 2}{\delta^p} \left(\frac{\omega_{j_0}}{\mu_{j_0}^+}\right)^{p-2} \left(\frac{1}{2^s - 1}\right)^{p-2} \theta \mu(B(r)) \end{aligned}$$

for almost every $t \in (t_0, t^*)$. Observe, that in the third inequality we used (6.11).

Now, by the annular decay property (2.3), we have

$$\begin{aligned} & \mu(\{x \in B(r) : u(x,t) > l\}) \\ & \leq \mu(B(r) \setminus B((1-\delta)r)) + \mu(\{x \in B((1-\delta)r) : u(x,t) > l\}) \\ & \leq C\delta^\alpha \mu(B(r)) + \mu(\{x \in B((1-\delta)r) : u(x,t) > l\}). \end{aligned}$$

For the first term, we choose δ small enough so that

$$C\delta^\alpha < \frac{\alpha_0}{16 \left(1 - \frac{\alpha_0}{2}\right)}$$

and for the second term we use the previous estimate. Indeed, by choosing s and n large enough so that

$$\frac{1 - \alpha_0}{1 - \frac{\alpha_0}{2}} \frac{n^2}{(n-1)^2} \left(\frac{1}{1 - \frac{1}{2^s}}\right)^{p-2} \leq 1 - \frac{3\alpha_0}{8 \left(1 - \frac{\alpha_0}{2}\right)}$$

and

$$\frac{Cn}{(\ln 2)\delta^p(n-1)^2} \left(\frac{1}{2^s - 1}\right)^{p-2} \leq \frac{\alpha_0}{16 \left(1 - \frac{\alpha_0}{2}\right)},$$

we get the claim for almost every $t \in (t_0, t^*)$ with $s^* = s + n$. Recall that

$$t^* - \theta r^p \leq t_0 \leq t^* - \frac{\theta \alpha_0}{2} r^p.$$

Finally, since the above holds for every cylinder $Q_{t^*}(\theta r^p, r) \subset Q(\eta\theta r^p, r)$, we can conclude that for almost every

$$t \geq -\eta\theta r^p + \theta r^p - \frac{\theta\alpha_0 r^p}{2} = \left(-\eta + 1 - \frac{\alpha_0}{2}\right)\theta r^p,$$

we have

$$\mu(\{x \in B(r) : u(x, t) > l\}) < \frac{1 - \frac{3\alpha_0}{4}}{1 - \frac{\alpha_0}{2}} \mu(B(r)).$$

□

Remark 6.13. Now we can choose η so large that the previous lemma holds for almost every $t \in \left(-\frac{\eta\theta}{2}r^p, 0\right)$, i.e.

$$-\eta + 1 - \frac{\alpha_0}{2} \leq -\frac{\eta}{2}$$

and hence

$$\eta = 2^{\lambda_1(p-2)+1} \geq 2 - \alpha_0.$$

But this is always guaranteed for $\lambda_1 > 1$ and $p \geq 2$.

We are ready to prove the final estimate, which, together with Lemma 4.24, gives the reduction of the oscillation. Let

$$E_\varrho(t) = \{x \in B(r) : u(x, t) > \mu_{j_0}^+ - \frac{\omega_{j_0}}{2^g}\}$$

and

$$E_\varrho = \{(x, t) \in Q\left(\frac{\eta\theta}{2}r^p, r\right) : u(x, t) - \mu_{j_0}^+ > \frac{\omega_{j_0}}{2^g}\}.$$

Then we have the following lemma.

Lemma 6.14. *For every $\alpha_1 \in (0, 1)$, there exists $\lambda_1 > 0$ such that*

$$\frac{\nu(E_{\lambda_1})}{\nu(Q(\frac{\eta\theta}{2}r^p, r))} \leq \alpha_1.$$

Proof. Denote

$$h = \mu_{j_0}^+ - \frac{\omega_{j_0}}{2^{s+1}}$$

and

$$k = \mu_{j_0}^+ - \frac{\omega_{j_0}}{2^s},$$

where $s > 0$ will be chosen large. Let also

$$v = \begin{cases} h - k, & u \geq h, \\ u - k, & k < u < h, \\ 0, & u \leq k. \end{cases}$$

By the previous lemma, we can choose s large enough, namely $s \geq s^*$ where s^* is from the previous lemma, so that, for almost every $t \in \left(-\frac{\eta\theta}{2}r^p, 0\right)$, we have

$$\begin{aligned} \mu(x \in B(r) : v(x, t) = 0) &= \mu(\{x \in B(r) : u(x, t) \leq k\}) \\ &\geq \frac{\alpha_0}{4 - 2\alpha_0} \mu(B(r) \times \{t\}) \\ &\geq \frac{\alpha_0}{4} \mu(B(r)). \end{aligned}$$

Thus, for almost every $t \in (-\frac{\eta\theta}{2}r^p, 0)$, we obtain

$$v_{B(r)}(t) = \int_{B(r) \times \{t\}} v \, d\mu \leq \left(1 - \frac{\alpha_0}{4}\right) (h - k)$$

and, consequently,

$$h - k - v_{B(r)}(t) \geq \frac{\alpha_0}{4}(h - k).$$

Using the (q, q) -Poincaré inequality for some $q < p$ (see (2.7) and the remark after that), yields

$$\begin{aligned} (h - k)^q \mu(E_{s+1}(t)) &\leq \left(\frac{4}{\alpha_0}\right)^q \int_{B(r) \times \{t\}} |v - v_{B(r)}(t)|^q \, d\mu \\ &\leq Cr^q \int_{B(r) \times \{t\}} |\nabla v|^q \, d\mu = Cr^q \int_{E_s(t) \setminus E_{s+1}(t)} |\nabla u|^q \, d\mu, \end{aligned}$$

for almost every $t \in (-\frac{\eta\theta}{2}r^p, 0)$. The constant $(4/\alpha_0)^q$ above was absorbed into the constant C . Now we integrate the above inequality over time to get

$$(h - k)^q \nu(E_{s+1}) \leq Cr^q \int_{E_s \setminus E_{s+1}} |\nabla u|^q \, d\nu.$$

Next, we introduce a cutoff function $\varphi \in C_0^\infty(Q(\eta\theta r^p, 2r))$ such that $0 \leq \varphi \leq 1$, $\varphi = 1$ in $Q(\frac{\eta\theta}{2}r^p, r)$ and

$$|\nabla \varphi| \leq \frac{C}{r} \quad \text{and} \quad \left| \frac{\partial \varphi}{\partial t} \right| \leq \frac{C}{\eta\theta r^p}.$$

Now Hölder's inequality gives

$$\begin{aligned} (h - k)^q \nu(E_{s+1}) &\leq Cr^q \left(\int_{E_s \setminus E_{s+1}} |\nabla u|^p \, d\nu \right)^{q/p} \nu(E_s \setminus E_{s+1})^{1-q/p} \\ &\leq Cr^q \left(\int_{Q(\eta\theta r^p, 2r)} |\nabla(u-k)_+|^p \varphi^p \, d\nu \right)^{q/p} \nu(E_s \setminus E_{s+1})^{1-q/p}. \end{aligned}$$

By choosing $\lambda_1 > s \geq s^*$ in the definition of η large enough, the first factor on the right hand side can be estimated by Lemma 4.6 and (4.2) as

$$\begin{aligned} &\int_{Q(\eta\theta r^p, 2r)} |\nabla(u-k)_+|^p \varphi^p \, d\nu \\ &\leq C \int_{Q(\eta\theta r^p, 2r)} (u-k)_+^p |\nabla \varphi|^p \, d\nu \\ &\quad + C \int_{Q(\eta\theta r^p, 2r)} \mathcal{J}((u-k)_+) \varphi^{p-1} \left| \frac{\partial \varphi}{\partial t} \right| \, d\nu \tag{6.15} \\ &\leq \frac{C}{r^p} \left(\left(\frac{\omega_{j_0}}{2^s}\right)^{p-2} + \frac{(\mu_{j_0}^+)^{p-2}}{\eta\theta} \right) \int_{Q(\eta\theta r^p, 2r)} (u-k)_+^2 \, d\nu \\ &\leq \frac{C}{r^p} \left(\frac{\omega_{j_0}}{2^s}\right)^p \nu \left(Q \left(\frac{\eta\theta}{2} r^p, r \right) \right). \end{aligned}$$

In the last inequality we used the doubling property of the measure ν .

We obtain

$$\left(\frac{\omega_{j_0}}{2^{s+1}}\right)^q \nu(E_{s+1}) \leq C \left(\frac{\omega_{j_0}}{2^s}\right)^q \nu\left(Q\left(\frac{\eta\theta}{2}r^p, r\right)\right)^{q/p} \nu(E_s \setminus E_{s+1})^{1-q/p}.$$

Finally, summing s over $s^*, \dots, \lambda_1 - 1$ gives

$$(\lambda_1 - s^*)\nu(E_{\lambda_1})^{p/(p-q)} \leq C\nu\left(Q\left(\frac{\eta\theta}{2}r^p, r\right)\right)^{q/(p-q)} \nu\left(Q\left(\frac{\eta\theta}{2}r^p, r\right)\right)$$

and hence

$$\nu(E_{\lambda_1}) \leq \frac{C}{(\lambda_1 - s^*)^{(p-q)/p}} \nu\left(Q\left(\frac{\eta\theta}{2}r^p, r\right)\right).$$

Choosing λ_1 large enough finishes the proof. \square

Lemma 6.16. *Suppose that (6.10) holds. Then there exists $0 < \sigma < 1$, depending only upon the data, such that*

$$\operatorname{ess\,osc}_{Q(\frac{\eta\theta}{2}r^p, r)} u \leq \sigma\omega_{j_0}.$$

Proof. Let

$$Q_n^+ = B(r_n) \times (-\gamma^+ r_n^p, 0),$$

$$r_n = \frac{r}{2} + \frac{r}{2^{n+1}}$$

and A_n^+ as before. Substituting $\gamma^+ = 2^{\lambda_1(p-2)}\theta$, $\varepsilon_+ = 1/2^{\lambda_1}$ and

$$k_n^+ = \mu_{j_0}^+ - \frac{\omega_{j_0}}{2^{\lambda_1+1}} - \frac{\omega_{j_0}}{2^{\lambda_1+n+1}}$$

in Lemma 4.13, and using (6.1) to bound

$$1 \leq \frac{1}{\gamma^+} \left(\frac{k_n^+}{\varepsilon_+ \omega_{j_0}}\right)^{p-2} \leq (2H_0 + 2)^{p-2},$$

yields

$$\frac{\nu(A_{n+1}^+)}{\nu(Q_{n+1}^+)} \leq C^{n+1} \left(\frac{\nu(A_n^+)}{\nu(Q_n^+)}\right)^{2-p/\kappa}.$$

By the previous Lemma, we can choose λ_1 large enough so that

$$\frac{\nu(A_0^+)}{\nu(Q_0^+)}$$

is as small as we please. Consequently, by Lemma 4.24, we obtain

$$\operatorname{ess\,sup}_{Q(\frac{\eta\theta}{2}(\frac{r}{2})^p, \frac{r}{2})} u \leq \mu_{j_0}^+ - \frac{\omega_{j_0}}{2^{\lambda_1+1}},$$

for some $\lambda_1 > 1$, which depends only upon the data. So if this alternative occurs, we choose

$$\mu_1^+ = \mu_{j_0}^+ - \frac{\omega_{j_0}}{2^{\lambda_1+1}}$$

and

$$\mu_1^- = \mu_{j_0}^-.$$

We also obtain

$$\operatorname{ess\,osc}_{Q(\frac{\eta\theta}{2}(\frac{r}{2})^p, \frac{r}{2})} u \leq \left(1 - \frac{1}{2^{\lambda_1+1}}\right) \omega_{j_0},$$

as required, with

$$\sigma = 1 - \frac{1}{2^{\lambda_1+1}}.$$

□

Now the Hölder continuity follows by standard iterative real analysis methods, see chapter III of [4], or [22].

REFERENCES

- [1] Filippo Chiarenza and Raul Serapioni. Degenerate parabolic equations and Harnack inequality. *Ann. Mat. Pura Appl. (4)*, 137:139–162, 1984.
- [2] Filippo M. Chiarenza and Raul P. Serapioni. A Harnack inequality for degenerate parabolic equations. *Comm. Partial Differential Equations*, 9(8):719–749, 1984.
- [3] Emmanuele DiBenedetto. On the local behaviour of solutions of degenerate parabolic equations with measurable coefficients. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 13(3):487–535, 1986.
- [4] Emmanuele DiBenedetto. *Degenerate parabolic equations*. Universitext. Springer-Verlag, New York, 1993.
- [5] Emmanuele DiBenedetto and Avner Friedman. Hölder estimates for nonlinear degenerate parabolic systems. *J. Reine Angew. Math.*, 357:1–22, 1985.
- [6] Emmanuele DiBenedetto, Ugo Gianazza, and Vincenzo Vespri. A geometric approach to the hölder continuity of solutions to certain singular parabolic partial differential equations. Preprint, 2010.
- [7] Emmanuele DiBenedetto, José Miguel Urbano, and Vincenzo Vespri. Current issues on singular and degenerate evolution equations. In *Evolutionary equations. Vol. I*, Handb. Differ. Equ., pages 169–286. North-Holland, Amsterdam, 2004.
- [8] Eugene B. Fabes, Carlos E. Kenig, and Raul P. Serapioni. The local regularity of solutions of degenerate elliptic equations. *Comm. Partial Differential Equations*, 7(1):77–116, 1982.
- [9] Ugo Gianazza, Mikhail Surnachev, and Vincenzo Vespri. A new proof of the hölder continuity of solutions to p -laplace type parabolic equations. Preprint, 2010.
- [10] Ugo Gianazza and Vincenzo Vespri. A Harnack inequality for solutions of doubly nonlinear parabolic equations. *J. Appl. Funct. Anal.*, 1(3):271–284, 2006.
- [11] Ugo Gianazza and Vincenzo Vespri. Parabolic De Giorgi classes of order p and the Harnack inequality. *Calc. Var. Partial Differential Equations*, 26(3):379–399, 2006.
- [12] Piotr Hajlasz and Pekka Koskela. Sobolev met Poincaré. *Mem. Amer. Math. Soc.*, 145(688):x+101, 2000.
- [13] Juha Heinonen, Tero Kilpeläinen, and Olli Martio. *Nonlinear potential theory of degenerate elliptic equations*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, 1993. , Oxford Science Publications.
- [14] Alexander V. Ivanov. Uniform hölder estimates for generalized solutions of quasilinear parabolic equations that admit double degeneration. *Algebra i Analiz*, 3(2):139–179, 1991. Translation in St. Petersburg Math. J. 3 (1992), no. 2, 363–403.
- [15] Alexander V. Ivanov. Hölder estimates for a natural class of equations of fast diffusion type hölder estimates for equations of fast diffusion type. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 229(11):29–322, 1995. Translation in J. Math. Sci. (New York) 89 (1998), no. 6, 1607–1630.
- [16] Stephen Keith and Xiao Zhong. The Poincaré inequality is an open ended condition. *Ann. of Math. (2)*, 167(2):575–599, 2008.
- [17] Juha Kinnunen and Tuomo Kuusi. Local behaviour of solutions to doubly nonlinear parabolic equations. *Math. Ann.*, 337(3):705–728, 2007.
- [18] Juha Kinnunen and Nageswari Shanmugalingam. Regularity of quasi-minimizers on metric spaces. *Manuscripta Math.*, 105(3):401–423, 2001.

- [19] Maria M. Porzio and Vincenzo Vespri. Hölder estimates for local solutions of some doubly nonlinear degenerate parabolic equations. *J. Differential Equations*, 103(1):146–178, 1993.
- [20] Mikhail Surnachev. A harnack inequality for weighted degenerate parabolic equations. To appear in *J. Differential Equations*, 2010.
- [21] Neil S. Trudinger. Pointwise estimates and quasilinear parabolic equations. *Comm. Pure Appl. Math.*, 21:205–226, 1968.
- [22] José Miguel Urbano. *The method of intrinsic scaling*, volume 1930 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2008. A systematic approach to regularity for degenerate and singular PDEs.
- [23] Vincenzo Vespri. On the local behaviour of solutions of a certain class of doubly nonlinear parabolic equations. *Manuscripta Math.*, 75(1):65–80, 1992.
- [24] Vincenzo Vespri. Harnack type inequalities for solutions of certain doubly nonlinear parabolic equations. *J. Math. Anal. Appl.*, 181(1):104–131, 1994.
- [25] Shulin Zhou. Parabolic Q -minima and their application. *J. Partial Differential Equations*, 7(4):289–322, 1994.

Addresses:

T.K.: Aalto University, Institute of Mathematics, P.O. Box 11100, FI-00076 Aalto, Finland.

E-mail: tuomo.kuusi@tkk.fi

J.S.: Aalto University, Institute of Mathematics, P.O. Box 11100, FI-00076 Aalto, Finland.

E-mail: juhana.siljander@tkk.fi

J.M.U.: CMUC, Department of Mathematics, University of Coimbra, 3001-454 Coimbra, Portugal.

E-mail: jmurb@mat.uc.pt