

On the equivalence of regularity criteria for triangular and tetrahedral finite element partitions

Jan Brandts, Sergey Korotov, Michal Křížek



On the equivalence of regularity criteria for triangular and tetrahedral finite element partitions

Jan Brandts, Sergey Korotov, Michal Křížek

Jan Brandts, Sergey Korotov, Michal Křížek : *On the equivalence of regularity criteria for triangular and tetrahedral finite element partitions* ; Helsinki University of Technology, Institute of Mathematics, Research Reports A505 (2006).

Abstract: *In this note we examine several regularity criteria for families of simplicial finite element partitions in \mathbf{R}^d , $d \in \{2, 3\}$. These are usually required in numerical analysis and computer implementations. We prove the equivalence of four different definitions of regularity often proposed in literature. The first one uses the volume of simplices. The others involve the inscribed and circumscribed ball conditions, and the minimal angle condition.*

AMS subject classifications: 65N12, 65N30

Keywords: finite element method, inscribed ball, circumscribed ball, regular family of simplicial partitions, Zlámal's condition

Correspondence

sergey.korotov@hut.fi

ISBN 951-22-8315-8
ISSN 0784-3143

Helsinki University of Technology
Department of Engineering Physics and Mathematics
Institute of Mathematics
P.O. Box 1100, 02015 HUT, Finland
email:math@hut.fi <http://www.math.hut.fi/>

1 Introduction

The finite element method (FEM) is nowadays one of the most powerful and popular numerical techniques widely used in various software packages that solve problems in, for instance, mathematical physics and mechanics. The initial step in FEM implementations is to establish an appropriate partition (also called mesh, grid, triangulation, etc.) on the solution domain. For a number of applications simplicial partitions are preferred over the others due to their flexibility. However, such partitions cannot be constructed arbitrarily from both theoretical and practical points of view. Thus, first of all we must ensure, at least theoretically, that the finite element approximations converge to the exact (weak) solution of the mathematical model under consideration when the associated partitions become finer. Mainly due to this reason the notions of regular families of partitions or nondegenerate partitions or shape-preserving partitions appeared. Second, the regularity is also important for real-life computations because degenerate partitions that contain flat elements may yield ill-conditioned stiffness matrices.

In 1968, Miloš Zlámal [19] introduced the so-called *minimal angle condition* that ensures the convergence of the finite element approximations for solving linear elliptic boundary value problems for $d = 2$. This condition requires that there exists a constant $\alpha_0 > 0$ such that the minimal angle α_S of each triangle S in all triangulations used satisfies

$$\alpha_S \geq \alpha_0.$$

Zlámal's condition can be generalized into \mathbf{R}^d for any $d \in \{2, 3, \dots\}$ so that all dihedral angles of simplicies and their lower-dimensional facets are bounded from below by a positive constant. Later, the so-called *inscribed ball condition* was introduced, see, e.g., [5, p. 124], which can also be used for nonsimplicial elements, and which has an elegant geometrical interpretation: the ratio of the radius of the inscribed ball of any element and the diameter of this element must be bounded from below by a positive constant over all partitions. Roughly speaking, no element of no partitions should degenerate to a hyperplane as the discretization parameter h (i.e., the maximal diameter of all elements in the corresponding partition \mathcal{T}_h) tends to zero. This property is called in [5] the *regularity of a family of partitions*. For triangular elements it is, obviously, equivalent to Zlámal's condition.

In 1985, Lin and Xu [17] introduced a somewhat stronger regularity assumption on triangular elements: each triangle $S \in \mathcal{T}_h$ contains a circle of radius $c_1 h$ and is contained in a circle of radius $c_2 h$, where c_1 and c_2 are positive constants that do not depend on S and h . Later, this assumption was modified as follows (see, e.g., [16]): a family of triangulations is called *strongly regular* if there exist two positive constants c_1 and c_2 such that for all $S \in \mathcal{T}_h$

$$c_1 h^2 \leq \text{meas}_2 S \leq c_2 h^2.$$

Notice that in this case no circle or angle conditions appear (cf. (1) and

(25)). Here and elsewhere in this paper meas_p stands for the p -dimensional measure.

Recently, in order to prove some superconvergence results, Brandts and Křížek [4] employed another regularity condition based on the circumscribed ball about simplicial elements, which is the unique sphere on which all vertices of the simplex lie (see (3)).

In the present paper we summarize the above proposed conditions into four different definitions of regularity and prove in detail that all these definitions are equivalent for simplicial elements in two and three dimensions.

2 Preliminaries

Let $\bar{\Omega} \subset \mathbf{R}^d, d \in \{1, 2, 3, \dots\}$ be a closed domain (i.e., the closure of a domain). If its boundary $\partial\bar{\Omega}$ is contained in a finite number of $(d - 1)$ -dimensional hyperplanes, we say that $\bar{\Omega}$ is *polytopic*. Moreover, if $\bar{\Omega}$ is bounded, it is called a *polytope*; in particular, $\bar{\Omega}$ is called a *polygon* for $d = 2$ and a *polyhedron* for $d = 3$.

A *simplex* S in \mathbf{R}^d is a convex hull of $d + 1$ points, A_1, A_2, \dots, A_{d+1} , that do not belong to the same hyperplane. We denote by h_S the length of the longest edge of S . Let F_i be the face of a simplex S opposite to the vertex A_i and let v_i be the altitude from the vertex A_i to the face F_i . For $d = 3$ angles between faces of a tetrahedron are called *dihedral*, whereas angles between its edges are called *solid*.

Next we define a *simplicial partition* \mathcal{T}_h over the polytope $\bar{\Omega} \subset \mathbf{R}^d$. We subdivide $\bar{\Omega}$ into a finite number of simplices (called *elements* or *simplicial elements*), so that their union is $\bar{\Omega}$, any two simplices have disjoint interiors and any facet of any simplex is a facet of another simplex from the partition or belongs to the boundary $\partial\bar{\Omega}$.

The set $\mathcal{F} = \{\mathcal{T}_h\}_{h>0}$ is called a *family of partitions* if for any $\varepsilon > 0$ there exists $\mathcal{T}_h \in \mathcal{F}$ with $h < \varepsilon$.

3 On the equivalence of various regularity conditions

The regularity conditions presented in the introduction can, in fact, be summarized into four conditions for the regularity of simplicial partitions which we will present below. In what follows, all constants C_i are independent of S and h , but can depend on the dimension $d \in \{2, 3\}$.

Condition 1: There exists a constant $C_1 > 0$ such that for any partition $\mathcal{T}_h \in \mathcal{F}$ and any simplex $S \in \mathcal{T}_h$ we have

$$\text{meas}_d S \geq C_1 h_S^d. \tag{1}$$

Condition 2: There exists a constant $C_2 > 0$ such that for any partition $\mathcal{T}_h \in \mathcal{F}$ and any simplex $S \in \mathcal{T}_h$ there exists a ball $B \subset S$ with radius r such that

$$r \geq C_2 h_S. \quad (2)$$

Condition 3: There exists a constant $C_3 > 0$ such that for any partition $\mathcal{T}_h \in \mathcal{F}$ and any simplex $S \in \mathcal{T}_h$ we have

$$\text{meas}_d S \geq C_3 \text{meas}_d B^S, \quad (3)$$

where $B^S \supset S$ is the circumscribed ball about S .

Condition 4: There exists a constant $C_4 > 0$ such that for any partition $\mathcal{T}_h \in \mathcal{F}$, any simplex $S \in \mathcal{T}_h$, and any dihedral or solid angle α of S , we have

$$\alpha \geq C_4. \quad (4)$$

Before we prove that the above four conditions are equivalent, we present three auxiliary lemmas.

Lemma 1. *For any simplex S , and any $i \in \{1, \dots, d+1\}$, $d \in \{2, 3\}$, we have*

$$\text{meas}_d S \leq h_S^d, \quad (5)$$

$$\text{meas}_{d-1} F_i \leq h_S^{d-1}, \quad (6)$$

$$v_i \leq h_S. \quad (7)$$

P r o o f: Relations (5) and (6) follow from the fact that the distance between any two points of a simplex S is not larger than h_S . Thus, S and any of its faces F_i (if $d = 3$), or edges F_i (if $d = 2$), are contained in a cube and a square with edges of length h_S . Inequality (7) is obvious. \square

Lemma 2. *For any simplex S we have*

$$2r^S \geq h_S > 2r, \quad (8)$$

where r^S is the radius of the circumscribed ball B^S about S , and r is a radius of any ball $B \subset S$.

P r o o f: Since $B \subset S \subset B^S$, their diameters are nondecreasing. The sharp inequality in (8) is evident. \square

Recall that for any $i \in \{1, \dots, d+1\}$ we have

$$\text{meas}_d S = \frac{1}{d} v_i \text{meas}_{d-1} F_i. \quad (9)$$

Lemma 3. *If condition (1) holds and $d \in \{2, 3\}$, then there exist positive constants C_5 , C_6 , and C_7 such that for any partition $\mathcal{T}_h \in \mathcal{F}$, any simplex $S \in \mathcal{T}_h$, and any $i \in \{1, \dots, d+1\}$, we have*

$$\text{meas}_{d-1} F_i \geq C_5 h_S^{d-1}, \quad (10)$$

$$v_i \geq C_6 h_S, \quad (11)$$

$$\sin \alpha \geq C_7, \quad (12)$$

where α is any dihedral or solid angle of the simplex S .

P r o o f: From (1), (9), and (7), we obtain

$$C_1 h_S^d \leq \text{meas}_d S = \frac{1}{d} v_i \text{meas}_{d-1} F_i \leq \frac{1}{d} h_S \text{meas}_{d-1} F_i, \quad (13)$$

which implies (10). Further, the inequality (11) follows from (13) if we use relation (6) in the last inequality of (13).

For any angle α of triangular elements or dihedral angle α for tetrahedral elements, we get

$$\sin \alpha \geq \frac{v_i}{h_S} \geq C_6.$$

Similar relations hold for the solid angles of the triangular faces of tetrahedron S (i.e., when $d = 3$), since in this case altitudes in the triangular faces are not greater than an appropriate altitude v_i of the tetrahedron. Thus, $C_7 = \arcsin C_6$. \square

Theorem 1. *For the dimension $d \in \{2, 3\}$, conditions 1, 2, 3, and 4 are equivalent.*

P r o o f: We prove that condition (1) is equivalent to each of conditions (2), (3), and (4).

(1) \implies (2): Let B_S be the inscribed ball of S with the radius r_S and the center O_S . We decompose S into $d+1$ subsimplicies – $\text{conv}\{O_S, F_i\}$, $i \in \{1, \dots, d+1\}$. All of them have the same altitude r_S , i.e., by (9), we get

$$\text{meas}_d S = \frac{1}{d} \sum_{i=1}^{d+1} r_S \text{meas}_{d-1} F_i. \quad (14)$$

Further, for any face of any simplex inequality (6) is valid, i.e., $d \text{meas}_d S \leq r_S(d+1)h_S^{d-1}$, and now using (1), we finally observe that

$$r_S(d+1)h_S^{d-1} \geq d \text{meas}_d S \geq C_1 d h_S^d, \quad (15)$$

which implies Condition 2 if we take $B = B_S$, $r = r_S$, and $C_2 = \frac{C_1 d}{d+1}$.

(2) \implies (1): Obviously, from the fact that $B \subset S$ and (2) we get

$$\text{meas}_d S \geq \text{meas}_d B \geq \pi r^d \geq \pi C_2^d h_S^d, \quad (16)$$

which implies Condition 1.

(1) \implies (3): From (1) and (5) we observe that $h_S^d \geq \text{meas}_d S \geq C_1 h_S^d$. Also, $\text{meas}_d B^S = C_8(d)(r^S)^d$, where $C_8(2) = \pi$ and $C_8(3) = \frac{4}{3}\pi$. We prove that under condition (1), there exists a constant $C_9 > 0$ such that for any simplex S from any partition $\mathcal{T}_h \in \mathcal{F}$ we have

$$r^S \leq C_9 h_S. \quad (17)$$

If (17) holds, then using (1) we immediately prove (3) as follows

$$\text{meas}_d S \geq C_1 h_S^d \geq C_1 \frac{(r^S)^d}{C_9^d} = \frac{C_1}{C_9^d C_8(d)} \text{meas}_d B^S. \quad (18)$$

Consider first the case $d = 2$. Let S denote the triangular element $A_1 A_2 A_3$. It is well-known that

$$r^S = \frac{|A_1 A_2| \cdot |A_2 A_3| \cdot |A_1 A_3|}{4 \text{meas}_2 S}. \quad (19)$$

Then in view of (1) (for $d = 2$) and the fact that any edge of S is of a length not greater than h_S we have

$$r^S \leq \frac{h_S^3}{4C_1 h_S^2} = \frac{1}{4C_1} h_S = C_9 h_S.$$

For the case $d = 3$ we use the following formula for the calculation of the circumradius presented in [6, p. 316] (cf. [14, p. 212]) for the tetrahedral element $S = A_1 A_2 A_3 A_4$

$$r^S = \frac{\sqrt{V_S}}{24 \text{meas}_3 S}, \quad (20)$$

where

$$V_S = 2l_1^2 l_2^2 l_4^2 l_5^2 + 2l_1^2 l_3^2 l_4^2 l_6^2 + 2l_2^2 l_3^2 l_5^2 l_6^2 - l_1^4 l_4^4 - l_2^4 l_5^4 - l_3^4 l_6^4. \quad (21)$$

In the above l_p and l_{p+3} are the lengths of opposite edges of S , $p = 1, 2, 3$. Obviously, using again the fact that $l_j \leq h_S$, $j = 1, \dots, 6$, we have $V_S \leq 6h_S^8$. Thus, from (1) (for $d = 3$) and (20) we get

$$r^S \leq \frac{\sqrt{6} h_S^8}{24 C_1 h_S^3} = \frac{\sqrt{6}}{24 C_1} h_S = C_9 h_S.$$

(3) \implies (1): In view of (3) and (8) we observe that

$$\text{meas}_d S \geq C_3 \text{meas}_d B^S = C_3 C_8(d) (r^S)^d \geq \frac{C_3 C_8(d)}{2^d} h_S^d, \quad (22)$$

which implies Condition 1.

(1) \implies (4): See (12).

(4) \implies (1): First we consider the case $d = 2$. Let S be again the triangular element $A_1 A_2 A_3$ and let $h_S = |A_1 A_3|$. Now, we cut out of the edge $A_1 A_3$




Figure 1: Illustration for the proof (4) \implies (1).

the segment $|MN|$ of the length $\frac{h_S}{2}$ with the endpoints M and N be at the distance $\frac{h_S}{4}$ from the vertices A_1 and A_3 , respectively (see the left of Figure 1).

Thus, $|A_1M| = |NA_3| = \frac{h_S}{4}$. Since the angles $\angle A_2A_1A_3$ and $\angle A_2A_3A_1$ are bounded from above and below due to (4), we can form a rectangle $KLMN$ inside of S (see the shadowed area in Figure 1 (left)) so that $|MK| = |LN| = \frac{h_S}{4} \tan C_4$. It is clear then that $\text{meas}_2 S \geq \text{meas}_2 KLMN = \frac{h_S}{2} \frac{h_S}{4} \tan C_4 = C_1 h_S^2$, where $C_1 = \frac{1}{8} \tan C_4$.

Consider now the case $d = 3$ and let S denote a tetrahedron $A_1A_2A_3A_4$ (see Figure 1 (right)). Using the above argumentation for the triangular faces $A_1A_2A_3$ and $A_1A_3A_4$ we can form two rectangles $KLMN$ and $PQNM$ with areas equal to $\frac{1}{8} h_S^2 \tan C_4$. Further, we consider the prism $KLNQPM$, which is inside of the tetrahedron S . Thus, $\text{meas}_3 S \geq C_1 h_S^3$, where $C_1 = \frac{1}{64} \tan^2 C_4 \sin C_4$, due to boundedness of the dihedral angle between faces $A_1A_2A_3$ and $A_1A_3A_4$, see (4). \square

Definition 1. A family of simplicial partitions is called *regular* if Condition 1 or 2 or 3 or 4 holds.

Remark 1. Consider a tetrahedron whose base is an equilateral triangle with a very high altitude (ending at the centre of the base). Then the three dihedral angles at the base are almost 90° and the remaining three angles are approximately 60° . However, some solid angles are very small. Therefore, in Condition 4 a positive lower bound on solid angles is prescribed.

Remark 2. Condition (1) seems to be simpler than the ball conditions or the angle condition, and therefore, it may be preferred in theoretical finite element analysis. On the other hand, the angle conditions are often used in finite element codes to keep simplices nondegenerating.

4 Final remarks

In 1957, Synge [18, p. 211] proved that linear triangular elements have optimal interpolation properties in the C^1 -norm provided there exists a positive

constant $\gamma_0 < \pi$ such that for any $\mathcal{T}_h \in \mathcal{F}$ and any triangle $S \in \mathcal{T}_h$ we have

$$\gamma_S \leq \gamma_0, \quad (23)$$

where γ_S is the maximal angle of S . We observe that in this case the minimal angles α_S may tend to zero as $h \rightarrow 0$. On the other hand, if Zlamal's condition holds, then the maximal angle condition (23) holds as well. In 1974, several authors [1, 2, 7, 8] independently proved the convergence of the finite element method under Synge's condition. According to [11], the maximal angle condition (23) is equivalent to the following circumscribed ball condition for $d = 2$: there exists a constant $C_{10} > 0$ such that for any partition $\mathcal{T}_h \in \mathcal{F}$ and any triangle $S \in \mathcal{T}_h$ we have (cf. (17))

$$r^S \leq C_{10} h_S. \quad (24)$$

The associated families of partitions are called *semiregular*. Each regular family is semiregular, but the converse implication does not hold. Therefore, (3) implies (24), but (24) does not imply (3).

Synge's condition (23) is generalized to the case of tetrahedra in [12] and [15]. However, an extension of (23) or (24) to \mathbf{R}^d so that simplicial finite elements preserve their optimal interpolation properties in Sobolev norms is still an open problem.

It is easy to verify that conditions (1) and (2) are equivalent also for non-simplicial finite elements.

Replacing (1) and (2) by

$$\text{meas}_d S \geq C'_1 h^d, \quad (25)$$

and

$$r_S \geq C'_2 h,$$

respectively, we can show that these conditions are also equivalent. The associated family of such partitions is called *strongly regular* (cf. [5, p. 147]).

Remark 3. Let us point out that for a strongly regular family of partitions the well-known inverse inequalities hold (see [5, p. 142]), e.g.,

$$\|v_h\|_1 \leq \frac{C}{h} \|v_h\|_0 \quad \forall v_h \in V_h, \quad (26)$$

where V_h are finite element subspaces of the Sobolev space $H^1(\Omega)$, the symbol C stands for a constant independent of h and $\|\cdot\|_k$ is the standard Sobolev norm. Inverse inequalities play an important role in proving convergence of the finite element method of various problems (see [15]).

Remark 4. In [3], [9], and [10] we show how to generate partitions with nonobtuse dihedral angles, i.e., all simplices satisfy the maximal angle condition in \mathbf{R}^d . Such partitions guarantee the discrete maximum principle for a class of nonlinear elliptic problems solved by linear simplicial elements (see [13]).

References

- [1] BABUŠKA, I., AZIZ, A. K., On the angle condition in the finite element method, *SIAM J. Numer. Anal.* 13 (1976), 214–226.
- [2] BARNHILL, R. E., GREGORY, J. A., Sard kernel theorems on triangular domains with application to finite element error bounds, *Numer. Math.* 25 (1975/1976), 215–229.
- [3] BRANDTS, J., KOROTOV, S., KRÍŽEK, M., Dissection of the path-simplex in R^n into n path-subsimplices, *Preprint A496 (2006), Helsinki University of Technology* (submitted).
- [4] BRANDTS, J., KRÍŽEK, M., Gradient superconvergence on uniform simplicial partitions of polytopes, *IMA J. Numer. Anal.* 23 (2003), 489–505.
- [5] CIARLET, P. G., *The Finite Element Method for Elliptic Problems*. North-Holland, Amsterdam, 1978.
- [6] FIEDLER, M., Geometrie simplexu v E_n , *Časopis Pěst. Mat.* XII (1954), 297–320.
- [7] GREGORY, J. A., Error bounds for linear interpolation on triangles, In *Proc. MAFELAP II* J.R. Whiteman (ed.) Academic Oress, London (1976), 163–170.
- [8] JAMET, P., Estimations de l’erreur pour des éléments finis droits presque dégénérés, *RAIRO Anal. Numér.* 10 (1976), 43–60.
- [9] KOROTOV, S., KRÍŽEK, M., Acute type refinements of tetrahedral partitions of polyhedral domains, *SIAM J. Numer. Anal.* 39 (2001), 724–733.
- [10] KOROTOV, S., KRÍŽEK, M., Global and local refinement techniques yielding nonobtuse tetrahedral partitions, *Comput. Math. Appl.* 50 (2005), 1105–1113.
- [11] KRÍŽEK, M., On semiregular families of triangulations and linear interpolation, *Appl. Math.* 36 (1991), 223–232.
- [12] KRÍŽEK, M., On the maximum angle condition for linear tetrahedral elements, *SIAM J. Numer. Anal.* 29 (1992), 513–520.
- [13] KRÍŽEK, M., LIN QUN, On diagonal dominance of stiffness matrices in 3D, *East-West J. Numer. Math.* 3 (1995), 59–69.
- [14] KRÍŽEK, M., STROUBOULIS, T., How to generate local refinements of unstructured tetrahedral meshes satisfying a regularity ball condition, *Numer. Methods Partial Differential Equations* 13 (1997), 201–214.

- [15] KRÍŽEK, M., NEITTAANMÄKI, P., *Mathematical and Numerical Methods in Modelling in Electrical Engineering: Theory and Applications*, Kluwer Academic Publishers, 1996.
- [16] LIN, J., LIN QUN, Global superconvergence of the mixed finite element methods for 2-d Maxwell equations, *J. Comput. Math.* 21 (2003), 637–646.
- [17] LIN QUN, XU, J., Linear finite elements with high accuracy, *J. Comput. Math.* 3 (1985), 115–133.
- [18] SYNGE, J. L., *The Hypercircle in Mathematical Physics*, Cambridge Univ. Press, Cambridge, 1957.
- [19] ZLÁMAL, M., On the finite element method, *Numer. Math.* 12 (1968), 394–409.

(continued from the back cover)

- A501 Marina Sirviö
On an inverse subordinator storage
June 2006
- A500 Outi Elina Maasalo , Anna Zatorska-Goldstein
Stability of quasiminimizers of the p -Dirichlet integral with varying p on metric spaces
April 2006
- A499 Mikko Parviainen
Global higher integrability for parabolic quasiminimizers in nonsmooth domains
April 2005
- A498 Marcus Ruter , Sergey Korotov , Christian Steenbock
Goal-oriented Error Estimates based on Different FE-Spaces for the Primal and the Dual Problem with Applications to Fracture Mechanics
March 2006
- A497 Outi Elina Maasalo
Gehring Lemma in Metric Spaces
March 2006
- A496 Jan Brandts , Sergey Korotov , Michal Krizek
Dissection of the path-simplex in \mathbf{R}^n into n path-subsimplices
March 2006
- A495 Sergey Korotov
A posteriori error estimation for linear elliptic problems with mixed boundary conditions
March 2006
- A494 Antti Hannukainen , Sergey Korotov
Computational Technologies for Reliable Control of Global and Local Errors for Linear Elliptic Type Boundary Value Problems
February 2006
- A493 Giovanni Formica , Stefania Fortino , Mikko Lyly
A *vartheta* method-based numerical simulation of crack growth in linear elastic fracture
February 2006

HELSINKI UNIVERSITY OF TECHNOLOGY INSTITUTE OF MATHEMATICS
RESEARCH REPORTS

The list of reports is continued inside. Electronical versions of the reports are available at <http://www.math.hut.fi/reports/> .

- A506 Sergey Korotov
Error control in terms of linear functionals based on gradient averaging techniques
July 2006
- A505 Jan Brandts , Sergey Korotov , Michal Krizek
On the equivalence of regularity criteria for triangular and tetrahedral finite element partitions
July 2006
- A504 Janos Karatson , Sergey Korotov , Michal Krizek
On discrete maximum principles for nonlinear elliptic problems
July 2006
- A503 Jan Brandts , Sergey Korotov , Michal Krizek , Jakub Solc
On acute and nonobtuse simplicial partitions
July 2006
- A502 Vladimir M. Miklyukov , Antti Rasila , Matti Vuorinen
Three spheres theorem for p -harmonic functions
June 2006

ISBN 951-22-8315-8

ISSN 0784-3143