

JUHA KINNUNEN

Measure and Integral

Contents

1 MEASURE THEORY	1
1.1 Outer measures	1
1.2 Measurable sets	6
1.3 Measures	13
1.4 The distance function	19
1.5 Characterizations of measurable sets	22
1.6 Metric outer measures	33
1.7 Lebesgue measure revisited	37
1.8 Invariance properties of the Lebesgue measure	50
1.9 Lebesgue measurable sets	52
1.10 A nonmeasurable set	57
1.11 The Cantor set	61
2 MEASURABLE FUNCTIONS	65
2.1 Calculus with infinities	65
2.2 Measurable functions	66
2.3 Cantor-Lebesgue function	72
2.4 Lipschitz mappings on \mathbb{R}^n	76
2.5 Limits of measurable functions	79
2.6 Almost everywhere	82
2.7 Approximation by simple functions	84
2.8 Modes of convergence	88
2.9 Egoroff's and Lusin's theorems	94
3 INTEGRATION	101
3.1 Integral of a nonnegative simple function	101
3.2 Integral of a nonnegative measurable function	104
3.3 Monotone convergence theorem	107
3.4 Fatou's lemma	112
3.5 Integral of a signed function	113
3.6 Dominated convergence theorem	116
3.7 Lebesgue integral	123
3.8 Cavalieri's principle	135

3.9 Lebesgue and Riemann	140
3.10 Fubini's theorem	144
3.11 Fubini's theorem for Lebesgue measure	149

All sets can be measured by an outer measure, which is monotone and countably subadditive function. Most important example is the Lebesgue outer measure, which generalizes the concept of volume to all sets. An outer measure has a proper measure theory on measurable sets. A set is Lebesgue measurable if it is almost a Borel set. Existence of a nonmeasurable set for the Lebesgue outer measure is shown by the axiom of choice.

1

Measure theory

1.1 Outer measures

We begin with a general definition of outer measure. Let X be a nonempty set and consider a mapping on the collection of subsets of X . Recall that if $A_i \subset X$ for $i = 1, 2, \dots$, then

$$\bigcup_{i=1}^{\infty} A_i = \{x \in X : x \in A_i \text{ for some } i\}$$

and

$$\bigcap_{i=1}^{\infty} A_i = \{x \in X : x \in A_i \text{ for every } i\}.$$

Moreover $X \setminus A = \{x \in X : x \notin A\}$.

Definition 1.1. A mapping $\mu^* : \{A : A \subset X\} \rightarrow [0, \infty]$ is an outer measure on X , if

- (1) $\mu^*(\emptyset) = 0$,
- (2) (monotonicity) $\mu^*(A) \leq \mu^*(B)$ whenever $A \subset B \subset X$ and
- (3) (countable subadditivity) $\mu^*(\bigcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \mu^*(A_i)$ whenever $A_i \subset X$, $i = 1, 2, \dots$.

T H E M O R A L : The sum of outer measures of countable many sets that cover a given set is at least the outer measure of the set, that is, volume may get smaller but not larger in countable coverings.

Remark 1.2. Countable subadditivity implies finite subadditivity, since we may add countably many empty sets to a finite collection of sets.

W A R N I N G : It may happen that the equality $\mu^*(A \cup B) = \mu^*(A) + \mu^*(B)$ fails for A and B with $A \cap B = \emptyset$. This means that an outer measure is not necessarily additive on pairwise disjoint sets. Note, however, that the inequality $\mu^*(A \cup B) \leq \mu^*(A) + \mu^*(B)$ always holds by subadditivity.




Figure 1.1: Countable subadditivity.

Example 1.3. Let $X = \{1, 2, 3\}$ and define μ^* by setting $\mu^*(\emptyset) = 0$, $\mu^*(X) = 2$ and $\mu^*(E) = 1$ for all other $E \subset X$. Then μ^* is an outer measure on X . However, if $A = \{1\}$ and $B = \{2\}$, then $A \cap B = \emptyset$ and

$$\mu^*(A \cup B) = \mu^*(\{1, 2\}) = 1 \neq 2 = \mu^*(A) + \mu^*(B).$$

Examples 1.4:

- (1) (The trivial measure) Let $\mu^*(A) = 0$ for every $A \subset X$. Then μ^* is an outer measure. The trivial measure is relatively useless, since all sets have measure zero.
- (2) (The discrete measure) Let

$$\mu^*(A) = \begin{cases} 1, & A \neq \emptyset, \\ 0, & A = \emptyset. \end{cases}$$

The discrete outer measure tells whether or not a set is empty.

- (3) (The Dirac measure) Let $x_0 \in X$ be a fixed point and let

$$\mu^*(A) = \begin{cases} 1, & x_0 \in A, \\ 0, & x_0 \notin A. \end{cases}$$

This is called the Dirac outer measure, or Dirac's delta, at x_0 . The Dirac measure tells whether or not a set contains the point x_0 .

(4) (The counting measure) Let $\mu^*(A)$ be the (possibly infinite) number of points in A . The counting outer measure tells the number of points of a set.

(5) (The Lebesgue measure) Let $X = \mathbb{R}^n$ and consider the n -dimensional closed interval (rectangle)

$$I = \{x \in \mathbb{R}^n : -\infty < a_i \leq x_i \leq b_i < \infty, i = 1, \dots, n\} = [a_1, b_1] \times \dots \times [a_n, b_n]$$

with sides parallel to the coordinate axes. We allow intervals to be degenerate, that is, $b_i = a_i$ for some i . The geometric volume of I is

$$\text{vol}(I) = (b_1 - a_1)(b_2 - a_2) \cdots (b_n - a_n).$$

The Lebesgue outer measure of a set $A \subset \mathbb{R}^n$ is defined as

$$m^*(A) = \inf \left\{ \sum_{i=1}^{\infty} \text{vol}(I_i) : A \subset \bigcup_{i=1}^{\infty} I_i \right\},$$

where the infimum is taken over all coverings of A by countably many intervals I_i , $i = 1, 2, \dots$.

T H E M O R A L : The Lebesgue outer measure of a set is the infimum of sums of volumes of countably many intervals that cover the set.

Observe that this includes coverings with a finite number of intervals, since we may add countably many intervals $I_i = \{x_i\}$ containing only one point with $\text{vol}(I_i) = 0$. The Lebesgue outer measure is nonnegative but may be infinite, that is, $0 \leq m^*(A) \leq \infty$.

- If $m^*(A) < \infty$, by the definition of infimum,

$$m^*(A) \leq \sum_{i=1}^{\infty} \text{vol}(I_i)$$

for every covering of A by countably many intervals I_i , $i = 1, 2, \dots$, and for every $\varepsilon > 0$, there exist intervals I_i , $i = 1, 2, \dots$, such that $A \subset \bigcup_{i=1}^{\infty} I_i$ and

$$\sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(A) + \varepsilon.$$

- $m^*(A) < \infty$, if there exist intervals I_i , $i = 1, 2, \dots$, such that $A \subset \bigcup_{i=1}^{\infty} I_i$ and $\sum_{i=1}^{\infty} \text{vol}(I_i) < \infty$.
- $m^*(A) = \infty$, if $\sum_{i=1}^{\infty} \text{vol}(I_i) = \infty$ for every covering of A by countably many intervals I_i , $i = 1, 2, \dots$.
- $m^*(A) = 0$, if for every $\varepsilon > 0$, there exist intervals I_i , $i = 1, 2, \dots$, such that $A \subset \bigcup_{i=1}^{\infty} I_i$ and $\sum_{i=1}^{\infty} \text{vol}(I_i) < \varepsilon$.




Figure 1.2: Covering by intervals.

We shall discuss more about the Lebesgue outer measure in Section 1.7, Section 1.8 and Section 1.9, but it generalizes the notion of n -dimensional volume to arbitrary subsets of \mathbb{R}^n .

Claim: m^* is an outer measure.

Reason. (1) Let $\varepsilon > 0$. Since

$$\emptyset \subset \left[-\frac{\varepsilon^{\frac{1}{n}}}{2}, \frac{\varepsilon^{\frac{1}{n}}}{2}\right] \times \cdots \times \left[-\frac{\varepsilon^{\frac{1}{n}}}{2}, \frac{\varepsilon^{\frac{1}{n}}}{2}\right] = \left[-\frac{\varepsilon^{\frac{1}{n}}}{2}, \frac{\varepsilon^{\frac{1}{n}}}{2}\right]^n,$$

we have

$$0 \leq m^*(\emptyset) \leq \text{vol}\left(\left[-\frac{\varepsilon^{\frac{1}{n}}}{2}, \frac{\varepsilon^{\frac{1}{n}}}{2}\right]^n\right) = \left(2 \frac{\varepsilon^{\frac{1}{n}}}{2}\right)^n = \varepsilon.$$

By letting $\varepsilon \rightarrow 0$, we conclude $m^*(\emptyset) = 0$. We could also cover \emptyset by the degenerate interval $[x_1, x_1] \times \cdots \times [x_n, x_n]$ for any $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and conclude the claim from this.

(2) Let $A \subset B$. We may assume that $m^*(B) < \infty$, for otherwise $m^*(A) \leq m^*(B) = \infty$ and there is nothing to prove. For every $\varepsilon > 0$ there exist intervals I_i , $i = 1, 2, \dots$, such that $B \subset \bigcup_{i=1}^{\infty} I_i$ and

$$\sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(B) + \varepsilon.$$

Since $A \subset B \subset \bigcup_{i=1}^{\infty} I_i$, we have

$$m^*(A) \leq \sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(B) + \varepsilon.$$

By letting $\varepsilon \rightarrow 0$, we conclude that $m^*(A) \leq m^*(B)$.

(3) Let $A_i \subset \mathbb{R}^n$, $i = 1, 2, \dots$. We may assume that $m^*(A_i) < \infty$ for every $i = 1, 2, \dots$, for otherwise $\sum_{i=1}^{\infty} m^*(A_i) = \infty$ and there is nothing to prove. Let $\varepsilon > 0$. For every $i = 1, 2, \dots$ there exist intervals $I_{i,j}$, $j = 1, 2, \dots$, such that $A_i \subset \bigcup_{j=1}^{\infty} I_{i,j}$ and

$$\sum_{j=1}^{\infty} \text{vol}(I_{i,j}) < m^*(A_i) + \frac{\varepsilon}{2^i}.$$

Then $\bigcup_{i=1}^{\infty} A_i \subset \bigcup_{i=1}^{\infty} \bigcup_{j=1}^{\infty} I_{i,j} = \bigcup_{i,j=1}^{\infty} I_{i,j}$ and

$$\begin{aligned} m^*\left(\bigcup_{i=1}^{\infty} A_i\right) &\leq \sum_{i,j=1}^{\infty} \text{vol}(I_{i,j}) = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \text{vol}(I_{i,j}) \\ &\leq \sum_{i=1}^{\infty} \left(m^*(A_i) + \frac{\varepsilon}{2^i}\right) = \sum_{i=1}^{\infty} m^*(A_i) + \varepsilon. \end{aligned}$$

The claim follows by letting $\varepsilon \rightarrow 0$. ■

(6) (The Hausdorff measure) Let $X = \mathbb{R}^n$, $0 < s < \infty$ and $0 < \delta \leq \infty$. Define

$$\mathcal{H}_{\delta}^s(A) = \inf \left\{ \sum_{i=1}^{\infty} (\text{diam}(B_i))^s : A \subset \bigcup_{i=1}^{\infty} B_i, \text{diam}(B_i) \leq \delta \right\}$$

and

$$\mathcal{H}^s(A) = \lim_{\delta \rightarrow 0} \mathcal{H}_{\delta}^s(A) = \sup_{\delta > 0} \mathcal{H}_{\delta}^s(A).$$

We call \mathcal{H}^s the s -dimensional Hausdorff outer measure on \mathbb{R}^n . This generalizes the notion of s -dimensional measure to arbitrary subsets of \mathbb{R}^n . See Remark 1.25 for the definition of the diameter of a set. We refer to [2, Section 3.8], [4, Chapter 2], [7, Chapter 19], [10, Chapter 7] and [16, Chapter 7] for more.

Observe that, for every $\delta > 0$, an arbitrary set $A \subset \mathbb{R}^n$ can be covered by $B(x, \frac{\delta}{2})$ with $x \in A$, that is,

$$A \subset \bigcup \left\{ B(x, \frac{\delta}{2}) : x \in A \right\},$$

where $B(x, r) = \{y \in \mathbb{R}^n : |y - x| < r\}$ denotes an open ball of radius $r > 0$ and center x . By Lindelöf's theorem every open covering in \mathbb{R}^n has a countable subcovering. This implies that there exist countably many balls $B_i = B(x_i, \frac{\delta}{2})$, $i = 1, 2, \dots$, such that $A \subset \bigcup_{i=1}^{\infty} B_i$. Moreover, we have $\text{diam}(B_i) \leq \delta$ for every $i = 1, 2, \dots$. This shows that the coverings in the definition of the Hausdorff outer measure exist.

(7) Let \mathcal{F} be a collection of subsets of X such that $\emptyset \in \mathcal{F}$ and there exist $A_i \in \mathcal{F}$, $i = 1, 2, \dots$, such that $X = \bigcup_{i=1}^{\infty} A_i$. Let $\rho : \mathcal{F} \rightarrow [0, \infty]$ be any function for which $\rho(\emptyset) = 0$. Then $\mu^* : \{A : A \subset X\} \rightarrow [0, \infty]$,

$$\mu^*(A) = \inf \left\{ \sum_{i=1}^{\infty} \rho(A_i) : A_i \in \mathcal{F}, A \subset \bigcup_{i=1}^{\infty} A_i \right\}$$

is an outer measure on X . Moreover, if ρ is monotone and countably subadditive on \mathcal{F} , then $\mu^* = \rho$ on \mathcal{F} (exercise). This gives a very general method to construct outer measures, see [2, Section 2.8].

(8) (Carathéodory's construction) Let $X = \mathbb{R}^n$, \mathcal{F} be a collection of subsets of X and $\rho : \mathcal{F} \rightarrow [0, \infty]$ be a function. We make the following assumptions on \mathcal{F} and ρ .

- For every $\delta > 0$ there exist $A_i \in \mathcal{F}$, $i = 1, 2, \dots$, such that $X = \bigcup_{i=1}^{\infty} A_i$ and $\text{diam}(A_i) \leq \delta$.
- For every $\delta > 0$ there exists $A \in \mathcal{F}$ such that $\rho(A) \leq \delta$ and $\text{diam}(A) \leq \delta$.

For $0 < \delta \leq \infty$ and $A \subset X$, we let

$$\mu_{\delta}^*(A) = \inf \left\{ \sum_{i=1}^{\infty} \rho(A_i) : A_i \in \mathcal{F}, A \subset \bigcup_{i=1}^{\infty} A_i, \text{diam}(A_i) \leq \delta \right\}.$$

The first assumption guarantees that we can cover any set A with sets in \mathcal{F} of arbitrarily small measure and the second assumption implies that $\mu_{\delta}^*(\emptyset) = 0$. It can be shown that μ_{δ}^* is an outer measure (exercise), but it is usually not additive on disjoint sets (see Theorem 1.12) and not a Borel outer measure (see Definition 1.37). Clearly,

$$\mu_{\delta'}^*(A) \leq \mu_{\delta}^*(A) \quad \text{for } 0 < \delta < \delta' \leq \infty.$$

Thus we may define

$$\mu^*(A) = \lim_{\delta \rightarrow 0} \mu_{\delta}^*(A) = \sup_{\delta > 0} \mu_{\delta}^*(A).$$

The outer measure μ^* has much better properties than μ_{δ}^* . For example, it is always a Borel outer measure (see Theorem 1.53 and Remarks 1.54). Moreover, if the members of \mathcal{F} are Borel sets, then μ^* is Borel regular (see Definition 1.37). This gives a very general method to construct Borel outer measures, see [2, Section 3.3].

T H E M O R A L : The examples above show that it is easy to construct outer measures. However, we have to restrict ourselves to a class of measurable sets in order to obtain a useful theory.

1.2 Measurable sets

We discuss so-called Carathéodory criterion for measurability for a general outer measure. The definition is perhaps not very intuitive, but it will be useful in the arguments. In practice it may be difficult to show directly from the definition that a set is measurable. However, Lemma 1.63 proves that every closed interval is Lebesgue measurable by using the definition below. Later we give a more

geometric characterizations of measurable sets for the Lebesgue and other outer measures.

Definition 1.5. A set $A \subset X$ is μ^* -measurable, if

$$\mu^*(E) = \mu^*(E \cap A) + \mu^*(E \setminus A)$$

for every $E \subset X$.

T H E M O R A L : A measurable set divides every set into two disjoint parts in an additive way. On the other hand, a nonmeasurable set divides some set into two parts in a nonadditive way.




Figure 1.3: A measurable set.

Remarks 1.6:

- (1) Since $E = (E \cap A) \cup (E \setminus A)$, by subadditivity

$$\mu^*(E) \leq \mu^*(E \cap A) + \mu^*(E \setminus A).$$

This means that an inequality always holds in Definition 1.5. To show that a set A is μ^* -measurable it is enough to prove that

$$\mu^*(E) \geq \mu^*(E \cap A) + \mu^*(E \setminus A).$$

for every $E \subset X$. Note that this certainly holds if $\mu^*(E) = \infty$, so that it is enough to consider sets E with $\mu^*(E) < \infty$. A set A is not μ^* -measurable, if $\mu^*(E) < \mu^*(E \cap A) + \mu^*(E \setminus A)$ for some $E \subset X$.

(2) If A is μ^* -measurable and B is an arbitrary subset of X with $A \cap B = \emptyset$, then by taking $E = A \cup B$ in Definition 1.5 we have

$$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B).$$

In particular, an outer measure behaves additively on two disjoint measurable subsets. A repeated application of this result implies that an outer measure is additive finitely many pairwise disjoint measurable sets. Theorem 1.12 below shows that this holds true also for countably many sets.

On the other hand, if there exist sets A and B such that $A \cap B = \emptyset$ and

$$\mu^*(A \cup B) \neq \mu^*(A) + \mu^*(B),$$

that is $\mu^*(A \cup B) < \mu^*(A) + \mu^*(B)$, then A and B are not μ^* -measurable.

Reason. Let $E = A \cup B$. Then

$$\mu^*(E) = \mu^*(A \cup B) \neq \mu^*(A) + \mu^*(B) = \mu^*(E \cap A) + \mu^*(E \setminus A)$$

and

$$\mu^*(E) = \mu^*(A \cup B) \neq \mu^*(A) + \mu^*(B) = \mu^*(E \setminus B) + \mu^*(E \cap B). \blacksquare$$

(3) If A is μ^* -measurable and $A \subset B$, where B is an arbitrary subset of X , by taking $E = B$ in Definition 1.5, we have

$$\mu^*(B) = \mu^*(\underbrace{B \cap A}_{=A}) + \mu^*(B \setminus A) = \mu^*(A) + \mu^*(B \setminus A).$$

(4) If $\mu^*(A) = 0$, then A is μ^* -measurable. In other words, sets of measure zero are always measurable.

Reason. Since $E \cap A \subset A$ and $E \setminus A \subset E$, we have

$$\mu^*(E \cap A) + \mu^*(E \setminus A) \leq \underbrace{\mu^*(A)}_{=0} + \mu^*(E) = \mu^*(E)$$

for every $E \subset X$. On the other hand, by (1) we always have the inequality in the reverse direction. This implies A is μ^* -measurable. \blacksquare

(5) \emptyset and X are μ^* -measurable. In other words, the empty set and the entire space are always measurable.

Reason.

$$\mu^*(E) = \mu^*(\underbrace{E \cap \emptyset}_{=\emptyset}) + \mu^*(\underbrace{E \setminus \emptyset}_{=E})$$

and

$$\mu^*(E) = \mu^*(\underbrace{E \cap X}_{=E}) + \mu^*(\underbrace{E \setminus X}_{=\emptyset}),$$

for every $E \subset X$. This shows that \emptyset and X are μ^* -measurable. \blacksquare

Another way is to apply Lemma 1.11 below, which asserts that A is μ^* -measurable if and only if $X \setminus A$ is μ^* -measurable. Hence $X = X \setminus \emptyset$ is μ^* -measurable, since \emptyset is μ^* -measurable as a set of measure zero.

- (6) The only measurable sets for the discrete measure are \emptyset and X (exercise).
In this case there are extremely few measurable sets.
- (7) All sets are measurable for the Dirac measure (exercise). In this case there are extremely many measurable sets.

Example 1.7. (Continuation of Example 1.3) Let $X = \{1, 2, 3\}$ and, define an outer measure μ^* such that $\mu^*(\emptyset) = 0$, $\mu^*(X) = 2$ and $\mu^*(E) = 1$ for all other $E \subset X$. If $a, b \in X$ are different points, $A = \{a\}$ and $E = \{a, b\}$, then

$$\mu^*(E) = \mu^*(\{a, b\}) = 1 < 2 = \mu^*(\{a\}) + \mu^*(\{b\}) = \mu^*(E \cap A) + \mu^*(E \setminus A).$$

This means that A is not μ^* -measurable. In the same way we can see that all sets consisting of two points are not μ^* -measurable. Thus only μ^* -measurable sets are \emptyset and X .

Next we discuss structural properties of measurable sets.

Definition 1.8. A collection \mathcal{A} of subsets of X is a σ -algebra, if

- (1) $\emptyset \in \mathcal{A}$,
- (2) $A \in \mathcal{A}$ implies $A^c = X \setminus A \in \mathcal{A}$ and
- (3) $A_i \in \mathcal{A}$ for every $i = 1, 2, \dots$ implies $\bigcup_{i=1}^{\infty} A_i \in \mathcal{A}$.

T H E M O R A L : A σ -algebra is closed under complements and countable unions.

Remark 1.9. Every σ -algebra \mathcal{A} has the following properties.

- (1) Since $X \setminus \emptyset = X$, by (1) and (2) in Definition 1.8 we have $X \in \mathcal{A}$.
- (2) If $A_1, \dots, A_k \in \mathcal{A}$, then $\bigcup_{i=1}^k A_i \in \mathcal{A}$. This follows from (3) in Definition 1.8 by taking $A_i = \emptyset$ for $i = k+1, k+2, \dots$
- (3) If $A_i \in \mathcal{A}$ for every $i = 1, 2, \dots$, then $\bigcap_{i=1}^{\infty} A_i \in \mathcal{A}$.

Reason. By de Morgan's law and Definition 1.8 (3), we have

$$X \setminus \bigcap_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} (X \setminus A_i) \in \mathcal{A}.$$

Thus by Definition 1.8 (2) we conclude that

$$\bigcap_{i=1}^{\infty} A_i = X \setminus \left(X \setminus \bigcap_{i=1}^{\infty} A_i \right) \in \mathcal{A}. \quad \blacksquare$$

The corresponding claim also holds for collections of finitely many sets. By taking $A_i = \emptyset$ for $i = k+1, k+2, \dots$, we have $\bigcap_{i=1}^k A_i \in \mathcal{A}$.

(4) If $A, B \in \mathcal{A}$, by (2) in Definition 1.8 and remark (2) above, we have $A \setminus B = A \cap (X \setminus B) \in \mathcal{A}$.

Examples 1.10:

- (1) Let \mathcal{A} be the collection of all subsets of X . Then \mathcal{A} is a σ -algebra. This is the largest σ -algebra of subsets of X .
- (2) Let $\mathcal{A} = \{\emptyset, X\}$. Then \mathcal{A} is a σ -algebra. This is the smallest σ -algebra of subsets of X .
- (3) Let X be an infinite set and let \mathcal{A} be the collection of all finite subsets of X . Then \mathcal{A} does not contain X and \mathcal{A} is not closed under complementation, so that it is not a σ -algebra.
- (4) Let X be an infinite set and let \mathcal{A} be the collection of subsets of X such that either A or $X \setminus A$ is finite. Then \mathcal{A} is not closed under countable unions, so that it is not a σ -algebra.

Lemma 1.11. The collection \mathcal{M} of μ^* -measurable sets is a σ -algebra.

T H E M O R A L : All sets constructed of measurable sets by countably many set theoretic operations of taking complements, unions and intersections are measurable.

Proof. (1) $\mu^*(\emptyset) = 0$ implies that $\emptyset \in \mathcal{M}$, see Remark 1.6.

(2) $\mu^*(E) = \mu^*(E \cap A) + \mu^*(E \setminus A) = \mu^*(E \setminus (X \setminus A)) + \mu^*(E \cap (X \setminus A))$ for every $E \subset X$. This shows that $A^c \in \mathcal{M}$.

(3) **Step 1:** First we show that $A_1, A_2 \in \mathcal{M}$ implies $A_1 \cup A_2 \in \mathcal{M}$.

$$\begin{aligned}
 \mu^*(E) &= \mu^*(E \setminus A_1) + \mu^*(E \cap A_1) \quad (A_1 \in \mathcal{M}, E \text{ test set}) \\
 &= \mu^*((E \setminus A_1) \cap A_2) + \mu^*((E \setminus A_1) \setminus A_2) + \mu^*(E \cap A_1) \\
 &\quad (A_2 \in \mathcal{M}, \text{ apply } E \setminus A_1 \text{ as a test set}) \\
 &= \mu^*((E \setminus A_1) \cap A_2) + \mu^*(E \setminus (A_1 \cup A_2)) + \mu^*(E \cap A_1) \\
 &\quad ((E \setminus A_1) \setminus A_2 = E \setminus (A_1 \cup A_2)) \\
 &\geq \mu^*((E \setminus A_1) \cap A_2) + \mu^*(E \setminus (A_1 \cup A_2)) \quad (\text{subadditivity}) \\
 &= \mu^*(E \cap (A_1 \cup A_2)) + \mu^*(E \setminus (A_1 \cup A_2)) \\
 &\quad ((E \setminus A_1) \cap A_2) \cup (E \cap A_1) = E \cap (A_1 \cup A_2)
 \end{aligned}$$

for every $E \subset X$. Recursively, the same result holds for finitely many sets: If $A_i \in \mathcal{M}, i = 1, 2, \dots, k$, then $\bigcup_{i=1}^k A_i \in \mathcal{M}$. By de Morgan's law, we also have $\bigcap_{i=1}^k A_i \in \mathcal{M}$.

Step 2: We construct pairwise disjoint sets C_i such that $C_i \subset A_i$ and $\bigcup_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} C_i$. Let $B_k = \bigcup_{i=1}^k A_i$, $k = 1, 2, \dots$. Then $B_k \subset B_{k+1}$ and

$$\bigcup_{i=1}^{\infty} A_i = B_1 \cup \left(\bigcup_{k=1}^{\infty} (B_{k+1} \setminus B_k) \right).$$

Let

$$C_1 = B_1 \quad \text{and} \quad C_{i+1} = B_{i+1} \setminus B_i, \quad i = 1, 2, \dots$$

Then $C_i \cap C_j = \emptyset$ whenever $i \neq j$ and the sets C_i , $i = 1, 2, \dots$, have the required properties. The sets $C_i \in \mathcal{M}$, since they are finite unions and intersections μ^* -measurable sets, see Step 1.

Note:

$$\begin{aligned} C_{i+1} &= B_{i+1} \setminus B_i = A_{i+1} \setminus \bigcup_{j=1}^i A_i = \bigcap_{j=1}^i (A_{i+1} \setminus A_i) \\ &= \bigcap_{j=1}^i (A_{i+1} \cap (X \setminus A_i)) = A_{i+1} \cap \bigcap_{j=1}^i (X \setminus A_i), \quad i = 1, 2, \dots \end{aligned}$$




Figure 1.4: Covering by disjoint sets.

Step 3: By the argument in Step 2 we may assume that the sets $A_i \in \mathcal{M}$, $i = 1, 2, \dots$, are pairwise disjoint, that is, $A_i \cap A_j = \emptyset$ for $i \neq j$. Let $B_k = \bigcup_{i=1}^k A_i$, $k = 1, 2, \dots$ We show by induction that

$$\mu^*(E \cap B_k) = \sum_{i=1}^k \mu^*(E \cap A_i), \quad k = 1, 2, \dots,$$

for every $E \subset X$.

Note: By choosing $E = X$, this implies finite additivity on pairwise disjoint measurable sets.

The case $k = 1$ is clear. Assume that the claim holds with index k . Then

$$\begin{aligned}
 \mu^*(E \cap B_{k+1}) &= \mu^*((E \cap B_{k+1}) \cap B_k) + \mu^*((E \cap B_{k+1}) \setminus B_k) \\
 &\quad (B_k \in \mathcal{M}, E \cap B_{k+1} \text{ as a test set}) \\
 &= \mu^*(E \cap B_k) + \mu^*(E \cap A_{k+1}) \\
 &\quad (B_k \subset B_{k+1}, A_i \text{ are pairwise disjoint implies } B_{k+1} \setminus B_k = A_{k+1}) \\
 &= \sum_{i=1}^k \mu^*(E \cap A_i) + \mu^*(E \cap A_{k+1}) \quad (\text{the induction assumption}) \\
 &= \sum_{i=1}^{k+1} \mu^*(E \cap A_i).
 \end{aligned}$$

Step 4: By Step 3 and monotonicity with $B_k \subset \bigcup_{i=1}^{\infty} A_i$, we have

$$\sum_{i=1}^k \mu^*(E \cap A_i) = \mu^*(E \cap B_k) \leq \mu^*\left(E \cap \bigcup_{i=1}^{\infty} A_i\right).$$

It follows that

$$\sum_{i=1}^{\infty} \mu^*(E \cap A_i) = \lim_{k \rightarrow \infty} \sum_{i=1}^k \mu^*(E \cap A_i) \leq \mu^*\left(E \cap \bigcup_{i=1}^{\infty} A_i\right).$$

On the other hand, by countable subadditivity

$$\mu^*\left(E \cap \bigcup_{i=1}^{\infty} A_i\right) = \mu^*\left(\bigcup_{i=1}^{\infty} (E \cap A_i)\right) \leq \sum_{i=1}^{\infty} \mu^*(E \cap A_i).$$

This shows that

$$\mu^*\left(E \cap \bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu^*(E \cap A_i)$$

whenever $A_i \in \mathcal{M}$, $i = 1, 2, \dots$, are pairwise disjoint.

Note: By choosing $E = X$, this implies countable additivity on pairwise disjoint measurable sets.

Step 5: Let $E \subset X$ and $A = \bigcup_{i=1}^{\infty} A_i$ with pairwise disjoint $A_i \in \mathcal{M}$, $i = 1, 2, \dots$.

Then

$$\begin{aligned}
 \mu^*(E) &= \mu^*(E \cap B_k) + \mu^*(E \setminus B_k) \quad (B_k \in \mathcal{M} \text{ by Step 1}) \\
 &= \sum_{i=1}^k \mu^*(E \cap A_i) + \mu^*(E \setminus B_k) \quad (\text{Step 3}) \\
 &\geq \sum_{i=1}^k \mu^*(E \cap A_i) + \mu^*(E \setminus A), \quad k = 1, 2, \dots \quad (B_k \subset A)
 \end{aligned}$$

This implies that

$$\begin{aligned}
 \mu^*(E) &\geq \lim_{k \rightarrow \infty} \sum_{i=1}^k \mu^*(E \cap A_i) + \mu^*(E \setminus A) \\
 &= \sum_{i=1}^{\infty} \mu^*(E \cap A_i) + \mu^*(E \setminus A) \\
 &= \mu^*(E \cap A) + \mu^*(E \setminus A). \quad (\text{Step 4})
 \end{aligned}$$

By Remark 1.6 (1) we have $A \in \mathcal{M}$. Note that the countable additivity on pairwise disjoint measurable sets in Step 4 is not really needed in the argument above. We could have used countable subadditivity instead. However, we apply the countable additivity on pairwise disjoint measurable sets in Step 4 in the proof of Theorem 1.12 below. \square

1.3 Measures

From the proof of Lemma 1.11 we see that an outer measure is countably additive on pairwise disjoint measurable sets. This is a very useful property. Example 1.3 shows this does not necessarily hold for sets that are not measurable. The overall idea is that an outer measure produces a proper measure theory when restricted to measurable sets.

Theorem 1.12. (Countable additivity) Assume that $A_i \subset X$, $i = 1, 2, \dots$, are pairwise disjoint ($A_i \cap A_j = \emptyset$ for $i \neq j$) and μ^* -measurable sets. Then

$$\mu^* \left(\bigcup_{i=1}^{\infty} A_i \right) = \sum_{i=1}^{\infty} \mu^*(A_i).$$

T H E M O R A L : An outer measure is countably additive on pairwise disjoint measurable sets. The outer measure is preserved under partitions a given measurable set into countably many pairwise disjoint measurable sets independently how the partition is done. In other words, volume is preserved in such partitions.

Proof. By Step 4 of the the proof of Lemma 1.11, we have

$$\mu^* \left(E \cap \bigcup_{i=1}^{\infty} A_i \right) = \sum_{i=1}^{\infty} \mu^*(E \cap A_i)$$

whenever $A_i \in \mathcal{M}$, $i = 1, 2, \dots$, are pairwise disjoint μ^* -measurable and $E \subset X$. The claim follows by choosing $E = X$. \square

Definition 1.13. Assume that \mathcal{M} is σ -algebra on X . A mapping $\mu: \mathcal{M} \rightarrow [0, \infty]$ is a measure on a measure space (X, \mathcal{M}, μ) , if

- (1) $\mu(\emptyset) = 0$ and
- (2) $\mu \left(\bigcup_{i=1}^{\infty} A_i \right) = \sum_{i=1}^{\infty} \mu(A_i)$ whenever $A_i \in \mathcal{M}$, $i = 1, 2, \dots$, are pairwise disjoint.

T H E M O R A L : A measure is a countably additive set function on pairwise disjoint sets in the σ -algebra. An outer measure is defined on all subsets, but a measure is defined only on sets in the σ -algebra.

Remarks 1.14:

- (1) A measure μ is monotone on \mathcal{M} .




Figure 1.5: Pairwise disjoint sets.

Reason.

$$\mu(B) = \mu((B \setminus A) \cup A) = \mu(A) + \mu(B \setminus A) \geq \mu(A)$$

for every $A, B \in \mathcal{M}$ with $A \subset B$. ■

(2) A measure μ is countably subadditive on \mathcal{M} .

Reason. Let $A_i \in \mathcal{M}$, $i = 1, 2, \dots$. Since \mathcal{M} is a σ -algebra, we have $\bigcup_{i=1}^{\infty} A_i \in \mathcal{M}$. As in the proof of Lemma 1.11 (Step 2), there exist pairwise disjoint sets $C_i \in \mathcal{M}$ such that $C_i \subset A_i$, $i = 1, 2, \dots$, and $\bigcup_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} C_i$. By (1) we have $\mu(C_i) \leq \mu(A_i)$, $i = 1, 2, \dots$, and thus

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \mu\left(\bigcup_{i=1}^{\infty} C_i\right) = \sum_{i=1}^{\infty} \mu(C_i) \leq \sum_{i=1}^{\infty} \mu(A_i). \quad \blacksquare$$

(3) It is possible to develop a theory also for signed or even complex valued measures. Assume that \mathcal{M} is σ -algebra on X . A mapping $\mu: \mathcal{M} \rightarrow [-\infty, \infty]$ is a signed measure on a measure space (X, \mathcal{M}, μ) , if $\mu(\emptyset) = 0$ and whenever $A_i \in \mathcal{M}$, $i = 1, 2, \dots$, are pairwise disjoint sets, then $\sum_{i=1}^{\infty} \mu(A_i)$ exists as an extended real number, that is the sum converges in $[-\infty, \infty]$, and

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu(A_i).$$

Every subset of a set of outer measure zero is measurable as a set of outer measure zero. In contrast, there is a delicate issue related to sets of measure zero for a measure defined on a σ -algebra.

Definition 1.15. A measure μ on a measure space (X, \mathcal{M}, μ) is said to be complete, if $B \in \mathcal{M}$, $\mu(B) = 0$ and $A \subset B$ implies $A \in \mathcal{M}$.

T H E M O R A L : A measure is complete, if every subset of a set of measure zero is measurable. This will be useful when we discuss properties that hold outside sets of measure zero, see Remark 2.31.

Example 1.16. It is possible that $A \subset B \in \mathcal{M}$ and $\mu(B) = 0$, but $A \notin \mathcal{M}$.

Reason. Let $X = \{1, 2, 3\}$ and $\mathcal{M} = \{\emptyset, \{1\}, \{2, 3\}, X\}$. Then \mathcal{M} is a σ -algebra. Define a measure μ on \mathcal{M} by setting $\mu(\emptyset) = 0$, $\mu(\{1\}) = 1$, $\mu(\{2, 3\}) = 0$ and $\mu(X) = 1$. In this case $\{2, 3\} \in \mathcal{M}$ and $\mu^*(\{2, 3\}) = 0$, but $\{2\} \notin \mathcal{M}$. ■

Remarks 1.17:

- (1) The measure space $(\mathbb{R}^n, \mathcal{M}, \mu)$, where \mathcal{M} denotes the collection of Lebesgue measurable sets and μ the Lebesgue outer measure is complete, see Remark 1.6 (4). On the other hand, the measure space $(\mathbb{R}^n, \mathcal{B}, \mu)$, where \mathcal{B} denotes the Borel sets and μ the Lebesgue outer measure is not complete, see Definition 1.35 and discussion in Section 2.3. The reason is that that there exist $B \in \mathcal{B}$ such that $\mu(B) = 0$ and B has a subset which is not a Borel set. However, the σ -algebra of Lebesgue measurable sets is the completion of the Borel σ -algebra (exercise), see Corollary 1.74 and Corollary 1.49.
- (2) Every measure space can be completed by adding sets of measure zero to the σ -algebra in the following way. Assume that (X, \mathcal{M}, μ) is a measure space and let $(X, \overline{\mathcal{M}}, \overline{\mu})$, where

$$\overline{\mathcal{M}} = \{A \cup M : A \in \mathcal{M}, M \subset N \text{ where } N \in \mathcal{M} \text{ satisfies } \mu(N) = 0\}$$

and $\overline{\mu}(A \cup M) = \mu(A)$. Then $(X, \overline{\mathcal{M}}, \overline{\mu})$ is a complete measure space such that $\overline{\mathcal{M}} \supset \mathcal{M}$ and $\overline{\mu}$ is the unique extension of μ to $\overline{\mathcal{M}}$ (exercise). See also [2, Theorem 2.26], [3, Proposition 1.5.1], [10, Exercise 2, p. 312] and [11, Exercise 1.4.6, p. 78].

The following finiteness condition is useful for us later.

Definition 1.18. A measure μ on a measure space (X, \mathcal{M}, μ) is σ -finite, if $X = \bigcup_{i=1}^{\infty} A_i$, where $A_i \in \mathcal{M}$ and $\mu(A_i) < \infty$ for every $i = 1, 2, \dots$

T H E M O R A L : If a measure is σ -finite, the entire space can be covered by measurable sets with finite measure. In many cases it is enough to assume that a measure is σ -finite instead of $\mu(X) < \infty$. By the proof of Lemma 1.11 (Step 2), we may assume that that the covering sets A_i are pairwise disjoint. The corresponding notion can be defined for outer measures as well.

Lemma 1.19. The Lebesgue outer measure m^* is σ -finite.

Proof. Clearly $\mathbb{R}^n = \bigcup_{i=1}^{\infty} B(0, i)$, where $B(0, i) = \{x \in \mathbb{R}^n : |x| < i\}$ is a ball with center at the origin and radius i . The Lebesgue outer measure of the ball $B(0, i)$ is finite, since

$$m^*(B(0, i)) \leq \text{vol}([-i, i]^n) = (2i)^n < \infty.$$

We shall show later that all open sets are Lebesgue measurable, see Lemma 1.55 and Lemma 1.67. \square

Remark 1.20. Every outer measure restricted to measurable sets induces a complete measure. On the other hand, every measure on a measure space (X, \mathcal{M}, μ) induces a regular (see Remark 1.38 (2)) outer measure

$$\mu^*(E) = \inf \left\{ \sum_{i=1}^{\infty} \mu(A_i) : E \subset \bigcup_{i=1}^{\infty} A_i, A_i \in \mathcal{M} \text{ for every } i = 1, 2, \dots \right\}.$$

Then every set in \mathcal{M} is μ^* -measurable and $\mu = \mu^*$ on \mathcal{M} . This means that μ^* is an extension of μ . This is sometimes called the Hahn-Kolmogorov theorem. The extension is unique, if the measure space (X, \mathcal{M}, μ) is σ -finite, see Definition 1.18. If the measure space (X, \mathcal{M}, μ) is complete, see Definition 1.15, then the class of μ^* -measurable sets is precisely \mathcal{M} (exercise). See also [2, p. 99–116], [7, Lemma 9.6], [10, p. 270–273] and [11, p. 153–156].

T H E M O R A L : If the measure is σ -finite and complete, then there will be no new measurable sets when we switch to the induced outer measure. In this sense the difference between outer measures and measures is small.

Next we give some examples of measures.

Examples 1.21:

- (1) (X, \mathcal{M}, μ) , where X is a set, μ is an outer measure on X and \mathcal{M} is the σ -algebra of μ -measurable sets.
- (2) $(\mathbb{R}^n, \mathcal{M}, m^*)$, where m^* is the Lebesgue outer measure and \mathcal{M} is the σ -algebra of Lebesgue measurable sets.
- (3) A measure space (X, \mathcal{M}, μ) with $\mu(X) = 1$ is called a probability space, μ a probability measure and sets belonging to \mathcal{M} events.

The next theorem shows that an outer measure has useful monotone convergence properties on measurable sets.

Theorem 1.22. Assume that μ^* is an outer measure on X and that sets $A_i \subset X$, $i = 1, 2, \dots$, are μ^* -measurable.

- (1) (Upwards monotone convergence) If $A_1 \subset A_2 \subset \dots$, then

$$\lim_{i \rightarrow \infty} \mu^*(A_i) = \mu^* \left(\bigcup_{i=1}^{\infty} A_i \right).$$

(2) (Downwards monotone convergence) If $A_1 \supset A_2 \supset \dots$, and $\mu^*(A_{i_0}) < \infty$ for some i_0 , then

$$\lim_{i \rightarrow \infty} \mu^*(A_i) = \mu^* \left(\bigcap_{i=1}^{\infty} A_i \right).$$

T H E M O R A L : The measure theory is compatible under taking limits, if we approximate a given measurable set with an increasing sequence of measurable sets from inside or a decreasing sequence of measurable sets from outside.




Figure 1.6: Monotone sequences of sets.

Remarks 1.23:

(1) The results do not hold, in general, without the measurability assumptions.

Reason. Let $X = \mathbb{N}$. Define an outer measure on \mathbb{N} by

$$\mu^*(A) = \begin{cases} 0, & A = \emptyset, \\ 1, & A \text{ finite,} \\ 2, & A \text{ infinite.} \end{cases}$$

Let $A_i = \{1, 2, \dots, i\}$, $i = 1, 2, \dots$. Then

$$\mu^* \left(\bigcup_{i=1}^{\infty} A_i \right) = 2 \neq 1 = \lim_{i \rightarrow \infty} \mu^*(A_i).$$

(2) The assumption $\mu^*(A_{i_0}) < \infty$ is essential in (2).

Reason. Let $X = \mathbb{R}$, m^* be the Lebesgue outer measure and $A_i = [i, \infty)$, $i = 1, 2, \dots$. Then $\bigcap_{i=1}^{\infty} A_i = \emptyset$ and $m^*(A_i) = \infty$ for every $i = 1, 2, \dots$. In this case

$$\lim_{i \rightarrow \infty} m^*(A_i) = \infty, \text{ but } m^*\left(\bigcap_{i=1}^{\infty} A_i\right) = m^*(\emptyset) = 0. \quad \blacksquare$$

(3) The following observation, see also Remark 1.6 (3) will be used several times in the proof of Theorem 1.22. Assume that A is μ^* -measurable and let $B \subset \mathbb{R}^n$ be any set with $A \subset B$ and $\mu^*(A) < \infty$. By Definition 1.5, we have

$$\mu^*(B) = \mu^*(B \cap A) + \mu^*(B \setminus A) = \mu^*(A) + \mu^*(B \setminus A)$$

and thus $\mu^*(B \setminus A) = \mu^*(B) - \mu^*(A)$. If both A and B are μ^* -measurable, we can conclude the same result from additivity on disjoint measurable sets as

$$\mu^*(B) = \mu^*(A \cup (B \setminus A)) = \mu^*(A) + \mu^*(B \setminus A).$$

Proof. (1) We may assume that $\mu^*(A_i) < \infty$ for every i , otherwise the claim follows from monotonicity. We write $\bigcup_{i=1}^{\infty} A_i$ as a union of countably many pairwise disjoint measurable sets as

$$\bigcup_{i=1}^{\infty} A_i = A_1 \cup \bigcup_{i=1}^{\infty} (A_{i+1} \setminus A_i).$$

By Theorem 1.12 and Remark 1.23 (3), we have

$$\begin{aligned} \mu^*\left(\bigcup_{i=1}^{\infty} A_i\right) &= \mu^*\left(A_1 \cup \bigcup_{i=1}^{\infty} (A_{i+1} \setminus A_i)\right) \\ &= \mu^*(A_1) + \sum_{i=1}^{\infty} \mu^*(A_{i+1} \setminus A_i) \quad (\text{disjointness and measurability}) \\ &= \mu^*(A_1) + \sum_{i=1}^{\infty} \left(\mu^*(A_{i+1}) - \mu^*(A_i)\right) \quad (\text{Remark 1.23 (3)}) \\ &= \lim_{k \rightarrow \infty} \left(\mu^*(A_1) + \sum_{i=1}^k \left(\mu^*(A_{i+1}) - \mu^*(A_i)\right)\right) \\ &= \lim_{k \rightarrow \infty} \mu^*(A_{k+1}) = \lim_{i \rightarrow \infty} \mu^*(A_i). \end{aligned}$$

(2) By replacing sets A_i by $A_i \cap A_{i_0}$, we may assume that $\mu^*(A_1) < \infty$. $A_{i+1} \subset A_i$ implies $A_1 \setminus A_i \subset A_1 \setminus A_{i+1}$ for every $i = 1, 2, \dots$. By (1) we have

$$\begin{aligned} \mu^*\left(\bigcup_{i=1}^{\infty} (A_1 \setminus A_i)\right) &= \lim_{i \rightarrow \infty} \mu^*(A_1 \setminus A_i) \\ &= \lim_{i \rightarrow \infty} (\mu^*(A_1) - \mu^*(A_i)) \quad (\text{Remark 1.23 (3)}) \\ &= \mu^*(A_1) - \lim_{i \rightarrow \infty} \mu^*(A_i). \end{aligned}$$

On the other hand, by de Morgan's law and Remark 1.23 (3), we have

$$\mu^* \left(\bigcup_{i=1}^{\infty} (A_1 \setminus A_i) \right) = \mu^* \left(A_1 \setminus \bigcap_{i=1}^{\infty} A_i \right) = \mu^*(A_1) - \mu^* \left(\bigcap_{i=1}^{\infty} A_i \right).$$

This implies that

$$\mu^*(A_1) - \mu^* \left(\bigcap_{i=1}^{\infty} A_i \right) = \mu^*(A_1) - \lim_{i \rightarrow \infty} \mu^*(A_i).$$

Since $\mu^*(A_1) < \infty$, we may conclude that

$$\mu^* \left(\bigcap_{i=1}^{\infty} A_i \right) = \lim_{i \rightarrow \infty} \mu^*(A_i). \quad \square$$

1.4 The distance function

The distance function will be a useful tool in the sequel.

Definition 1.24. Let $A \subset \mathbb{R}^n$ with $A \neq \emptyset$. The distance from a point $x \in \mathbb{R}^n$ to A is

$$\text{dist}(x, A) = \inf\{|x - y| : y \in A\}.$$

Remarks 1.25:

(1) The distance between the nonempty sets $A, B \subset \mathbb{R}^n$ is

$$\text{dist}(A, B) = \inf\{|x - y| : x \in A \text{ and } y \in B\}.$$

(2) The diameter of the nonempty set $A \subset \mathbb{R}^n$ is

$$\text{diam}(A) = \sup\{|x - y| : x, y \in A\}.$$

Lemma 1.26. Let $A \subset \mathbb{R}^n$ with $A \neq \emptyset$. For every $x \in \mathbb{R}^n$, there exist a point $x_0 \in \overline{A}$ such that $\text{dist}(x, A) = |x_0 - x|$.

T H E M O R A L : There is a closest point in the closure of the set. If A is closed, then the closest point belongs to A . In general, the closest point is not unique.

Proof. Let $x \in \mathbb{R}^n$. There exists a sequence $y_i \in A$, $i = 1, 2, \dots$, such that

$$\lim_{i \rightarrow \infty} |x - y_i| = \text{dist}(x, A).$$

The sequence (y_i) is bounded and by Bolzano-Weierstrass theorem it has a converging subsequence (y_{j_k}) such that $y_{j_k} \rightarrow x_0$ as $k \rightarrow \infty$ for some $x_0 \in \mathbb{R}^n$. Since \overline{A} is a closed set and $y_{j_k} \in A$ for every k , we have $x_0 \in \overline{A}$. Since $y \mapsto |x - y|$ is a continuous function, we conclude

$$|x - x_0| = \lim_{k \rightarrow \infty} |x - y_{j_k}| = \text{dist}(x, A). \quad \square$$

Lemma 1.27. Let $A \subset \mathbb{R}^n$ with $A \neq \emptyset$. Then

$$|\text{dist}(x, A) - \text{dist}(y, A)| \leq |x - y|$$

for every $x, y \in \mathbb{R}^n$.

T H E M O R A L : The distance function is a Lipschitz continuous function with the Lipschitz constant one. In particular, the distance function does not increase distances between points.

Proof. Let $x, y \in \mathbb{R}^n$. By the triangle inequality $|x - z| \leq |x - y| + |y - z|$ for every $z \in A$. For every $\varepsilon > 0$ there exists $z' \in A$ such that $|y - z'| \leq \text{dist}(y, A) + \varepsilon$. Thus

$$\text{dist}(x, A) \leq |x - z'| \leq |x - y| + \text{dist}(y, A) + \varepsilon,$$

which implies

$$\text{dist}(x, A) - \text{dist}(y, A) \leq |x - y| + \varepsilon.$$

By switching the roles of x and y , we obtain

$$|\text{dist}(x, A) - \text{dist}(y, A)| \leq |x - y| + \varepsilon.$$

This holds for every $\varepsilon > 0$, so that

$$|\text{dist}(x, A) - \text{dist}(y, A)| \leq |x - y|. \quad \square$$

Lemma 1.28. Let $A \subset \mathbb{R}^n$ be an open set with $\partial A \neq \emptyset$ and let

$$A_i = \{x \in A : \text{dist}(x, \partial A) > \frac{1}{i}\}, \quad i = 1, 2, \dots$$

Then the sets A_i are open, $A_i \subset A_{i+1}$, $i = 1, 2, \dots$, and $A = \bigcup_{i=1}^{\infty} A_i$.

T H E M O R A L : Any open set can be exhausted by an increasing sequence of distance sets.

Proof. Recall that a function is continuous if and only if the preimage of every open set is open. By Lemma 1.27 the function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $f(x) = \text{dist}(x, \partial A)$ is continuous and thus

$$\{x \in A : \text{dist}(x, \partial A) > \frac{1}{i}\} = f^{-1}((\frac{1}{i}, \infty))$$

is an open set. It is immediate that $A_i \subset A_{i+1}$, $i = 1, 2, \dots$.

Since $A_i \subset A$ for every $i = 1, 2, \dots$, we have $\bigcup_{i=1}^{\infty} A_i \subset A$. On the other hand, since A is open, for every $x \in A$ there exists $\varepsilon > 0$ such that $B(x, \varepsilon) \subset A$. This implies $\text{dist}(x, \partial A) \geq \varepsilon$. Thus we may choose i large enough so that $x \in A_i$. This shows that $A \subset \bigcup_{i=1}^{\infty} A_i$. \square

Remarks 1.29:

(1) Let $A \subset \mathbb{R}^n$ be an open set with $\partial A \neq \emptyset$ and let

$$A_i = \{x \in A : \text{dist}(x, \partial A) \geq \frac{1}{i}\}, \quad i = 1, 2, \dots$$

Then the sets A_i are closed, $A_i \subset A_{i+1}$, $i = 1, 2, \dots$, and $A = \bigcup_{i=1}^{\infty} A_i$.

(2) Let $A \subset \mathbb{R}^n$, $A \neq \emptyset$, be a closed set with and let

$$A_i = \{x \in \mathbb{R}^n : \text{dist}(x, A) < \frac{1}{i}\}, \quad i = 1, 2, \dots$$

Then the sets A_i are open, $A_{i+1} \subset A_i$, $i = 1, 2, \dots$, and $A = \bigcap_{i=1}^{\infty} A_i$. Thus any closed set can be represented as an intersection of a decreasing sequence of open sets. The corresponding claim with closed sets is obtained by considering

$$A_i = \{x \in \mathbb{R}^n : \text{dist}(x, A) \leq \frac{1}{i}\}, \quad i = 1, 2, \dots$$

Lemma 1.30. If $A \subset \mathbb{R}^n$ is an open set with $\partial A \neq \emptyset$, then $\text{dist}(K, \partial A) > 0$ for every compact subset K of A .

Proof. Since $x \mapsto \text{dist}(x, \partial A)$ is a continuous function, it attains its minimum on any compact set. Thus there exists $z \in K$ such that $\text{dist}(z, \partial A) = \text{dist}(K, \partial A)$. Since A is open and $z \in A$, there exists $\varepsilon > 0$ such that $B(z, \varepsilon) \subset A$. This implies

$$\text{dist}(K, \partial A) = \text{dist}(z, \partial A) \geq \varepsilon > 0.$$

□

W A R N I N G : The corresponding claim does not hold if $K \subset A$ only assumed to be closed. For example, $A = \{(x, y) \in \mathbb{R}^2 : y > 0\}$ is open, $K = \{(x, y) \in \mathbb{R}^2 : y \geq e^x\}$ is closed and $K \subset A$. However, $\text{dist}(K, A) = 0$.

Remark 1.31. In addition, the distance function has the following properties (exercise):

- (1) $x \in \overline{A}$ if and only if $\text{dist}(x, A) = 0$,
- (2) $\emptyset \neq A \subset B$ implies $\text{dist}(x, A) \geq \text{dist}(x, B)$,
- (3) $\text{dist}(x, A) = \text{dist}(x, \overline{A})$ for every $x \in \mathbb{R}^n$ and
- (4) $\overline{A} = \overline{B}$ if and only if $\text{dist}(x, A) = \text{dist}(x, B)$ for every $x \in \mathbb{R}^n$.

Remark 1.32. The distance function can be used to construct a cutoff function, which is useful in localization arguments and partitions of unity. Assume that $G \subset \mathbb{R}^n$ is open and $F \subset G$ closed. Then there exist a continuous function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

- (1) $0 \leq f(x) \leq 1$ for every $x \in \mathbb{R}^n$,
- (2) $f(x) = 1$ for every $x \in F$ and
- (3) $f(x) = 0$ for every $x \in \mathbb{R}^n \setminus G$.

Reason. The claim is trivial if F or $\mathbb{R}^n \setminus G$ is empty. Thus we may assume that both sets are nonempty. Define

$$f(x) = \frac{\text{dist}(x, \mathbb{R}^n \setminus G)}{\text{dist}(x, \mathbb{R}^n \setminus G) + \text{dist}(x, F)}.$$

This function has the desired properties. The claim (1) is clear. To prove (2), let $x \in F$. Then $\text{dist}(x, F) = 0$. On the other hand, since $x \in F \subset G$ and G is open, there exists $r > 0$ such that $B(x, r) \subset G$. This implies $\text{dist}(x, \mathbb{R}^n \setminus G) \geq r > 0$ and thus $f(x) = 1$. The claim (3) is clear. ■

1.5 Characterizations of measurable sets

In this section we assume that $X = \mathbb{R}^n$ even though most of the results hold true in a more general context. We discuss a useful method to construct σ -algebras, see Definition 1.8. A σ -algebra generated by a collection of sets \mathcal{E} is the smallest σ -algebra containing \mathcal{E} . Next we show that this definition well-stated.

Lemma 1.33. Let \mathcal{E} be a collection of subsets of X . There exists a unique smallest σ -algebra \mathcal{A} containing \mathcal{E} , that is, \mathcal{A} is a σ -algebra, $\mathcal{E} \subset \mathcal{A}$, and if \mathcal{B} is any other σ -algebra with $\mathcal{E} \subset \mathcal{B}$, then $\mathcal{A} \subset \mathcal{B}$.

Proof. Let \mathcal{S} be the collection of all σ -algebras \mathcal{B} that contain \mathcal{E} and consider

$$\mathcal{A} = \bigcap_{\mathcal{B} \in \mathcal{S}} \mathcal{B} = \{A \subset X : \text{if } \mathcal{B} \text{ is a } \sigma\text{-algebra with } \mathcal{E} \subset \mathcal{B}, \text{ then } A \in \mathcal{B}\}$$

The collection \mathcal{A} is a σ -algebra, since the intersection of σ -algebras is a σ -algebra. It is easy to verify that \mathcal{A} has the required properties (exercise). □

Example 1.34. Let $A \subset X$. Then the smallest σ -algebra containing A is $\{\emptyset, X, A, X \setminus A\}$.

Definition 1.35. The collection \mathcal{B} of Borel sets is the smallest σ -algebra containing all open subsets of \mathbb{R}^n .

Remarks 1.36:

- (1) Since any σ -algebra is closed with respect to complements, the collection \mathcal{B} of Borel sets can be also defined as the smallest σ -algebra containing, for example, the closed subsets of \mathbb{R}^n . In fact \mathcal{B} is generated by open and closed intervals, because every open set is a countable union of open (or closed) intervals by the Lindelöf theorem.
- (2) Note that the collection of Borel sets does not only contain open and closed sets, but it also contains, for example, the G_δ -sets which are countable intersections of open sets and the F_σ -sets which are countable unions of closed sets. Note that every closed set is G_δ and every open set is F_σ . On

the other hand, the half-open interval $[0, 1)$ is not open nor closed, but both G_δ and F_σ , since it can be expressed as both a countable union of closed sets and a countable intersection of open sets

$$[0, 1) = \bigcup_{i=1}^{\infty} \left[0, 1 - \frac{1}{i}\right] = \bigcap_{i=1}^{\infty} \left(-\frac{1}{i}, 1\right).$$

We can apply the operations σ and δ repeatedly and obtain the classes

$$G \subset G_\delta \subset G_{\delta\sigma} \subset G_{\delta\sigma\delta} \dots \quad \text{and} \quad F \subset F_\sigma \subset F_{\sigma\delta} \subset F_{\sigma\delta\sigma} \dots,$$

where G denotes the collection of open sets, F the collection of closed sets, σ the operation of countable unions and δ the operation of countable intersections. Note that $G = G_\sigma$, $G_{\sigma\sigma} = G_\sigma, \dots$ and $F = F_\delta$, $F_{\delta\delta} = F_\delta, \dots$. By Remark 1.29 every open set can be represented as a union of countably many closed sets and every closed set can be represented as an intersection of countably many open sets. This implies that $G \subset F_\sigma$ and $F \subset G_\delta$. It also holds that $G_\delta \subset F_{\sigma\delta}$, $G_{\delta\sigma} \subset F_{\sigma\delta\sigma}, \dots$ and $F_\sigma \subset G_{\delta\sigma}$, $F_{\sigma\delta} \subset G_{\delta\sigma\delta}, \dots$. It turns out that these are all different classes of sets and that there are Borel sets that do not belong to any of them, see [3, Chapter 8].

- (3) One way to show that Borel sets have a certain property is to construct a σ -algebra containing open (or closed) sets, or open (or closed) intervals, that satisfies the required property. Since Borel sets is the smallest σ -algebra with this property, every Borel set satisfies the required property.

Next we discuss locally finite Borel regular outer measures.

Definition 1.37. Let μ^* be an outer measure on \mathbb{R}^n .

- (1) μ^* is called a Borel outer measure, if all Borel sets are μ^* -measurable.
- (2) A Borel outer measure μ^* is called Borel regular, if for every set $A \subset \mathbb{R}^n$ there exists a Borel set B such that $A \subset B$ and $\mu^*(A) = \mu^*(B)$.
- (3) μ^* is a Radon outer measure, if μ^* is Borel regular and $\mu^*(K) < \infty$ for every compact set $K \subset \mathbb{R}^n$.

T H E M O R A L : We shall see that the Lebesgue outer measure is a Radon outer measure. Radon outer measures have many good approximation properties similar to the Lebesgue measure. There is also a natural way to construct Radon outer measures by the Riesz representation theorem, but this will be discussed in the real analysis course.

Remarks 1.38:

- (1) In particular, all open and closed sets are measurable for a Borel outer measure. Thus the collection of measurable sets is relatively large.

(2) In general, an outer measure μ^* is called regular, if for every set $A \subset X$ there exists a μ^* -measurable set B such that $A \subset B$ and $\mu^*(A) = \mu^*(B)$. Roughly speaking, regularity enables us to pass some properties of Borel sets to measurable sets. Many natural constructions of outer measures give regular measures, see Remark 1.20. On the other hand, the outer measure discussed in Example 1.3 and Example 1.7 is not regular (exercise).

(3) The local finiteness condition $\mu^*(K) < \infty$ for every compact set $K \subset \mathbb{R}^n$ is equivalent with the condition $\mu^*(B(x, r)) < \infty$ for every $x \in \mathbb{R}^n$ and $r > 0$. In particular, this implies that μ^* is σ -finite, see Definition 1.18.

Examples 1.39:

(1) The Dirac outer measure is a Radon outer measure (exercise).

(2) The counting measure is Borel regular on any metric space X , but it is a Radon outer measure only if every compact subset of X is finite (exercise).

Lemma 1.40. The Lebesgue outer measure m^* is Borel regular.

Proof. We may assume that $m^*(A) < \infty$, for otherwise we may take $B = \mathbb{R}^n$ with $m^*(\mathbb{R}^n) = \infty$, see Remark 1.68 (4). For every $i = 1, 2, \dots$ there exist intervals $I_{i,j}$, $j = 1, 2, \dots$, such that $A \subset \bigcup_{j=1}^{\infty} I_{i,j}$ and

$$m^*(A) \leq \sum_{j=1}^{\infty} \text{vol}(I_{i,j}) < m^*(A) + \frac{1}{i}.$$

Denote $B_i = \bigcup_{j=1}^{\infty} I_{i,j}$, $i = 1, 2, \dots$. The set B_i , $i = 1, 2, \dots$, is a Borel set as a countable union of closed intervals. This implies that $B = \bigcap_{i=1}^{\infty} B_i$ is a Borel set. Moreover, since $A \subset B_i$, we have $A \subset B \subset B_i$ for every $i = 1, 2, \dots$. By monotonicity and the definition of the Lebesgue outer measure, we obtain

$$m^*(A) \leq m^*(B) \leq m^*(B_i) = m^*\left(\bigcup_{j=1}^{\infty} I_{i,j}\right) \leq \sum_{j=1}^{\infty} \text{vol}(I_{i,j}) < m^*(A) + \frac{1}{i}.$$

By letting $i \rightarrow \infty$, we conclude $m^*(A) = m^*(B)$. Later we show that all Borel sets are Lebesgue measurable, see Lemma 1.55 and Lemma 1.67. \square

The next result asserts that the Lebesgue outer measure is locally finite.

Lemma 1.41. The Lebesgue outer measure satisfies $m^*(K) < \infty$ for every compact set $K \subset \mathbb{R}^n$.

Proof. Since $K \subset \mathbb{R}^n$ is compact it is closed and bounded. Thus there exists an interval $I = [a_1, b_1] \times \dots \times [a_n, b_n]$, $a_i, b_i \in \mathbb{R}$, $i = 1, 2, \dots, n$, such that $K \subset I$. By the definition of the Lebesgue outer measure, we have

$$m^*(K) \leq \text{vol}(I) = (b_1 - a_1) \cdots (b_n - a_n) < \infty.$$

\square

Remark 1.42. The Hausdorff measures, defined in Example 1.4 (6), are not necessarily locally finite. For example, the one-dimensional Hausdorff measure \mathcal{H}^1 is a Borel regular outer measure but not a Radon outer measure on \mathbb{R}^2 , because $\mathcal{H}^1(\overline{B(0,1)}) = \infty$ and the closed unit ball $\overline{B(0,1)} = \{x \in \mathbb{R}^2 : |x| \leq 1\}$ is a compact subset of \mathbb{R}^2 . We shall show later that all Borel sets are measurable with respect to the Hausdorff outer measure, see Remark 1.54.

We discuss an approximation result for a measurable set with respect to a Radon outer measure. In Lemma 1.43 we assume that $\mu^*(\mathbb{R}^n) < \infty$, but Theorem 1.48 shows that the result holds also when $\mu^*(\mathbb{R}^n) = \infty$.

Lemma 1.43. Let μ^* be a Radon outer measure on \mathbb{R}^n with $\mu^*(\mathbb{R}^n) < \infty$ and assume that $A \subset \mathbb{R}^n$ is a μ^* -measurable set. Then, for every $\varepsilon > 0$, there exists a closed set F and an open set G such that

$$F \subset A \subset G, \quad \mu^*(A \setminus F) < \varepsilon \quad \text{and} \quad \mu^*(G \setminus A) < \varepsilon.$$

T H E M O R A L : A measurable set with respect to a Radon outer measure can be approximated by closed sets from inside and open sets from outside up to a set of arbitrarily small measure.




Figure 1.7: Approximation of a measurable set.

Proof. **Step 1:** Let

$$\mathcal{F} = \{A \subset \mathbb{R}^n : A \text{ is } \mu^*\text{-measurable, for every } \varepsilon > 0 \text{ there exists a closed } F \subset A \text{ such that } \mu^*(A \setminus F) < \varepsilon \text{ and an open } G \supset A \text{ such that } \mu^*(G \setminus A) < \varepsilon\}$$

be the collection of measurable sets that has the required approximation property.

S T R A T E G Y : We show that \mathcal{F} is a σ -algebra that contains all closed sets. Since the collection of Borel sets is the smallest σ -algebra with this property, every Borel set belongs to \mathcal{F} . This implies that every Borel set has the desired approximation property. Borel regularity takes care of the rest.

It is clear that $\emptyset \in \mathcal{F}$ and that $A \in \mathcal{F}$ implies $\mathbb{R}^n \setminus A \in \mathcal{F}$. Let $A_i \in \mathcal{F}$, $i = 1, 2, \dots$. We show that $\bigcap_{i=1}^{\infty} A_i \in \mathcal{F}$. By de Morgan's law this implies that $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$. Let $\varepsilon > 0$. Since $A_i \in \mathcal{F}$ there exist a closed set F_i and an open set G_i such that

$$F_i \subset A_i \subset G_i, \quad \mu^*(A_i \setminus F_i) < \frac{\varepsilon}{2^{i+1}} \quad \text{and} \quad \mu^*(G_i \setminus A_i) < \frac{\varepsilon}{2^{i+1}}$$

for every $i = 1, 2, \dots$. It follows that

$$\begin{aligned} \mu^*\left(\bigcap_{i=1}^{\infty} A_i \setminus \bigcap_{i=1}^{\infty} F_i\right) &\leq \mu^*\left(\bigcup_{i=1}^{\infty} (A_i \setminus F_i)\right) \quad (\text{monotonicity}) \\ &\leq \sum_{i=1}^{\infty} \mu^*(A_i \setminus F_i) \quad (\text{subadditivity}) \\ &\leq \varepsilon \sum_{i=1}^{\infty} \frac{1}{2^{i+1}} < \varepsilon. \end{aligned}$$

Since $\bigcap_{i=1}^{\infty} F_i$ is a closed set, it will do as an approximation from inside. On the other hand, since $\mu^*(\mathbb{R}^n) < \infty$, Theorem 1.22 (2) implies that

$$\lim_{k \rightarrow \infty} \mu^*\left(\bigcap_{i=1}^k G_i \setminus \bigcap_{i=1}^{\infty} A_i\right) = \mu^*\left(\bigcap_{i=1}^{\infty} G_i \setminus \bigcap_{i=1}^{\infty} A_i\right) < \varepsilon.$$

The last inequality is proved as above. Consequently, there exists an index k such that

$$\mu^*\left(\bigcap_{i=1}^k G_i \setminus \bigcap_{i=1}^{\infty} A_i\right) < \varepsilon.$$

As an intersection of finitely many open sets, $\bigcap_{i=1}^k G_i$ is an open set, and it will do as an approximation from outside. This shows that \mathcal{F} is a σ -algebra.

Next we show that \mathcal{F} contains closed sets. Assume that $A \subset \mathbb{R}^n$ is a closed set and let $\varepsilon > 0$. Then $\mu^*(A \setminus A) = 0 < \varepsilon$, so that A itself will do as an approximation from inside in the definition of \mathcal{F} . To obtain an approximation from outside, we represent A as an intersection of countably many open sets by using the distance function as in Lemma 1.28, see also Remark 1.29. Since A is closed we have

$$A = \bigcap_{i=1}^{\infty} A_i, \quad \text{where} \quad A_i = \{x \in \mathbb{R}^n : \text{dist}(x, A) < \frac{1}{i}\}, \quad i = 1, 2, \dots$$

The sets A_i , $i = 1, 2, \dots$, are open, because $x \mapsto \text{dist}(x, A)$ is continuous, see Lemma 1.27. Since $\mu^*(\mathbb{R}^n) < \infty$ and $A_1 \supset A_2 \supset \dots$, Theorem 1.22 implies

$$\lim_{i \rightarrow \infty} \mu^*(A_i \setminus A) = \mu^*\left(\bigcap_{i=1}^{\infty} A_i \setminus A\right) = \mu^*(\emptyset) = 0,$$

and there exists an index i such that $\mu^*(A_i \setminus A) < \varepsilon$. This A_i will do as an approximation from outside in the definition of \mathcal{F} .

Thus \mathcal{F} is σ -algebra containing closed sets and consequently also Borel sets. This follows from the fact that the collection of Borel sets is the smallest σ -algebra with this property. This proves the claim in Lemma 1.43 for Borel sets.

Step 2: Then assume that $A \subset \mathbb{R}^n$ is a general μ^* -measurable set. By Borel regularity there exists a Borel set $B_1 \supset A$ with $\mu^*(B_1) = \mu^*(A)$ and a Borel set B_2 with $\mathbb{R}^n \setminus A \subset \mathbb{R}^n \setminus B_2$ and $\mu^*(\mathbb{R}^n \setminus B_2) = \mu^*(\mathbb{R}^n \setminus A)$. By Step 1 of the proof there exist a closed set F and an open set G such that

$$F \subset B_2 \subset A \subset B_1 \subset G, \quad \mu^*(G \setminus B_1) < \varepsilon \quad \text{and} \quad \mu^*(B_2 \setminus F) < \varepsilon.$$

It follows that

$$\begin{aligned} \mu^*(G \setminus A) &\leq \mu^*(G \setminus B_1) + \mu^*(B_1 \setminus A) \quad (\text{subadditivity, even = holds}) \\ &< \varepsilon + \mu^*(B_1) - \mu^*(A) \quad (\mu^*(B_1 \setminus A) = \mu^*(B_1) - \mu^*(A), \mu^*(A) < \infty) \\ &= \varepsilon \quad (\mu^*(B_1) - \mu^*(A) = 0) \end{aligned}$$

and

$$\begin{aligned} \mu^*(A \setminus F) &\leq \mu^*(A \setminus B_2) + \mu^*(B_2 \setminus F) \\ &\leq \mu^*\left((\mathbb{R}^n \setminus B_2) \setminus (\mathbb{R}^n \setminus A)\right) + \varepsilon \\ &= \mu^*(\mathbb{R}^n \setminus B_2) - \mu^*(\mathbb{R}^n \setminus A) + \varepsilon \quad (\mathbb{R}^n \setminus A \text{ measurable, } \mu^*(\mathbb{R}^n \setminus A) < \infty) \\ &= \varepsilon. \quad (\mu^*(\mathbb{R}^n \setminus B_2) - \mu^*(\mathbb{R}^n \setminus A) = 0, \mu^*(\mathbb{R}^n) < \infty) \quad \square \end{aligned}$$

Definition 1.44. Let μ^* be an outer measure on \mathbb{R}^n and E an arbitrary subset of \mathbb{R}^n . Then the restriction of μ^* to E is defined to be

$$(\mu^*|E)(A) = \mu^*(A \cap E)$$

for every $A \subset \mathbb{R}^n$.

T H E M O R A L : The restriction is a useful tool to make an outer measure finite by considering a restriction to a set with finite measure.

Remarks 1.45:

- (1) $\mu^*|E$ is an outer measure (exercise).
- (2) Any μ^* -measurable set is also $\mu^*|E$ -measurable (exercise). This holds for every set $E \subset \mathbb{R}^n$. In particular, the set E does not have to be μ^* -measurable. Note that not all $\mu^*|E$ -measurable sets need not be μ^* -measurable.
- (3) It is useful to consider restrictions of the Hausdorff measures, defined in Example 1.4, to sets with a lower Hausdorff dimension than n . Consider, for example, the one-dimensional Hausdorff measure \mathcal{H}^1 on \mathbb{R}^2 . By Remark 1.42 we have $\mathcal{H}^1(\mathbb{R}^2) = \infty$, but $(\mathcal{H}^1|_{\partial B(0,1)})(\mathbb{R}^2) < \infty$.

Lemma 1.46. Let μ^* be a Borel regular outer measure on \mathbb{R}^n . Assume that $E \subset \mathbb{R}^n$ is μ^* -measurable and $\mu^*(E) < \infty$. Then $\mu^*|E$ is a Radon outer measure.

Remarks 1.47:

- (1) The assumption $\mu^*(E) < \infty$ cannot be removed, see Remark 1.42.
- (2) If E is a Borel set, then $\mu^*|E$ is Borel regular even if $\mu^*(E) = \infty$ (exercise).

Proof. Let $\nu = \mu^*|E$. Since every μ^* -measurable set is ν -measurable, ν is a Borel outer measure. If $K \subset \mathbb{R}^n$ is compact, then

$$\nu(K) = \mu^*(K \cap E) \leq \mu^*(E) < \infty.$$

C L A I M : ν is Borel regular.

Since μ^* is Borel regular, there exists a Borel set B_1 such that $E \subset B_1$ and $\mu^*(B_1) = \mu^*(E)$. Then

$$\begin{aligned} \mu^*(B_1) &= \mu^*(B_1 \cap E) + \mu^*(B_1 \setminus E) \quad (E \text{ is } \mu^*\text{-measurable}) \\ &= \mu^*(E) + \mu^*(B_1 \setminus E) \quad (E \subset B_1) \end{aligned}$$

Since $\mu^*(E) < \infty$, we have $\mu^*(B_1 \setminus E) = \mu^*(B_1) - \mu^*(E) = 0$.

Let $A \subset \mathbb{R}^n$. Since μ^* is Borel regular, there exists a Borel set B_2 such that $B_1 \cap A \subset B_2$ and $\mu^*(B_2) = \mu^*(B_1 \cap A)$. Then $A \subset B_2 \cup (\mathbb{R}^n \setminus B_1) = C$ and C is a Borel set as a union of two Borel sets. We have

$$\begin{aligned} (\mu^*|E)(C) &= \mu^*(C \cap E) \leq \mu^*(B_1 \cap C) \quad (E \subset B_1) \\ &= \mu^*(B_1 \cap B_2) \quad (B_1 \cap C = B_1 \cap (B_2 \cup (\mathbb{R}^n \setminus B_1)) = B_1 \cap B_2) \\ &\leq \mu^*(B_2) = \mu^*(B_1 \cap A) \\ &= \mu^*((B_1 \cap A) \cap E) + \mu^*((B_1 \cap A) \setminus E) \quad (E \text{ is } \mu^*\text{-measurable}) \\ &\leq \mu^*(E \cap A) + \mu^*(B_1 \setminus E) \quad (\text{monotonicity}) \\ &= (\mu^*|E)(A). \quad (\mu^*(B_1 \setminus E) = 0) \end{aligned}$$

On the other hand, since $A \subset C$ we have $(\mu^*|E)(A) \leq (\mu^*|E)(C)$. Consequently $(\mu^*|E)(A) = (\mu^*|E)(C)$ and $\mu^*|E$ is Borel regular. \square

Next we discuss the first characterization of measurable sets with respect to a Radon outer measure.

Theorem 1.48. Let μ^* be a Radon outer measure on \mathbb{R}^n . Then the following conditions are equivalent.

- (1) $A \subset \mathbb{R}^n$ is μ^* -measurable.
- (2) For every $\varepsilon > 0$ there exists a closed set F and an open set G such that $F \subset A \subset G$, $\mu^*(A \setminus F) < \varepsilon$ and $\mu^*(G \setminus A) < \varepsilon$.

T H E M O R A L : This is a topological characterization of a measurable set by squeezing it between closed and open sets. A set is measurable for a Radon outer measure if and only if it can be approximated by closed sets from inside and open sets from outside up to a set of arbitrarily small measure. Observe that the original Carathéodory criterion for measurability in Definition 1.5 depends only on the outer measure and there is no reference to open or closed sets. The assumption that a Radon measure is Borel regular plays an important role here.

Proof. $\boxed{(1) \implies (2)}$ Let $v_i = \mu^*|B(0, i)$, with $B(0, i) = \{x \in \mathbb{R}^n : |x| < i\}$, $i = 1, 2, \dots$. By Lemma 1.46, v_i is a Radon outer measure and $v_i(\mathbb{R}^n) \leq \mu^*(\overline{B(0, i)}) < \infty$ for every $i = 1, 2, \dots$. Since A is μ^* -measurable, A is also v_i -measurable.

By Lemma 1.43, there exists an open set $G_i \supset A$ such that

$$v_i(G_i \setminus A) < \frac{\varepsilon}{2^{i+1}},$$

for every $i = 1, 2, \dots$. Let

$$G = \bigcup_{i=1}^{\infty} (G_i \cap B(0, i))$$

As a union of open sets, the set G is open and $G \supset A$. Moreover, we have

$$\begin{aligned} \mu^*(G \setminus A) &= \mu^*\left(\left(\bigcup_{i=1}^{\infty} (G_i \cap B(0, i))\right) \setminus A\right) \\ &= \mu^*\left(\bigcup_{i=1}^{\infty} ((G_i \setminus A) \cap B(0, i))\right) \\ &\leq \sum_{i=1}^{\infty} \mu^*((G_i \setminus A) \cap B(0, i)) \\ &\leq \sum_{i=1}^{\infty} v_i(G_i \setminus A) \leq \sum_{i=1}^{\infty} \frac{\varepsilon}{2^{i+1}} < \varepsilon. \end{aligned}$$

Thus G will do as an approximation from outside.

By considering the complements, there exists a closed set F such that

$$\mathbb{R}^n \setminus F \supset \mathbb{R}^n \setminus A \quad \text{and} \quad \mu^*(A \setminus F) = \mu^*((\mathbb{R}^n \setminus F) \setminus (\mathbb{R}^n \setminus A)) < \varepsilon.$$

The set F is closed and will do as an approximation from inside.

$\boxed{(2) \implies (1)}$ For every $i = 1, 2, \dots$ there exists a closed set $F_i \subset A$ such that $\mu^*(A \setminus F_i) < \frac{1}{i}$. Then $F = \bigcup_{i=1}^{\infty} F_i$ is a Borel set (not necessarily closed) and $F \subset A$. Moreover,

$$0 \leq \mu^*(A \setminus F) = \mu^*\left(A \setminus \bigcup_{i=1}^{\infty} F_i\right) = \mu^*\left(\bigcap_{i=1}^{\infty} (A \setminus F_i)\right) \leq \mu^*(A \setminus F_i) < \frac{1}{i}$$

for every $i = 1, 2, \dots$. This implies that $\mu^*(A \setminus F) = 0$. Observe that $A = F \cup (A \setminus F)$, where F is a Borel set and hence μ^* -measurable. On the other hand, $\mu^*(A \setminus F) = 0$ so that $A \setminus F$ is μ^* -measurable. The set A is μ^* -measurable as a union of two measurable sets. \square




Figure 1.8: Approximation by open sets in measure.

By the previous theorem we can approximate measurable sets for a Radon outer measure μ^* by open or closed sets up to a set of arbitrary small but, in general, nonzero measure. By taking countable intersections of open sets or countable unions of closed sets, we can approximate measurable sets by Borel sets up to sets of measure zero.

Corollary 1.49. Let μ^* be a Radon outer measure on \mathbb{R}^n . Then the following claims are equivalent for a set $A \subset \mathbb{R}^n$.

- (1) A is μ^* -measurable.
- (2) There exists a G_δ set G such that $A \subset G$ and $\mu^*(G \setminus A) = 0$.
- (3) There exists an F_σ set F such that $F \subset A$ and $\mu^*(A \setminus F) = 0$.

T H E M O R A L : An arbitrary measurable set with respect to a Radon measure differs from a Borel set only by a set of measure zero.

Proof. $\boxed{(1) \iff (2)}$ By Theorem 1.48, for every $i = 1, 2, \dots$, there exists an open set $G_i \supset A$ such that $\mu^*(G_i \setminus A) < \frac{1}{i}$. Then $G = \bigcap_{i=1}^{\infty} G_i$ is a G_δ set such that $A \subset G$. Since $G \setminus A \subset G_i \setminus A$ for every $i = 1, 2, \dots$, we have

$$0 \leq \mu^*(G \setminus A) \leq \mu^*(G_i \setminus A) < \frac{1}{i}$$

for every $i = 1, 2, \dots$. This implies that $\mu^*(G \setminus A) = 0$.

Then assume that $A \subset \mathbb{R}^n$ is a set such that there exists a G_δ set G such that $A \subset G$ and $\mu^*(G \setminus A) = 0$. The set $G \setminus A$ is μ^* -measurable as a set of measure zero and thus $A = G \setminus (G \setminus A)$ is μ^* -measurable as a union of two μ^* -measurable sets.

(1) \iff (3) By Theorem 1.48, for every $i = 1, 2, \dots$, there exists a closed set $F_i \subset A$ such that $\mu^*(A \setminus F_i) < \frac{1}{i}$. Then $F = \bigcup_{i=1}^{\infty} F_i$ is an \mathcal{F}_σ -set and $F \subset A$. The claim follows as in the proof of the implication (2) \implies (1) in Theorem 1.48.

Then assume that $A \subset \mathbb{R}^n$ is a set such that there exists an F_σ set F such that $F \subset A$ and $\mu^*(A \setminus F) = 0$. Observe that The set $A \setminus F$ is μ^* -measurable as a set of measure zero and thus $A = F \cup (A \setminus F)$ is μ^* -measurable as a union of two μ^* -measurable sets. \square

Remarks 1.50:

- (1) It follows from the proof of Corollary 1.49 that a set $A \subset \mathbb{R}^n$ is μ^* -measurable if and only if for every $\varepsilon > 0$ there exists an open set G such that $G \supset A$ and $\mu^*(G \setminus A) < \varepsilon$.
- (2) Alternatively a set $A \subset \mathbb{R}^n$ is μ^* -measurable, if for every $\varepsilon > 0$ there exists a closed set F with $F \supset A$ and $\mu^*(A \setminus F) < \varepsilon$.
- (3) The claim that $A \subset \mathbb{R}^n$ if and only if there exists a Borel set B such that $A \subset B$ and $\mu^*(B \setminus A) = 0$ follows directly from Borel regularity. Assume that $\mu^*(A) < \infty$. Since μ^* is Borel regular, for every $A \subset \mathbb{R}^n$, there exists a Borel set B such that $A \subset B$ and $\mu^*(B) = \mu^*(A)$. If A is μ^* -measurable, then $\mu^*(B) = \mu^*(B \setminus A) + \mu^*(A)$ and we may conclude that $\mu^*(B \setminus A) = 0$. The other implication follows as above.

T H E M O R A L : Measurability of a set can be characterized separately by approximations be open sets form outside or closed sets from inside, compare with Theorem 1.48.

Corollary 1.51. Let μ^* be a Radon outer measure on \mathbb{R}^n .

- (1) (Outer measure) For every set $A \subset \mathbb{R}^n$,

$$\mu^*(A) = \inf \{\mu^*(G) : A \subset G, G \text{ open}\}.$$

- (2) (Inner measure) For every μ^* -measurable set $A \subset \mathbb{R}^n$,

$$\mu^*(A) = \sup \{\mu^*(K) : K \subset A, K \text{ compact}\}.$$

T H E M O R A L : The inner and outer measures coincide for a measurable set with respect to a Radon measure on \mathbb{R}^n . In this case, the measure can be determined by compact sets from inside or open sets from outside.

Proof. (1) If $\mu^*(A) = \infty$, the claim is clear. Hence we may assume that $\mu^*(A) < \infty$.

Step 1: Assume that A is a Borel set and let $\varepsilon > 0$. Since μ^* is a Borel outer measure, the set A is μ^* -measurable. By Theorem 1.48, there exists an open set

$G \supset A$ such that $\mu^*(G \setminus A) < \varepsilon$. Moreover, we have

$$\begin{aligned}\mu^*(G) &= \mu^*(G \cap A) + \mu^*(G \setminus A) \quad (A \text{ is } \mu^*\text{-measurable}) \\ &= \mu^*(A) + \mu^*(G \setminus A) < \mu^*(A) + \varepsilon. \quad (A \subset G)\end{aligned}$$

This implies the claim.

Step 2: Assume then that $A \subset \mathbb{R}^n$ is an arbitrary set. Since μ^* is Borel regular, there exists a Borel set $B \supset A$ with $\mu^*(B) = \mu^*(A)$. It follows that

$$\begin{aligned}\mu^*(A) &= \mu^*(B) = \inf\{\mu^*(G) : B \subset G, G \text{ open}\} \quad (\text{Step 1}) \\ &\geq \inf\{\mu^*(G) : A \subset G, G \text{ open}\}. \quad (A \subset B)\end{aligned}$$

On the other hand, by monotonicity, we have

$$\mu^*(A) \leq \inf\{\mu^*(G) : A \subset G, G \text{ open}\}$$

and, consequently, the equality holds.

(2) Assume first that $\mu^*(A) < \infty$ and let $\varepsilon > 0$. By Theorem 1.48, there exists a closed set $F \subset A$ such that $\mu^*(A \setminus F) < \varepsilon$. Since F is μ^* -measurable and $\mu^*(A) < \infty$, we have

$$\mu^*(A) - \mu^*(F) = \mu^*(A \setminus F) < \varepsilon$$

and thus $\mu^*(F) > \mu^*(A) - \varepsilon$. This implies that

$$\mu^*(A) = \sup\{\mu^*(F) : F \subset A, F \text{ closed}\}$$

Then we consider the case $\mu^*(A) = \infty$. Let $B_i = \{x \in \mathbb{R}^n : i-1 \leq |x| < i\}$, $i = 1, 2, \dots$. Then $A = \bigcup_{i=1}^{\infty} (A \cap B_i)$ and Theorem 1.12 implies that

$$\sum_{i=1}^{\infty} \mu^*(A \cap B_i) = \mu^*(A) = \infty$$

because the sets $A \cap B_i$, $i = 1, 2, \dots$, are pairwise disjoint and μ^* -measurable. Since μ^* is a Radon outer measure, $\mu^*(A \cap B_i) \leq \mu^*(\overline{B_i}) < \infty$. By the beginning of the proof, there exists a closed set $F_i \subset A \cap B_i$ such that

$$\mu^*(F_i) > \mu^*(A \cap B_i) - \frac{1}{2^i}$$

for every $i = 1, 2, \dots$. Clearly $\bigcup_{i=1}^{\infty} F_i \subset A$ and

$$\begin{aligned}\lim_{k \rightarrow \infty} \mu^*\left(\bigcup_{i=1}^k F_i\right) &= \mu^*\left(\bigcup_{i=1}^{\infty} F_i\right) \quad (\text{Theorem 1.22}) \\ &= \sum_{i=1}^{\infty} \mu^*(F_i) \quad (F_i \text{ disjoint } (F_i \subset A \cap B_i), \text{ Theorem 1.12}) \\ &\geq \sum_{i=1}^{\infty} \left(\mu^*(A \cap B_i) - \frac{1}{2^i}\right) = \infty.\end{aligned}$$

The set $F = \bigcup_{i=1}^k F_i$ is closed as a union of finitely many closed sets and hence

$$\mu^*(A) = \sup \{\mu^*(F) : F \subset A, F \text{ closed}\} = \infty.$$

Finally we pass over to compact sets. Assume that F is closed. Then the sets $F \cap \overline{B(0, i)}$, $i = 1, 2, \dots$, are closed and bounded and hence compact. By Theorem 1.22,

$$\mu^*(F) = \mu^*\left(\bigcup_{i=1}^{\infty}(F \cap \overline{B(0, i)})\right) = \lim_{i \rightarrow \infty} \mu^*(F \cap \overline{B(0, i)})$$

and consequently

$$\sup \{\mu^*(K) : K \subset A, K \text{ compact}\} = \sup \{\mu^*(F) : F \subset A, F \text{ closed}\}.$$

□

Remarks 1.52:

(1) Let μ^* be a Radon outer measure on \mathbb{R}^n . A set $A \subset \mathbb{R}^n$ is μ^* -measurable if and only if

$$\mu^*(A) = \sup \{\mu^*(K) : K \subset A, K \text{ compact}\}.$$

A μ^* -measurable set A has the property above by Corollary 1.51 (2). To prove the other implication, assume that the property above holds for $A \subset \mathbb{R}^n$ with $\mu^*(A) < \infty$. Then for every $i = 1, 2, \dots$, there exists a compact set $K_i \subset A$ such that $\mu^*(K_i) > \mu^*(A) - \frac{1}{i}$. Since K_i is μ^* -measurable, by Remark 1.6 (3) we have $\mu(A \setminus K_i) = \mu^*(A) - \mu^*(K_i) < \frac{1}{i}$, $i = 1, 2, \dots$. Then $K = \bigcup_{i=1}^{\infty} K_i$ is an \mathcal{F}_σ -set and $K \subset A$. Moreover,

$$0 \leq \mu^*(A \setminus K) = \mu^*\left(A \setminus \bigcup_{i=1}^{\infty} K_i\right) = \mu^*\left(\bigcap_{i=1}^{\infty}(A \setminus K_i)\right) \leq \mu^*(A \setminus K_i) < \frac{1}{i}$$

for every $i = 1, 2, \dots$. This implies that $\mu^*(A \setminus K) = 0$. Observe that $A = K \cup (A \setminus K)$, where K is a Borel set and hence μ^* -measurable. On the other hand, $\mu^*(A \setminus K) = 0$ so that $A \setminus K$ is μ^* -measurable. The set A is μ^* -measurable as a union of two measurable sets.

(2) Let μ^* be a Radon outer measure on \mathbb{R}^n . There is a delicate issue related to the approximation by open sets. Let $A \subset \mathbb{R}^n$ with $\mu^*(A) < \infty$. By Corollary 1.51 (1), for every $\varepsilon > 0$, there exists an open set G such that $A \subset G$ and $\mu^*(G) < \mu^*(A) + \varepsilon$. This holds for an arbitrary set A . Since $G = A \cup (G \setminus A)$ if $A \subset G$, by subadditivity, we have $\mu^*(G) \leq \mu^*(A) + \mu^*(G \setminus A)$. In this case we cannot conclude from $\mu^*(G) < \mu^*(A) + \varepsilon$ that $\mu^*(G \setminus A) < \varepsilon$. However, if A is μ^* -measurable, then $\mu^*(G) = \mu^*(A) + \mu^*(G \setminus A)$ and $\mu^*(G) < \mu^*(A) + \varepsilon$ implies that $\mu^*(G \setminus A) < \varepsilon$.

1.6 Metric outer measures

Next we give a useful method to show that Borel sets are measurable.

Theorem 1.53. Let μ^* be an outer measure on \mathbb{R}^n . If μ^* is a metric outer measure, that is,

$$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B)$$

for every $A, B \subset \mathbb{R}^n$ with $\text{dist}(A, B) > 0$, then μ^* is a Borel outer measure.

T H E M O R A L : If an outer measure is additive on separated sets, then all Borel sets are measurable. This is a practical way to show that Borel sets are measurable. This means that very useful properties of an outer measure on measurable sets are available for a large class of sets.

Proof. We shall show that every closed set $F \subset \mathbb{R}^n$ is μ^* -measurable. By Remark 1.6 (1), it is enough to show that

$$\mu^*(E) \geq \mu^*(E \cap F) + \mu^*(E \setminus F)$$

for every $E \subset \mathbb{R}^n$. By Remark 1.6 (1) we may assume that $\mu^*(E) < \infty$. The set $G = \mathbb{R}^n \setminus F$ is open. We separate the set $A = E \setminus F$ from F by considering the sets

$$A_i = \{x \in A : \text{dist}(x, F) \geq \frac{1}{i}\}, \quad i = 1, 2, \dots,$$

Then $\text{dist}(A_i, F) \geq \frac{1}{i}$ for every $i = 1, 2, \dots$, $A_i \subset A_{i+1}$ for $i = 1, 2, \dots$ and $A = \bigcup_{i=1}^{\infty} A_i$, see Lemma 1.28 and Remark 1.29.




Figure 1.9: Exhaustion by distance sets.

Claim: $\lim_{i \rightarrow \infty} \mu^*(A_i) = \mu^*(A)$.

Reason. $A_i \subset A_{i+1}$ and $A_i \subset A$, $i = 1, 2, \dots$, imply

$$\lim_{i \rightarrow \infty} \mu^*(A_i) \leq \mu^*(A).$$

Then we prove the reverse inequality. Let

$$B_i = A_{i+1} \setminus A_i = \{x \in A : \frac{1}{i+1} \leq \text{dist}(x, F) < \frac{1}{i}\}, \quad i = 1, 2, \dots$$

Since $A_i \subset A_{i+1}$ for $i = 1, 2, \dots$, we have $A = \bigcup_{i=1}^{\infty} A_i = A_i \cup \bigcup_{j=i}^{\infty} B_j$ and by countable subadditivity

$$\mu^*(A) \leq \mu^*(A_i) + \sum_{j=i}^{\infty} \mu^*(B_j).$$

It follows that

$$\mu^*(A) \leq \lim_{i \rightarrow \infty} \mu^*(A_i) + \lim_{i \rightarrow \infty} \sum_{j=i}^{\infty} \mu^*(B_j),$$

where

$$\lim_{i \rightarrow \infty} \sum_{j=i}^{\infty} \mu^*(B_j) = 0, \text{ provided } \sum_{j=1}^{\infty} \mu^*(B_j) < \infty.$$




Figure 1.10: Exhaustion by separated distance sets.

By the construction $\text{dist}(B_j, B_l) > 0$ whenever $l \geq j + 2$. By the assumption we have

$$\sum_{j=1}^k \mu^*(B_{2j}) = \mu^*\left(\bigcup_{j=1}^k B_{2j}\right) \leq \mu^*(E) < \infty$$

and

$$\sum_{j=0}^k \mu^*(B_{2j+1}) = \mu^*\left(\bigcup_{j=0}^k B_{2j+1}\right) \leq \mu^*(E) < \infty.$$

These together imply

$$\sum_{j=1}^{\infty} \mu^*(B_j) = \lim_{k \rightarrow \infty} \left(\sum_{j=1}^k \mu^*(B_{2j}) + \sum_{j=0}^k \mu^*(B_{2j+1}) \right) \leq 2\mu^*(E) < \infty.$$

Thus

$$\mu^*(A) \leq \lim_{i \rightarrow \infty} \mu^*(A_i)$$

and consequently

$$\lim_{i \rightarrow \infty} \mu^*(A_i) = \mu^*(A). \quad \blacksquare$$

Finally

$$\begin{aligned} \mu^*(E \cap F) + \mu^*(E \setminus F) &= \mu^*(E \cap F) + \mu^*(A) \quad (A = E \setminus F) \\ &= \lim_{i \rightarrow \infty} (\mu^*(E \cap F) + \mu^*(A_i)) \quad (\text{above}) \\ &= \lim_{i \rightarrow \infty} \mu^*((E \cap F) \cup A_i) \quad (\text{dist}(A_i, F) > 0) \\ &\leq \mu^*(E). \quad (A_i \subset E) \quad \square \end{aligned}$$

Remarks 1.54:

- (1) The converse holds as well, so that the previous theorem gives a characterization for a Borel outer measure, see [2, Theorem 3.7]. Observe, that there may be also other measurable sets than Borel sets, because an arbitrary measurable set for a Radon measure can be represented as a union of Borel set and a set of measure zero, see Corollary 1.49.
- (2) The Carathéodory construction in Example 1.4 (8) always gives a metric outer measure. In particular, all Borel sets are measurable. Moreover, if the members of covering family in the construction are Borel sets, then the measure is Borel regular (exercise). Thus many natural constructions give a Borel regular outer measure.
- (3) The Hausdorff measure, defined in Example 1.4 (6), is a metric outer measures (exercise). Thus all Borel sets are measurable with respect to a Hausdorff measure. See also [2, Section 3.8], [4, Chapter 2], [7, Chapter 19], [10, Chapter 7] and [16, Chapter 7].

Lemma 1.55. The Lebesgue outer measure m^* is a metric outer measure.

T H E M O R A L : Theorem 1.53 implies that the Lebesgue outer measure m^* is a Borel outer measure. Thus all Borel sets are m^* -measurable, in particular closed and open sets, are Lebesgue measurable. By Lemma 1.40 and Lemma 1.41 we can conclude that m^* is a Radon outer measure, see Definition 1.37.

Proof. Let $A, B \subset \mathbb{R}^n$ with $\text{dist}(A, B) > 0$. Subadditivity implies that $m^*(A \cup B) \leq m^*(A) + m^*(B)$. Hence it is enough to prove the reverse inequality. We may assume that $m^*(A \cup B) < \infty$. For every $\varepsilon > 0$ there exist intervals I_i , $i = 1, 2, \dots$, such that $A \cup B \subset \bigcup_{i=1}^{\infty} I_i$ and

$$\sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(A \cup B) + \varepsilon.$$

By subdividing each I_i into smaller intervals, we may assume that $\text{diam}(I_i) < \text{dist}(A, B)$ for every $i = 1, 2, \dots$. This can be done, for example, by diving recursively every side of an interval into two equally long parts as long as diameters of all intervals are less than $\text{dist}(A, B)$. Note that the sum of volumes of the intervals remains unchanged in the subdivision. Every such interval I_i intersects at most one of the sets A and B .

THE MORAL: We can assume that the diameter of the intervals in the definition of the Lebesgue measure is as small as we want.

We consider two subfamilies I'_i and I''_i , $i = 1, 2, \dots$, where the intervals I'_i , $i = 1, 2, \dots$, have nonempty intersections with A and the intervals I''_i , $i = 1, 2, \dots$, have nonempty intersections with B . Note that there may be intervals that do not intersect $A \cup B$, but this is not a problem. Thus

$$m^*(A) + m^*(B) \leq \sum_{i=1}^{\infty} \text{vol}(I'_i) + \sum_{i=1}^{\infty} \text{vol}(I''_i) \leq \sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(A \cup B) + \varepsilon. \quad \square$$

By letting $\varepsilon \rightarrow 0$, we obtain $m^*(A \cup B) \geq m^*(A) + m^*(B)$.

1.7 Lebesgue measure revisited

We have already discussed the definition of the Lebesgue outer measure in Example 1.4 (5). Recall that the Lebesgue outer measure of an arbitrary set $A \subset \mathbb{R}^n$ is

$$m^*(A) = \inf \left\{ \sum_{i=1}^{\infty} \text{vol}(I_i) : A \subset \bigcup_{i=1}^{\infty} I_i \right\},$$

where the infimum is taken over all coverings of A by countably many closed intervals I_i , $i = 1, 2, \dots$. We discuss examples and properties that are characteristic for the Lebesgue outer measure. One of the goals is to show that measure theory for the Lebesgue outer measure can be developed directly from the definition.

Remarks 1.56:

- (1) We cannot upgrade countable subadditivity of the Lebesgue outer measure to uncountable subadditivity. For example, \mathbb{R}^n is an uncountable union of points, each of which has Lebesgue outer measure zero, but \mathbb{R}^n has infinite Lebesgue outer measure.




Figure 1.11: Lebesgue measure is a metric measure.

(2) If we consider coverings with finitely many intervals, we obtain the Jordan outer measure defined as

$$m^{*,J}(A) = \inf \left\{ \sum_{i=1}^k \text{vol}(I_i) : A \subset \bigcup_{i=1}^k I_i, k = 1, 2, \dots \right\},$$

where $A \subset \mathbb{R}^n$ is a bounded set. The Jordan outer measure will not be an outer measure, it is only finitely subadditive instead of countably subadditive. It has the property $J^*(A) = J^*(\overline{A})$ for every bounded $A \subset \mathbb{R}^n$. We can define the corresponding Jordan inner measure by

$$m_{*,J}(A) = \sup \left\{ \sum_{i=1}^k \text{vol}(I_i) : A \supset \bigcup_{i=1}^k I_i, k = 1, 2, \dots \right\}$$

and say that a bounded set $A \subset \mathbb{R}^n$ is Jordan measurable if the inner and outer Jordan measures coincide. It can be shown that a bounded set $A \subset \mathbb{R}^n$ is Jordan measurable if and only if the Jordan outer measure of ∂A is zero. For example,

$$m^{*,J}(\mathbb{Q} \cap [0, 1]) = 1 \quad \text{and} \quad m_{*,J}(\mathbb{Q} \cap [0, 1]) = 0,$$

while $m^*(\mathbb{Q} \cap [0, 1]) = 0$, since $\mathbb{Q} \cap [0, 1]$ is a countable set. In particular, $\mathbb{Q} \cap [0, 1]$ is Lebesgue measurable but not Jordan measurable. This example also shows that the Jordan outer measure is not countably additive. See [11] for more on the Jordan outer measure.

Remark 1.57. The closed intervals in the definition of the Lebesgue outer measure can be replaced by open intervals.

Reason. Assume that $m^*(A) < \infty$. Let $\varepsilon > 0$. Let I_i , $i = 1, 2, \dots$, be closed intervals such that $A \subset \bigcup_{i=1}^{\infty} I_i$ and

$$\sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(A) + \frac{\varepsilon}{2}.$$

Let J_i be an open interval containing I_i with

$$\text{vol}(\overline{J_i}) \leq \text{vol}(I_i) + \frac{\varepsilon}{2^{i+1}}$$

for $i = 1, 2, \dots$. Here $\overline{J_i}$ is the closure of J_i , that is, the corresponding closed interval. Then $A \subset \bigcup_{i=1}^{\infty} I_i \subset \bigcup_{i=1}^{\infty} \overline{J_i}$ and

$$\begin{aligned} m^*(A) &\leq \sum_{i=1}^{\infty} \text{vol}(\overline{J_i}) \leq \sum_{i=1}^{\infty} \left(\text{vol}(I_i) + \frac{\varepsilon}{2^i} \right) \leq \sum_{i=1}^{\infty} \text{vol}(I_i) + \frac{\varepsilon}{2} \\ &< m^*(A) + \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = m^*(A) + \varepsilon. \end{aligned}$$

Then we discuss the case $m^*(A) = \infty$. Let J_i , $i = 1, 2, \dots$, be open intervals such that $A \subset \bigcup_{i=1}^{\infty} J_i$. Since $m^*(A) = \infty$, we have $\sum_{i=1}^{\infty} \text{vol}(I_i) = \infty$ for every countable collection closed intervals I_i , $i = 1, 2, \dots$, such that $A \subset \bigcup_{i=1}^{\infty} I_i$. Since $A \subset \bigcup_{i=1}^{\infty} J_i \subset \bigcup_{i=1}^{\infty} \overline{J_i}$, this implies that $\sum_{i=1}^{\infty} \text{vol}(J_i) = \sum_{i=1}^{\infty} \text{vol}(\overline{J_i}) = \infty$. ■

The closed intervals in the definition of the Lebesgue outer measure can also be replaced by cubes. Cubes are intervals whose side lengths are equal, that is $b_1 - a_1 = \dots = b_n - a_n$. Even balls will do, but this is more subtle (exercise).

We begin with some facts on the volumes of intervals, which may look obvious but are somewhat tedious to prove. We say that closed intervals I_i , $i = 1, 2, \dots$, are almost disjoint if their interiors are pairwise disjoint. Interior of a closed interval is the corresponding open interval. Thus almost disjoint intervals may touch at most on their boundaries.

Lemma 1.58. Let I be an interval and assume that $I = \bigcup_{i=1}^k I_i$, where I_i , $i = 1, 2, \dots, k$, are almost disjoint intervals. Then

$$\text{vol}(I) = \sum_{i=1}^k \text{vol}(I_i).$$

T H E M O R A L : The volume of a closed interval is preserved in finite partitions into almost disjoint intervals.

Proof. By extending the sides of intervals I_1, \dots, I_k to hyperplanes, we may decompose I into a collection of almost disjoint intervals J_j , $j = 1, \dots, l$, such that

$$I = \bigcup_{j=1}^l J_j \quad \text{and} \quad I_i = \bigcup_{j \in \mathcal{J}_i} J_j, \quad i = 1, \dots, k.$$




Figure 1.12: The grid formed by the intervals I_i .

Here \mathcal{J}_i , $i = 1, \dots, k$, consists of those indices $j = 1, \dots, l$ for which $J_j \subset I_i$. Since the obtained grid partitions the sides of I and every J_j consists of products of the intervals in these partitions, we have

$$\text{vol}(I) = \sum_{j=1}^l \text{vol}(J_j).$$

Since this also holds for the intervals I_1, \dots, I_k , we obtain

$$\text{vol}(I) = \sum_{j=1}^l \text{vol}(J_j) = \sum_{i=1}^k \sum_{j \in \mathcal{J}_i} \text{vol}(J_j) = \sum_{i=1}^k \text{vol}(I_i).$$

□

Next we discuss a subadditivity result for the volume of coverings of an interval with finitely many possibly overlapping intervals.

Lemma 1.59. Let I be an interval and assume that $I \subset \bigcup_{i=1}^k I_i$, where I_i , $i = 1, 2, \dots, k$, are intervals. Then

$$\text{vol}(I) \leq \sum_{i=1}^k \text{vol}(I_i).$$

Proof. Since the intervals I_i , $i = 1, \dots, k$, cover I , there exist almost disjoint intervals J_j , $j = 1, \dots, l$, such that $I = \bigcup_{j=1}^l J_j$ and

$$\sum_{j=1}^l \text{vol}(J_j) \leq \sum_{i=1}^k \text{vol}(I_i).$$

To obtain the intervals J_j , we replace I_i by the interval $I \cap I_i$ and then decompose these possibly overlapping intervals into an almost disjoint collection of subintervals with the same union. By discarding overlaps the sum of the volumes can only decrease. By Lemma 1.58, we have

$$\text{vol}(I) = \sum_{j=1}^l \text{vol}(J_j) \leq \sum_{i=1}^k \text{vol}(I_i).$$

□

Next we show that the Lebesgue outer measure of a closed interval is equal to its volume. Note that this is not obvious from the definition.

Lemma 1.60. Let $I \subset \mathbb{R}^n$ be a closed interval. Then $m^*(I) = \text{vol}(I)$.

THEOREM: The definition of the Lebesgue outer measure is consistent for closed intervals.

Proof. It is clear that $m^*(I) \leq \text{vol}(I)$, since the interval I itself is an admissible covering the definition of the Lebesgue outer measure. Hence it remains to prove that $\text{vol}(I) \leq m^*(I)$. For every $\varepsilon > 0$ there exist intervals I_i , $i = 1, 2, \dots$, such that $I \subset \bigcup_{i=1}^{\infty} I_i$ and

$$\sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(I) + \varepsilon.$$

For every $i = 1, 2, \dots$ there exists an open interval J_i such that $I_i \subset J_i$ and

$$\text{vol}(\bar{J}_i) \leq \text{vol}(I_i) + \frac{\varepsilon}{2^i}.$$

Here \bar{J}_i is the closure of J_i , that is, the corresponding closed interval. It follows that

$$\sum_{i=1}^{\infty} \text{vol}(\bar{J}_i) \leq \sum_{i=1}^{\infty} \left(\text{vol}(I_i) + \frac{\varepsilon}{2^i} \right) = \sum_{i=1}^{\infty} \text{vol}(I_i) + \sum_{i=1}^{\infty} \frac{\varepsilon}{2^i} = \sum_{i=1}^{\infty} \text{vol}(I_i) + \varepsilon.$$

The collection of intervals J_i , $i = 1, 2, \dots$, is an open covering of the compact set I and thus there exists a finite subcovering J_i , $i = 1, 2, \dots, k$. Then \bar{J}_i , $i = 1, 2, \dots, k$, are closed intervals that cover I and Lemma 1.59 implies

$$\text{vol}(I) \leq \sum_{i=1}^k \text{vol}(\bar{J}_i) \leq \sum_{i=1}^{\infty} \text{vol}(\bar{J}_i) \leq \sum_{i=1}^{\infty} \text{vol}(I_i) + \varepsilon \leq m^*(I) + 2\varepsilon.$$

The claim follows by letting $\varepsilon \rightarrow 0$. □

Remark 1.61. The key point of the previous proof is a countable subadditivity result for the volume of intervals. Let $I \subset \mathbb{R}^n$ be an interval and let I_i , $i = 1, 2, \dots$, be intervals such that $I \subset \bigcup_{i=1}^{\infty} I_i$. Then

$$\text{vol}(I) \leq \sum_{i=1}^{\infty} \text{vol}(I_i).$$

Note that this follows from the countable subadditivity of the Lebesgue measure if we know that Lemma 1.60 is true.

Remark 1.62. The assertion $m^*(I) = \text{vol}(I)$ in Lemma 1.60 holds for an open interval $I \subset \mathbb{R}^n$ as well. Since I is covered by its closure \bar{I} , we have $m^*(I) \leq \text{vol}(\bar{I}) = \text{vol}(I)$. To prove the reverse inequality, let $\varepsilon > 0$ and let J be a closed interval contained in I with $\text{vol}(I) \leq \text{vol}(J) + \varepsilon$. By monotonicity $m^*(J) \leq m^*(I)$ and by Lemma 1.60 we obtain

$$\text{vol}(I) \leq \text{vol}(J) + \varepsilon = m^*(J) + \varepsilon \leq m^*(I) + \varepsilon.$$

By letting $\varepsilon \rightarrow 0$, we have $\text{vol}(I) \leq m^*(I)$.

By Lemma 1.55 and Theorem 1.53 we know that all Borel sets are Lebesgue measurable. However, we discuss a direct argument to show that every closed interval is Lebesgue measurable.

Lemma 1.63. Every closed interval $I \subset \mathbb{R}^n$ is Lebesgue measurable.

T H E M O R A L : Lemma 1.67 below shows that every open set can be decomposed into countably many almost disjoint closed intervals. This implies that open sets are Lebesgue measurable and, consequently, Borel sets are Lebesgue measurable.

Proof. Let I be an interval in \mathbb{R}^n and $E \subset \mathbb{R}^n$. By Remark 1.6 (1), we may assume that $m^*(E) < \infty$. For every $\varepsilon > 0$ there exist intervals I_j , $j = 1, 2, \dots$, such that $E \subset \bigcup_{j=1}^{\infty} I_j$ and

$$\sum_{j=1}^{\infty} \text{vol}(I_j) < m^*(E) + \varepsilon.$$

We decompose every I_j into finitely many almost disjoint intervals $J_j, R_{j,1}, \dots, R_{j,k_j}$ such that

$$I_j = J_j \cup \bigcup_{i=1}^{k_j} R_{j,i}, \quad J_j = I_j \cap I \subset I \quad \text{and} \quad R_{j,i} \subset \bar{I}^c.$$

Here $I^c = \mathbb{R}^n \setminus I$ denotes the complement of I .

Lemma 1.58 implies

$$\text{vol}(I_j) = \text{vol}(J_j) + \sum_{i=1}^{k_j} \text{vol}(R_{j,i})$$

and we obtain

$$m^*(E) + \varepsilon > \sum_{j=1}^{\infty} \text{vol}(I_j) = \sum_{j=1}^{\infty} \text{vol}(J_j) + \sum_{j=1}^{\infty} \sum_{i=1}^{k_j} \text{vol}(R_{j,i}).$$

By relabeling $R_{j,i}$ as R_j we have

$$m^*(E) + \varepsilon > \sum_{j=1}^{\infty} \text{vol}(J_j) + \sum_{j=1}^{\infty} \text{vol}(R_j).$$




Figure 1.13: Decomposition of the covering intervals.

Since the intervals J_j , $j = 1, 2, \dots$, cover $E \cap I$ and the intervals R_j , $j = 1, 2, \dots$, cover $E \cap I^C$, we have

$$m^*(E \cap I) \leq \sum_{j=1}^{\infty} \text{vol}(J_j) \quad \text{and} \quad m^*(E \cap I^C) \leq \sum_{j=1}^{\infty} \text{vol}(R_j).$$

Thus we have

$$m^*(E) + \varepsilon > m^*(E \cap I) + m^*(E \cap I^C).$$

By letting $\varepsilon \rightarrow 0$, we have

$$m^*(E) \geq m^*(E \cap I) + m^*(E \cap I^C) = m^*(E \cap I) + m^*(E \setminus I).$$

This shows that I is Lebesgue measurable, see Definition 1.5. \square

Remark 1.64. An open interval $I = (a_1, b_1) \times \dots \times (a_n, b_n) \subset \mathbb{R}^n$ can be written as $I = \bigcup_{i=1}^{\infty} I_i$, where

$$I_i = \left[a_1 + \frac{1}{i}, b_1 - \frac{1}{i} \right] \times \dots \times \left[a_n + \frac{1}{i}, b_n - \frac{1}{i} \right], \quad i = 1, 2, \dots$$

Since the collection of Lebesgue measurable sets is a σ -algebra, this shows that every open interval is Lebesgue measurable. By Theorem 1.22 and Lemma 1.60

we have

$$\begin{aligned} m^*(I) &= m^*\left(\bigcup_{i=1}^{\infty} I_i\right) = \lim_{i \rightarrow \infty} m^*(I_i) = \lim_{i \rightarrow \infty} \text{vol}(I_i) \\ &= \lim_{i \rightarrow \infty} (b_1 - a_1 - \frac{2}{i}) \cdots (b_n - a_n - \frac{2}{i}) = (b_1 - a_1) \cdots (b_n - a_n) \\ &= \text{vol}(\bar{I}) = m^*(\bar{I}). \end{aligned}$$

Thus the Lebesgue measure of a closed interval and the corresponding open interval coincide. Moreover $\partial I = \bar{I} \setminus I$ and since \bar{I} and I are Lebesgue measurable, by Remark 1.6 (3) we have

$$m^*(\partial I) = m^*(\bar{I} \setminus I) = m^*(\bar{I}) - m^*(I) = 0.$$

Thus the boundary of an interval is a set of Lebesgue measure zero. This can be also proved directly for the definition of the Lebesgue measure by covering ∂I with appropriate intervals whose volumes sum up to less than $\varepsilon > 0$.

Next we discuss the Lebesgue outer measure of countable unions of intervals. Let I_i , $i = 1, 2, \dots$, be pairwise disjoint closed intervals. Since intervals are Lebesgue measurable, we have

$$m^*\left(\bigcup_{i=1}^{\infty} I_i\right) = \sum_{i=1}^{\infty} m^*(I_i).$$

Next we discuss the corresponding result for almost disjoint closed intervals.

Lemma 1.65. If $I_i \subset \mathbb{R}^n$, $i = 1, 2, \dots$ are almost disjoint closed intervals, then

$$m^*\left(\bigcup_{i=1}^{\infty} I_i\right) = \sum_{i=1}^{\infty} m^*(I_i).$$

T H E M O R A L : Lemma 1.67 below shows that every open can be decomposed into countably many almost disjoint closed intervals. Thus its Lebesgue outer measure equals the sum of the volumes of the intervals. Moreover, the sum is independent of the decomposition. \square

Proof. Since the intervals I_i , $i = 1, 2, \dots$, cover $\bigcup_{i=1}^{\infty} I_i$, by the definition of the Lebesgue outer measure we have

$$m^*\left(\bigcup_{i=1}^{\infty} I_i\right) \leq \sum_{i=1}^{\infty} \text{vol}(I_i).$$

Then we prove the reverse inequality. For every $i = 1, 2, \dots$, let J_i be a closed interval contained in I_i with $J_i \cap \partial I_i = \emptyset$ and

$$\text{vol}(I_i) \leq \text{vol}(J_i) + \frac{\varepsilon}{2^i}.$$

For every $k = 1, 2, \dots$, the intervals J_1, \dots, J_k are pairwise disjoint compact sets and thus $\text{dist}(J_i, J_j) > 0$ for $i \neq j$. Since the Lebesgue outer measure is a metric outer measure, see Lemma 1.55, we have

$$\begin{aligned} m^* \left(\bigcup_{i=1}^k J_i \right) &= \sum_{i=1}^k m^*(J_i) = \sum_{i=1}^k \text{vol}(J_i) \\ &\geq \sum_{i=1}^k \left(\text{vol}(I_i) - \frac{\varepsilon}{2^i} \right) \geq \sum_{i=1}^k \text{vol}(I_i) - \varepsilon \end{aligned}$$

Here we also used Lemma 1.60. Since $\bigcup_{i=1}^k J_i \subset \bigcup_{i=1}^{\infty} I_i$, by monotonicity we have

$$m^* \left(\bigcup_{i=1}^{\infty} I_i \right) \geq m^* \left(\bigcup_{i=1}^k J_i \right) \geq \sum_{i=1}^k \text{vol}(I_i) - \varepsilon$$

for every $k = 1, 2, \dots$. By letting $k \rightarrow \infty$, we obtain

$$\sum_{i=1}^{\infty} \text{vol}(I_i) = \lim_{k \rightarrow \infty} \sum_{i=1}^k \text{vol}(I_i) \leq m^* \left(\bigcup_{i=1}^{\infty} I_i \right) + \varepsilon.$$

Finally, by letting $\varepsilon \rightarrow 0$, we have

$$\sum_{i=1}^{\infty} \text{vol}(I_i) \leq m^* \left(\bigcup_{i=1}^{\infty} I_i \right). \quad \square$$

Remark 1.66. The proof above is only based on the definition of the Lebesgue measure and does not apply general results on outer measures. We discuss another proof, which applies countable additivity on pairwise disjoint measurable sets. By countable subadditivity, we have

$$m^* \left(\bigcup_{i=1}^{\infty} I_i \right) \leq \sum_{i=1}^{\infty} m^*(I_i).$$

To prove the reverse inequality, we let J_i be interior of I_i , that is, the corresponding open interval. The open intervals J_i are pairwise disjoint Lebesgue measurable sets with $J_i \subset I_i$, $i = 1, 2, \dots$. By countable additivity on pairwise disjoint Lebesgue measurable sets, we have

$$m^* \left(\bigcup_{i=1}^{\infty} I_i \right) \geq m^* \left(\bigcup_{i=1}^{\infty} J_i \right) = \sum_{i=1}^{\infty} m^*(J_i) = \sum_{i=1}^{\infty} m^*(I_i).$$

Here we also applied Remark 1.64.

In the one-dimensional case every nonempty open set is a union of countably many disjoint open intervals, see [10, Theorem 1.3, p. 6]. By Lemma 1.65 the Lebesgue outer measure of an open set is the sum of volumes of these intervals. Next we consider this question in the higher dimensional case.

A half-open dyadic cube is of the form

$$\left[\frac{i_1}{2^k}, \frac{i_1+1}{2^k} \right) \times \cdots \times \left[\frac{i_n}{2^k}, \frac{i_n+1}{2^k} \right), \quad i_1, \dots, i_n \in \mathbb{Z}, \quad k \in \mathbb{Z}.$$

The collection of dyadic cubes \mathcal{D}_k , $k \in \mathbb{Z}$, consists of the dyadic cubes with the side length 2^{-k} . The collection of all dyadic cubes in \mathbb{R}^n is

$$\mathcal{D} = \bigcup_{k \in \mathbb{Z}} \mathcal{D}_k.$$

Observe that \mathcal{D}_k consist of cubes whose vertices lie on the lattice $2^{-k} \mathbb{Z}^n$ and whose side length is 2^{-k} . The dyadic cubes in the k th generation can be defined as $\mathcal{D}_k = 2^{-k}([0, 1]^n + \mathbb{Z}^n)$, $k \in \mathbb{Z}$. The cubes in \mathcal{D}_k cover the whole \mathbb{R}^n and are pairwise disjoint. Dyadic cubes have a very useful nesting property which states that any




Figure 1.14: Dyadic cubes.

two dyadic cubes are either disjoint or one of them is contained in the other.

Lemma 1.67. Every nonempty open set G in \mathbb{R}^n is a union of countably many pairwise disjoint dyadic cubes.

T H E M O R A L : Since dyadic cubes are Lebesgue measurable by Lemma 1.63 every nonempty open set is Lebesgue measurable as a countable union of dyadic cubes. Moreover, the Lebesgue measure of an open set can be computed as a sum of the volumes of the related dyadic cubes by Lemma 1.65. Since the collection of Lebesgue measurable sets is a σ -algebra, we conclude that all Borel sets are Lebesgue measurable.




Figure 1.15: The nesting property of the dyadic cubes.

Proof. Consider dyadic cubes in \mathcal{D}_1 that are contained in G and denote

$$\mathcal{D}_1 = \{Q \in \mathcal{D}_1 : Q \subset G\}.$$

Then consider dyadic cubes in \mathcal{D}_2 that are contained in G and do not intersect any of the cubes in \mathcal{D}_1 and denote

$$\mathcal{D}_2 = \{Q \in \mathcal{D}_2 : Q \subset G, Q \cap J = \emptyset \text{ for every } J \in \mathcal{D}_1\}.$$

Recursively define

$$\mathcal{D}_k = \left\{ Q \in \mathcal{D}_k : Q \subset G, Q \cap J = \emptyset \text{ for every } J \in \bigcup_{i=1}^{k-1} \mathcal{D}_i \right\}$$

for every $k = 2, 3, \dots$. Then $\mathcal{D} = \bigcup_{k=1}^{\infty} \mathcal{D}_k$ is a countable collection of pairwise disjoint cubes.

Claim: $G = \bigcup_{Q \in \mathcal{D}} Q$.

Reason. It is clear from the construction that $\bigcup_{Q \in \mathcal{D}} Q \subset G$. For the reverse inclusion, let $x \in G$. Let k be so large that the common diameter of the cubes in \mathcal{D}_k is smaller than r , that is, $\sqrt{n}2^{-k} < r$. Since G is open, there exists a ball $B(x, r) \subset G$ with $r > 0$. Since the dyadic \mathcal{D}_k cubes cover \mathbb{R}^n , there exists a dyadic cube $Q \in \mathcal{D}_k$ with $x \in Q$ and $Q \subset B(x, r) \subset G$. There are two possibilities $Q \in \mathcal{D}_k$ or $Q \notin \mathcal{D}_k$. If $Q \in \mathcal{D}_k$, then $x \in Q \subset \bigcup_{Q \in \mathcal{D}} Q$. If $Q \notin \mathcal{D}_k$, there exists $J \in \bigcup_{i=1}^{k-1} \mathcal{D}_i$ with $J \cap Q \neq \emptyset$. The nesting property of dyadic cubes implies $Q \subset J$ and $x \in Q \subset J \subset \bigcup_{Q \in \mathcal{D}} Q$. ■




Figure 1.16: Dyadic partition.

The argument above shows that every nonempty open set is a union of countably many almost disjoint closed cubes

$$\left[\frac{i_1}{2^k}, \frac{i_1+1}{2^k} \right] \times \cdots \times \left[\frac{i_n}{2^k}, \frac{i_n+1}{2^k} \right], \quad i_1, \dots, i_n \in \mathbb{Z}, \quad k \in \mathbb{Z}.$$

In most of the cases we are not interested in the precise value of the Lebesgue outer measure of a set $A \subset \mathbb{R}^n$. Instead, it is enough to know whether $m^*(A) = 0$, $0 < m^*(A) < \infty$ or $m^*(A) = \infty$, see Examples 1.4 (5). It follows directly from the definition of the Lebesgue measure that a set $A \subset \mathbb{R}^n$ is of Lebesgue outer measure zero if and only if for every $\varepsilon > 0$ there exist intervals I_i , $i = 1, 2, \dots$, such that

$$A \subset \bigcup_{i=1}^{\infty} I_i \quad \text{and} \quad \sum_{i=1}^{\infty} \text{vol}(I_i) < \varepsilon.$$

Observe that we do not need measure theory in order to be able to define sets of Lebesgue measure zero.

T H E M O R A L : A set is of Lebesgue measure zero if it can be covered by intervals such that the sum of the volumes of the intervals is arbitrarily small.

Examples 1.68:

(1) Any one point set is of Lebesgue measure zero, that is, $m^*(\{x\}) = 0$ for every $x = (x_1, \dots, x_n) \in \mathbb{R}^n$. We give two ways to prove the claim.

(1) Let $\varepsilon > 0$ and

$$Q = \left[x_1 - \frac{\varepsilon^{\frac{1}{n}}}{2}, x_1 + \frac{\varepsilon^{\frac{1}{n}}}{2} \right] \times \cdots \times \left[x_n - \frac{\varepsilon^{\frac{1}{n}}}{2}, x_n + \frac{\varepsilon^{\frac{1}{n}}}{2} \right].$$

Observe that Q is a cube with center x and all side lengths equal to $\varepsilon^{\frac{1}{n}}$.

Then

$$m^*(\{x\}) \leq \text{vol}(Q) = \varepsilon,$$

which implies that $m^*(\{x\}) = 0$.

(2) We can cover $\{x\}$ by the degenerate interval $[x_1, x_1] \times \cdots \times [x_n, x_n]$ with zero volume and conclude the claim from this.

(2) Any countable set $A = \{x_1, x_2, \dots\}$, $x_i \in \mathbb{R}^n$, is of Lebesgue measure zero. We give two ways to prove the claim.

(1) Let $\varepsilon > 0$ and Q_i , $i = 1, 2, \dots$, be a closed n -dimensional cube with center x_i and side length $(\frac{\varepsilon}{2^i})^{\frac{1}{n}}$. Then

$$m^*(A) \leq \sum_{i=1}^{\infty} \text{vol}(Q_i) \leq \sum_{i=1}^{\infty} \frac{\varepsilon}{2^i} = \varepsilon,$$

which implies that $m^*(A) = 0$.

(2) By subadditivity

$$m^*(A) = m^*\left(\bigcup_{i=1}^{\infty} \{x_i\}\right) \leq \sum_{i=1}^{\infty} m^*(\{x_i\}) = 0.$$

(3) Let $A = \{x = (x_1, x_2) \in \mathbb{R}^2 : x_2 = 0\} \subset \mathbb{R}^2$. Then the 2-dimensional Lebesgue measure of A is zero.

Reason. Let $A_i = \{x = (x_1, x_2) \in \mathbb{R}^2 : i \leq x_1 < i+1, x_2 = 0\}$, $i \in \mathbb{Z}$. Then $A = \bigcup_{i \in \mathbb{Z}} A_i$. Let $\varepsilon > 0$ and $I = [i, i+1] \times [-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}]$. Then $A_i \subset I$ and $\text{vol}(I) = \varepsilon$. This implies $m^*(A_i) = 0$ and

$$m^*(A) \leq \sum_{i \in \mathbb{Z}} m^*(A_i) = 0. \quad \blacksquare$$

(4) $m^*(\mathbb{R}^n) = \infty$.

Reason. Let I_i , $i = 1, 2, \dots$, be a collection of closed intervals such that $\mathbb{R}^n \subset \bigcup_{i=1}^{\infty} I_i$. Consider the cubes $Q_j = [-j, j]^n = [-j, j] \times \cdots \times [-j, j]$, $j = 1, 2, \dots$. Then $Q_j \subset \bigcup_{i=1}^{\infty} I_i$ and by Lemma 1.60 the Lebesgue outer measure of a closed interval coincides with its volume. Thus we have

$$(2j)^n = \text{vol}(Q_j) = m^*(Q_j) \leq \sum_{i=1}^{\infty} \text{vol}(I_i).$$

Letting $j \rightarrow \infty$, we see that $\sum_{i=1}^{\infty} \text{vol}(I_i) = \infty$ for every covering. This implies that $m^*(\mathbb{R}^n) = \infty$. \blacksquare

(5) Every nonempty open set has positive Lebesgue outer measure.

Reason. Let $G \subset \mathbb{R}^n$ be open. Then for every $x \in G$, there exists a ball $B(x, r) \subset G$ with $r > 0$. The ball $B(x, r)$ contains the cube Q with the center

x and diameter $\frac{r}{2}$. On the other hand, the $\text{diam}(Q) = \sqrt{n}l(Q)$, where $l(Q)$ is the side length of Q . From this we conclude that $l(Q) = r/(2\sqrt{n})$ and thus

$$Q = \left[x_1 - \frac{r}{4\sqrt{n}}, x_1 + \frac{r}{4\sqrt{n}} \right] \times \cdots \times \left[x_n - \frac{r}{4\sqrt{n}}, x_n + \frac{r}{4\sqrt{n}} \right].$$

By Lemma 1.60 this implies

$$m^*(G) \geq m^*(Q) = \text{vol}(Q) = \left(\frac{r}{2\sqrt{n}} \right)^n > 0. \quad \blacksquare$$

This argument is based on the fact that every nonempty open set contains a ball. Observe that a general Lebesgue measurable set of positive Lebesgue measure does not necessarily contain a ball. For example, the set of irrational numbers contained in $(0, 1)$ has Lebesgue measure one, but this set does not contain an open interval.




Figure 1.17: A cube inside a ball.

Observe, that every nonempty open set contains uncountably many points, since all countable sets have Lebesgue measure zero.

1.8 Invariance properties of the Lebesgue measure

The following invariance properties of the Lebesgue measure follow from the corresponding properties of the volume of an interval.

(1) (Translation invariance) Let $A \subset \mathbb{R}^n$, $x_0 \in \mathbb{R}^n$ and denote $A + x_0 = \{x + x_0 \in \mathbb{R}^n : x \in A\}$. Then

$$m^*(A + x_0) = m^*(A).$$

This means that the Lebesgue outer measure is invariant in translations.

Reason. Intervals are mapped to intervals in translations and

$$A \subset \bigcup_{i=1}^{\infty} I_i \iff A + x_0 \subset \bigcup_{i=1}^{\infty} (I_i + x_0).$$

Clearly $\text{vol}(I_i) = \text{vol}(I_i + x_0)$, $i = 1, 2, \dots$, and thus

$$\begin{aligned} m^*(A + x_0) &= \inf \left\{ \sum_{i=1}^{\infty} \text{vol}(I_i + x_0) : A + x_0 \subset \bigcup_{i=1}^{\infty} (I_i + x_0) \right\} \\ &= \inf \left\{ \sum_{i=1}^{\infty} \text{vol}(I_i) : A \subset \bigcup_{i=1}^{\infty} I_i \right\} = m^*(A). \quad \blacksquare \end{aligned}$$

Moreover, A is Lebesgue measurable if and only if $A + x_0$ is Lebesgue measurable.

Reason. Assume that A is Lebesgue measurable. Then

$$\begin{aligned} m^*(E \cap (A + x_0)) + m^*(E \setminus (A + x_0)) \\ &= m^*((E - x_0) \cap A) + x_0 + m^*((E - x_0) \setminus A) + x_0 \\ &= m^*((E - x_0) \cap A) + m^*((E - x_0) \setminus A) \quad (\text{translation invariance}) \\ &= m^*(E - x_0) \quad (A \text{ is measurable}) \\ &= m^*(E) \quad (\text{translation invariance}) \end{aligned}$$

for every $E \subset \mathbb{R}^n$. This shows that $A + x_0$ is Lebesgue measurable. The equivalence follows from this. This claim can also be proved using Theorem 1.48 or Theorem 1.73. \blacksquare

(2) (Reflection invariance) Let $A \subset \mathbb{R}^n$ and denote $-A = \{-x \in \mathbb{R}^n : x \in A\}$. Then

$$m^*(-A) = m^*(A).$$

This means that the Lebesgue outer measure is invariant in reflections.

(3) (Scaling property) Let $A \subset \mathbb{R}^n$, $\delta \geq 0$ and denote $\delta A = \{\delta x \in \mathbb{R}^n : x \in A\}$. Then

$$m^*(\delta A) = \delta^n m^*(A).$$

This shows that the Lebesgue outer measure behaves as a volume is expected in dilations.

(4) (Change of variables) Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a general linear mapping. Then

$$m^*(L(A)) = |\det L| m^*(A).$$

This is a change of variables formula, see [6] pages 65–80 or [16] pages 612–619. Moreover, if A is Lebesgue measurable, then $L(A)$ is Lebesgue measurable. However, if $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $m < n$, then $L(A)$ need not be Lebesgue measurable. We shall return to this question later.

(5) (Rotation invariance) A rotation is a linear mapping $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with $LL^* = I$, where L^* is the transpose of L and I is the identity mapping. Since $\det L = \det L^*$ it follows that $|\det L| = 1$. The change of variables formula implies that

$$m^*(L(A)) = m^*(A)$$

and thus the Lebesgue outer measure is invariant in rotations. This also shows that the Lebesgue outer measure is invariant in orthogonal linear mappings $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$. Recall that L is orthogonal, if $L^{-1} = L^*$. Moreover, the Lebesgue outer measure is invariant under rigid motions $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\Phi(x) = x_0 + Lx$, where L is orthogonal.

T H E M O R A L : The Lebesgue measure is invariant in rigid motions and is consistent with scalings. In Remark 1.76 we shall see that the Lebesgue measure is essentially the only measure with these properties.

Example 1.69. Let $B(x, r) = \{y \in \mathbb{R}^n : |y - x| < r\}$ be a ball with the center $x \in \mathbb{R}^n$ and radius $r > 0$. By the translation invariance

$$m(B(x, r)) = m(B(x', r)) \quad \text{for every } x' \in \mathbb{R}^n$$

and by the scaling property

$$m(B(x, ar)) = a^n m(B(x, r)) \quad \text{for every } a > 0.$$

In particular, $m(B(x, r)) = r^n m(B(0, 1))$ for every $r > 0$. Thus the Lebesgue measure of any ball is uniquely determined by the measure of the unit ball. This question will be discussed further in Example 3.36.

1.9 Lebesgue measurable sets

Next we discuss measurable sets for the Lebesgue outer measure. We have already shown that the Lebesgue outer measure is a Radon outer measure, see the discussion after Lemma 1.55. In particular, all Borel sets are Lebesgue measurable, see also Lemma 1.67.

T H E M O R A L : Open and closed sets are Lebesgue measurable and all sets obtained from these sets by countably many set theoretic operations, as complements intersections and unions, are Lebesgue measurable sets. Thus the majority of sets that we actually encounter in real analysis will be Lebesgue measurable.

However, there exist sets which are not Lebesgue measurable, as we shall see soon.

We begin with discussing how much does a Lebesgue measurable set of finite measure differ from a finite union of intervals. For a compact set it is enough to consider finite coverings in the definition of the Lebesgue measure, see the proof of Lemma 1.60. For a more general Lebesgue measurable set, we apply the symmetric difference of sets A and B defined as

$$A \Delta B = (A \setminus B) \cup (B \setminus A).$$

Remark 1.70. If $A \subset \mathbb{R}^n$ is a Lebesgue measurable set and $B \subset \mathbb{R}^n$ is a set with $m^*(A \Delta B) = 0$, then B is Lebesgue measurable.

Reason. Since $A \Delta B = (A \setminus B) \cup (B \setminus A)$, by monotonicity, we have

$$m^*(A \setminus B) \leq m^*(A \Delta B) = 0 \quad \text{and} \quad m^*(B \setminus A) \leq m^*(A \Delta B) = 0.$$

By Remark 1.6 (4), we conclude that $A \setminus B$ and $B \setminus A$ are Lebesgue measurable. Since $A \cap B = A \setminus (A \setminus B)$, we conclude that $A \cap B$ is Lebesgue measurable. Since $B = (A \cap B) \cup (B \setminus A)$, we conclude that B is Lebesgue measurable. Note that this holds true for general outer measures as well. ■

Theorem 1.71. Let $A \subset \mathbb{R}^n$ be a Lebesgue measurable set with $m^*(A) < \infty$. Then for every $\varepsilon > 0$, there exists a finite union $B = \bigcup_{i=1}^k I_i$ of closed intervals such that $m^*(A \Delta B) < \varepsilon$.

T H E M O R A L : A Lebesgue measurable set with finite measure differs from a finite union of intervals by a set of arbitrarily small measure.

Proof. Let $\varepsilon > 0$. Let I_i , $i = 1, 2, \dots$, be intervals such that $A \subset \bigcup_{i=1}^{\infty} I_i$ and

$$\sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(A) + \frac{\varepsilon}{2}.$$

Since $m^*(A) < \infty$ the series on the left-hand side converges and thus there exists k such that

$$\sum_{i=k+1}^{\infty} \text{vol}(I_i) < \frac{\varepsilon}{2}.$$

Let $B = \bigcup_{i=1}^k I_i$. Since A and B are Lebesgue measurable and $(A \setminus B) \cap (B \setminus A) = \emptyset$,

we have

$$\begin{aligned}
m^*(A \Delta B) &= m^*((A \setminus B) \cup (B \setminus A)) \\
&= m^*(A \setminus B) + m^*(B \setminus A) \\
&\leq m^*\left(\bigcup_{i=k+1}^{\infty} I_i\right) + m^*\left(\bigcup_{i=1}^k I_i \setminus A\right) \\
&\leq m^*\left(\bigcup_{i=k+1}^{\infty} I_i\right) + m^*\left(\bigcup_{i=1}^{\infty} I_i \setminus A\right) \\
&= m^*\left(\bigcup_{i=k+1}^{\infty} I_i\right) + m^*\left(\bigcup_{i=1}^{\infty} I_i\right) - m^*(A) \\
&\leq \sum_{i=k+1}^{\infty} \text{vol}(I_i) + \sum_{i=1}^{\infty} \text{vol}(I_i) - m^*(A) < \varepsilon. \quad \square
\end{aligned}$$

Remark 1.72. The previous result does not hold without the assumption $m^*(A) < \infty$. We discuss a one-dimensional example. Let $A = \bigcup_{i=1}^{\infty} A_i$, where $A_i = (i - \frac{1}{2}, i) \subset \mathbb{R}$, $i = 1, 2, \dots$. The set A is open and thus Lebesgue measurable as a countable union of open intervals A_i . Since the sets A_i , $i = 1, 2, \dots$, are pairwise disjoint, we have

$$m^*(A) = \sum_{i=1}^{\infty} m^*(A_i) = \sum_{i=1}^{\infty} \frac{1}{2} = \infty.$$

Let I_j , $j = 1, \dots, k$, be an arbitrary finite collection of closed intervals in \mathbb{R} . Then $\bigcup_{j=1}^k I_j$ is a bounded set and there exists $m \in \mathbb{N}$ such that $\bigcup_{j=1}^k I_j \subset (-m, m)$. It follows that

$$A \Delta \bigcup_{j=1}^k I_j \supset A \setminus \bigcup_{j=1}^k I_j \supset A \setminus (-m, m) = \bigcup_{i=m+1}^{\infty} A_i.$$

Thus we have

$$m^*\left(A \Delta \bigcup_{j=1}^k I_j\right) \geq m^*\left(\bigcup_{i=m+1}^{\infty} A_i\right) = \sum_{i=m+1}^{\infty} m^*(A_i) = \sum_{i=m+1}^{\infty} \frac{1}{2} = \infty.$$

We revisit approximation properties of Lebesgue measurable sets that are already known from Theorem 1.48 and Corollary 1.51. Certain arguments are easier for the Lebesgue outer measure than for a general Radon outer measure.

Theorem 1.73. If $A \subset \mathbb{R}^n$ is Lebesgue measurable, then the following claims are true.

- (1) For every $\varepsilon > 0$, there exists an open set $G \supset A$ such that $m^*(G \setminus A) < \varepsilon$.
- (2) For every $\varepsilon > 0$, there exists a closed set $F \subset A$ such that $m^*(A \setminus F) < \varepsilon$.
- (3) If $m^*(A) < \infty$, for every $\varepsilon > 0$, there exists a compact set $K \subset A$ such that $m^*(A \setminus K) < \varepsilon$.
- (4) $m^*(A) = \inf\{m^*(G) : A \subset G, G \text{ open}\}$. This holds for every $A \subset \mathbb{R}^n$.
- (5) $m^*(A) = \sup\{m^*(K) : K \subset A, K \text{ compact}\}$.

Proof. (1) Assume that $m^*(A) < \infty$. Let $\varepsilon > 0$. Let I_i , $i = 1, 2, \dots$, be open intervals such that $A \subset \bigcup_{i=1}^{\infty} I_i$ and

$$\sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(A) + \varepsilon,$$

see Remark 1.57. Let $G = \bigcup_{i=1}^{\infty} I_i$. Then G is an open set and we have

$$m^*(G) \leq \sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(A) + \varepsilon.$$

Since A is Lebesgue measurable and $A \subset G$, as in Remark 1.6 (3), we have

$$m^*(G) = m^*(G \cap A) + m^*(G \setminus A) = m^*(A) + m^*(G \setminus A).$$

This also follows from additivity on pairwise disjoint measurable sets. Since $m^*(A) < \infty$, we obtain

$$m^*(G \setminus A) = m^*(G) - m^*(A) < \varepsilon.$$

In the case $m^*(A) = \infty$ we consider the exhaustion $A = \bigcup_{i=1}^{\infty} (A \cap B(0, i))$ as in the proof of Theorem 1.48.

(2) Let $\varepsilon > 0$. By Lemma 1.11 the set $\mathbb{R}^n \setminus A$ is measurable and by (1) there exists an open set $G \supset \mathbb{R}^n \setminus A$ with $m^*(G \setminus (\mathbb{R}^n \setminus A)) < \varepsilon$. Let $F = \mathbb{R}^n \setminus G$. Then F is closed, $F \subset A$, $A \setminus F = G \setminus (\mathbb{R}^n \setminus A)$ and

$$m^*(A \setminus F) = m^*(G \setminus (\mathbb{R}^n \setminus A)) < \varepsilon.$$

(3) Let $\varepsilon > 0$. By (2) there is a closed set $F \subset A$ such that $m^*(A \setminus F) < \varepsilon$. Consider the closed balls centered at the origin $\overline{B(0, i)} = \{x \in \mathbb{R}^n : |x| \leq i\}$ and let $K_i = F \cap \overline{B(0, i)}$, $i = 1, 2, \dots$. The sets K_i , $i = 1, 2, \dots$, are compact, since they are closed and bounded. Then $A \setminus K_i$, $i = 1, 2, \dots$, is a decreasing sequence of measurable sets with $\bigcap_{i=1}^{\infty} (A \setminus K_i) = A \setminus F$. Since $m^*(A) < \infty$, Theorem 1.22 (2) implies that

$$\lim_{i \rightarrow \infty} m^*(A \setminus K_i) = m^*\left(\bigcap_{i=1}^{\infty} (A \setminus K_i)\right) = m^*(A \setminus F) < \varepsilon.$$

It follows that $m^*(A \setminus K_i) < \varepsilon$ for large enough i .

(4) Let $\varepsilon > 0$. Let I_i , $i = 1, 2, \dots$, be open intervals such that $A \subset \bigcup_{i=1}^{\infty} I_i$ and

$$\sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(A) + \varepsilon,$$

see Remark 1.57. Let $G = \bigcup_{i=1}^{\infty} I_i$. Then $G \supset A$ and

$$m^*(G) \leq \sum_{i=1}^{\infty} \text{vol}(I_i) < m^*(A) + \varepsilon.$$

(5) The claim follows as in the proof of Corollary 1.51. \square

The following characterization of Lebesgue measurable sets is a reformulation of Corollary 1.49 and Remark 1.50.

Corollary 1.74. The following claims are equivalent for a set $A \subset \mathbb{R}^n$.

- (1) A is Lebesgue measurable.
- (2) For every $\varepsilon > 0$, there exists an open set $G \supset A$ such that $m^*(G \setminus A) < \varepsilon$.
- (3) For every $\varepsilon > 0$, there exists a closed set $F \subset A$ such that $m^*(A \setminus F) < \varepsilon$.
- (4) There exists a G_δ set G such that $A \subset G$ and $m^*(G \setminus A) = 0$.
- (5) There exists an F_σ set F such that $F \subset A$ and $m^*(A \setminus F) = 0$.

Do there exist Lebesgue measurable sets that are not Borel sets? We shall see in Section 2.3, that

- (1) there are Lebesgue measurable sets that are not Borel sets and
- (2) the restriction of the Lebesgue measure to the Borel sets is not a complete measure, see Definition 1.15.

Lebesgue measurable sets arise as a completion of the σ -algebra of Borel sets, that is, adding all sets of measure zero as in Remark 1.17.

Definition 1.75. The Lebesgue measure is defined to be the Lebesgue outer measure on the σ -algebra of Lebesgue measurable sets. We denote the Lebesgue measure by m . In particular, the Lebesgue measure is countably additive on pairwise disjoint Lebesgue measurable sets.

Remark 1.76. The Lebesgue measure is unique in the sense that it is the only measure on the σ -algebra of Lebesgue measurable sets satisfying the following conditions.

- (1) (Translation invariance) If $A \subset \mathbb{R}^n$ is a Lebesgue measurable set and $x \in \mathbb{R}^n$, then $\mu(A + x) = \mu(A)$.
- (2) (Normalization) $\mu([0, 1]^n) = 1$.

Reason. Subdivide the half open unit cube $Q = [0, 1]^n$ to a union of 2^{kn} pairwise disjoint half open dyadic intervals Q_i of side length 2^{-k} , $k = 1, 2, \dots$. By translation invariance all cubes Q_i have the same measure, that is, $m(Q_i) = m(Q_j)$ for $i, j = 1, 2, \dots$, and by countable additivity on pairwise disjoint measurable sets the sum of their measures equals the measure of the entire cube Q which, by normalization, has measure 1. This implies

$$2^{kn} m(Q_i) = \sum_{i=1}^{2^{kn}} m(Q_i) = m(Q) = 1 = \mu(Q) = \sum_{i=1}^{2^{kn}} \mu(Q_i) = 2^{kn} \mu(Q_i).$$

A similar argument can be done for $k = 0, -1, -2, \dots$ and thus $\mu(Q) = m(Q)$ for all dyadic cubes $Q \subset \mathbb{R}^n$. By Lemma 1.67 every open set can be represented as a

union of pairwise disjoint half open dyadic cubes, countable additivity on pairwise disjoint measurable sets implies that $\mu(G) = m(G)$ for all open sets $G \subset \mathbb{R}^n$. Let $A \subset \mathbb{R}^n$ be a Lebesgue measurable set and $G \subset \mathbb{R}^n$ be an open set with $A \subset G$. Then $\mu(A) \leq \mu(G) = m(G)$ and by Theorem 1.73 (4) we have

$$\mu(A) \leq \inf\{m(G) : A \subset G, G \text{ open}\} = m(A).$$

To show the reverse inequality, we first assume that $A \subset \mathbb{R}^n$ is a bounded Lebesgue measurable set and that $G \subset \mathbb{R}^n$ is a bounded open set with $A \subset G$. By the inequality above, we have $\mu(A) \leq m(A)$ and $\mu(G \setminus A) \leq m(G \setminus A)$ and thus

$$\mu(G) = \mu(A) + \mu(G \setminus A) \leq m(A) + m(G \setminus A) = m(G).$$

Since $\mu(G) = m(G)$, we have

$$m(A) + m(G \setminus A) = \mu(A) + \mu(G \setminus A) \leq \mu(A) + m(G \setminus A)$$

and thus $m(A) \leq \mu(A)$. This implies that $\mu(A) = m(A)$ for all bounded Lebesgue measurable sets $A \subset \mathbb{R}^n$. Finally, we observe that an arbitrary Lebesgue measurable set $A \subset \mathbb{R}^n$ can be represented as a countable union of pairwise disjoint bounded Lebesgue measurable sets $A_i = A \cap (B(0, i+1) \setminus B(0, i))$, $i = 1, 2, \dots$. Thus we obtain

$$\mu(A) = \mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu(A_i) = \sum_{i=1}^{\infty} m(A_i) = m\left(\bigcup_{i=1}^{\infty} A_i\right) = m(A). \quad \blacksquare$$

1.10 A nonmeasurable set

The Lebesgue outer measure m^* on \mathbb{R}^n measuring the n -dimensional volume of subsets of \mathbb{R}^n would ideally have the following properties:

- (1) $m^*(A)$ is defined for every set $A \subset \mathbb{R}^n$,
- (2) m^* is an outer measure,
- (3) m^* is countably additive: $m^*(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} m^*(A_i)$ for pairwise disjoint sets $A_i \subset \mathbb{R}^n$, $i = 1, 2, \dots$
- (4) m^* is translation invariant: $m^*(A) = m^*(A + x)$ for $x \in \mathbb{R}^n$

However, it is impossible to satisfy all these are simultaneously. The Lebesgue outer measure satisfies (1), (2) and (4). The Lebesgue outer measure also satisfies (3), if the sets A_i , $i = 1, 2, \dots$, are Lebesgue measurable, but additivity breaks down for nonmeasurable sets. We have shown that Borel sets are Lebesgue measurable, but we have not yet ruled out the possibility that all sets are Lebesgue measurable. Next we shall show that there exists a nonmeasurable set for the Lebesgue outer measure on \mathbb{R} . Such a set can be constructed by using the axiom of choice.

Remark 1.77. The axiom of choice states that if E is a set and $\{E_\alpha\}$ is a collection of nonempty subsets of E , then there exists a function (a choice function) $\alpha \mapsto x_\alpha$ such that $x_\alpha \in E_\alpha$ for every α . The indexing set of α 's is not assumed to be countable.

T H E M O R A L : The axiom of choice states that we have a set which contains exactly one point from each set in an uncountable collection of sets.

Theorem 1.78. There exists a set $E \subset [0, 1]$ which is not Lebesgue measurable.

T H E M O R A L : It is not possible to define the Lebesgue measure of all subsets in a geometrically reasonable way.

S T R A T E G Y : We show that there exists a set $E \subset [0, 1]$ of positive Lebesgue outer measure such that the translated sets $E + q$, $q \in [-1, 1] \cap \mathbb{Q}$, form a pairwise disjoint covering of the interval $[0, 1]$. As the Lebesgue measure is translation invariant and countably additive on measurable sets, the set E cannot be Lebesgue measurable.

Proof. Define an equivalence relation on \mathbb{R} by

$$x \sim y \iff x - y \in \mathbb{Q}.$$

Claim: \sim is an equivalence relation.

Reason. It is clear that $x \sim x$ and that if $x \sim y$ then $y \sim x$. To prove transitivity, assume that $x \sim y$ and $y \sim z$. Then $x - y = q_1$ and $y - z = q_2$, where $q_1, q_2 \in \mathbb{Q}$ and

$$x - z = (x - y) + (y - z) = q_1 + q_2 \in \mathbb{Q}.$$

This implies that $x \sim z$. ■

Denote the equivalence class containing a point $x \in \mathbb{R}$ by E_x . Note that $E_x = x + \mathbb{Q}$, that is, every equivalence class is a translate of \mathbb{Q} . Note also that if $x \in \mathbb{Q}$, then $E_x = \mathbb{Q}$. A crucial property of equivalence classes is that if $x, y \in \mathbb{R}$, then $E_x = E_y$ or $E_x \cap E_y = \emptyset$. That is, two equivalence classes either coincide or are disjoint. The equivalence relation \sim decomposes \mathbb{R} into disjoint equivalence classes. In other words, \mathbb{R} is a union of pairwise disjoint translates of \mathbb{Q} . Note that each equivalence class is countable and, since \mathbb{R} is uncountable, there must be uncountably many equivalence classes. Each equivalence class is dense in \mathbb{R} and has a nonempty intersection with $[0, 1]$. By the axiom of choice, there exists a set E which consists of precisely one element of each equivalence class belonging to $[0, 1]$. If x and y are arbitrary points of E , then $x - y$ is an irrational number, for otherwise they would belong to the same equivalence class, contrary to the definition of E .

We claim that E is not Lebesgue measurable. Assume for a contradiction that E is Lebesgue measurable. Then the translated sets

$$E + q = \{x + q : x \in E\}, \quad q \in \mathbb{Q},$$

are Lebesgue measurable.

Claim: The sets $E + q$, $q \in \mathbb{Q}$, are pairwise disjoint, that is,

$$(E + q) \cap (E + r) = \emptyset \quad \text{whenever} \quad q, r \in \mathbb{Q}, q \neq r.$$

Reason. For a contradiction, assume that $y \in (E + q) \cap (E + r)$ with $q \neq r$. Then $y = x + q$ and $y = z + r$ for some $x, z \in E$. Thus

$$x - z = (y - q) - (y - r) = r - q \in \mathbb{Q},$$

which implies that $x \sim z$. Since E contains exactly one element of each equivalence class, we have $x = z$ and consequently $r = q$. \blacksquare

Claim: $[0, 1] \subset \bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q) \subset [-1, 2]$.

Reason. Let $x \in [0, 1]$ and let y be the representative of the equivalence class E_x belonging to E . In particular, $x \sim y$ from which it follows that $x - y \in \mathbb{Q}$. Let $q = x - y$. Since $x, y \in [0, 1]$ we have $q \in [-1, 1]$ and $x = y + q \in E + q$. This proves the first inclusion. The second inclusion is clear. \blacksquare

Since the sets $E + q$, $q \in [-1, 1] \cap \mathbb{Q}$, are pairwise disjoint and Lebesgue measurable, by countable additivity and translation invariance,

$$m^* \left(\bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q) \right) = \sum_{q \in [-1, 1] \cap \mathbb{Q}} m^*(E + q) = \sum_{q \in [-1, 1] \cap \mathbb{Q}} m^*(E),$$

which is 0 if $m^*(E) = 0$ and ∞ if $m^*(E) > 0$. On the other hand, since $[0, 1] \subset \bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q)$, by monotonicity

$$\sum_{q \in [-1, 1] \cap \mathbb{Q}} m^*(E) = m^* \left(\bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q) \right) \geq m^*([0, 1]) = 1 > 0,$$

which implies that $m^*(E) > 0$ and, consequently,

$$m^* \left(\bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q) \right) = \infty.$$

Since $\bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q) \subset [-1, 2]$, by monotonicity

$$\infty = m^* \left(\bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q) \right) \leq m^*([-1, 2]) = 3.$$

This is a contradiction and thus E cannot be Lebesgue measurable. \square

Remark 1.79. The proof shows that $E \subset [0, 1]$ is not Lebesgue measurable, $m^*(E) > 0$, the sets $E + q$, $q \in [-1, 1] \cap \mathbb{Q}$, are pairwise disjoint,

$$m^*\left(\bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q)\right) \leq m^*([-1, 2]) = 3 < \infty$$

and

$$\sum_{q \in [-1, 1] \cap \mathbb{Q}} m^*(E + q) = \sum_{q \in [-1, 1] \cap \mathbb{Q}} m^*(E) = \infty.$$

Thus

$$m^*\left(\bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q)\right) \neq \sum_{q \in [-1, 1] \cap \mathbb{Q}} m^*(E + q)$$

and countable additivity on pairwise disjoint sets fails.

Remark 1.80. A Lebesgue nonmeasurable set is not a Borel set, since all Borel sets are Lebesgue measurable.

Remark 1.81. In the n -dimensional case, we may consider an equivalence relation $x \sim y \iff x - y \in \mathbb{Q}^n$ on \mathbb{R}^n (exercise). Here \mathbb{Q}^n denotes the set of points in \mathbb{R}^n with rational coordinates.

Remark 1.82. By a modification of the above proof we see that any set $A \subset \mathbb{R}$ with $m^*(A) > 0$ contains a set B which is not Lebesgue measurable.

Reason. Let $A \subset \mathbb{R}$ be a set with $m^*(A) > 0$. Then there must be at least one interval $[i, i+1]$, $i \in \mathbb{Z}$, such that $m^*(A \cap [i, i+1]) > 0$, otherwise

$$m^*(A) = m^*\left(\bigcup_{i \in \mathbb{Z}} (A \cap [i, i+1])\right) \leq \sum_{i \in \mathbb{Z}} m^*(A \cap [i, i+1]) = 0.$$

By a translation, we may assume that $m^*(A \cap [0, 1]) > 0$ and $A \subset [0, 1]$. By the notation of the proof of the previous theorem,

$$A = \bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (E + q) \cap A.$$

Again, by countable subadditivity, at least one of the sets $(E + q) \cap A$, $q \in [-1, 1]$, has positive Lebesgue outer measure. Set $B = (E + q) \cap A$ with $m^*(B) > 0$.

The same argument as in the proof of the previous theorem shows that B is not Lebesgue measurable. Indeed, assume that B is Lebesgue measurable. Since the translated sets $B + q$, $q \in [-1, 1] \cap \mathbb{Q}$, are disjoint and Lebesgue measurable, by countable additivity and translation invariance,

$$m^*\left(\bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (B + q)\right) = \sum_{q \in [-1, 1] \cap \mathbb{Q}} m^*(B + q) = \sum_{q \in [-1, 1] \cap \mathbb{Q}} m^*(B) = \infty,$$

since $m^*(B) > 0$. On the other hand,

$$m^*\left(\bigcup_{q \in [-1, 1] \cap \mathbb{Q}} (B + q)\right) \leq m^*([-1, 2]) = 3 < \infty.$$

This is a contradiction and thus B cannot be Lebesgue measurable. ■

Remark 1.83. Hausdorff (1914) has shown that for any dimension $n = 1, 2, \dots$, there is no countably additive measure defined on all subsets of \mathbb{R}^n that is invariant under isometries (translations and rotations) and assigns measure one to the unit cube. The Banach-Tarski (1924) paradox shows that the unit ball in \mathbb{R}^n with $n = 3, 4, \dots$ can be cut into a finite number of pairwise disjoint pieces, which can then be reassembled (after translating and rotating each of the pieces) to form two disjoint copies of the original ball (or a ball of any given radius), see [3, Appendix G] and [13]. The pieces used in this decomposition are irregular sets and their construction applies the axiom of choice. In particular, the pieces cannot be Lebesgue measurable, because otherwise additivity on pairwise disjoint sets fails. Banach-Tarski paradox does not hold in \mathbb{R}^2 . This is because \mathbb{R}^2 has less symmetries compared to \mathbb{R}^3 (this has nothing to do with the existence of non-measurable sets).

1.11 The Cantor set

Cantor sets constructed in this section give several examples of unexpected features in analysis. The middle thirds Cantor set is a subset of the interval $C_0 = [0, 1]$. The construction will proceed in steps. At the first step, let $I_{1,1}$ denote the open interval $(\frac{1}{3}, \frac{2}{3})$. Then $I_{1,1}$ is the open middle third of C_0 . At the second step we denote two open intervals $I_{2,1}$ and $I_{2,2}$ each being the open middle third of one of the two intervals comprising $I \setminus I_{1,1}$ and so forth. At the k th step, we obtain 2^{k-1} pairwise disjoint open intervals $I_{k,i}$, $i = 1, \dots, 2^{k-1}$, and denote

$$C_0 = [0, 1], \quad C_k = C_{k-1} \setminus \bigcup_{i=1}^{2^{k-1}} I_{k,i}, \quad k = 1, 2, \dots$$

Note:

$$C_k = \bigcup_{a_1, \dots, a_k \in \{0, 2\}} \left[\sum_{i=1}^k \frac{a_i}{3^i}, \sum_{i=1}^k \frac{a_i}{3^i} + \frac{1}{3^k} \right].$$

Thus C_k consists of 2^k closed intervals of length $\frac{1}{3^k}$. Let us denote these intervals by $J_{k,i}$, $i = 1, 2, \dots, 2^k$.

The (middle thirds) Cantor set is the intersection of all sets C_k , that is,

$$C = \bigcap_{k=0}^{\infty} C_k.$$

Note:

$$C = \left\{ \sum_{i=1}^{\infty} \frac{a_i}{3^i} : a_i \in \{0, 2\}, i = 1, 2, \dots \right\}.$$

Note that C contains more points than the end points $\{\frac{1}{3}, \frac{2}{3}, \frac{1}{9}, \frac{2}{9}, \frac{7}{9}, \frac{8}{9}, \frac{1}{27}, \dots\}$ of the extracted open intervals. For example, $\frac{1}{4} \in C$, but it is not an end point of any of the intervals (exercise).

Since every C_k , $k = 0, 1, 2, \dots$, is closed, the intersection C is closed. Since C is also bounded, it is a compact subset of $[0, 1]$.




Figure 1.18: The Cantor construction.

Claim: C is uncountable.

Reason. For a contradiction, assume that $C = \{x_1, x_2, \dots\}$ is countable. Let J_1 be one of the closed intervals $J_{1,i}$, $i = 1, 2$, in the first step of construction of the Cantor set with $x_1 \notin J_1$. We continue recursively. Let J_2 be one of the closed intervals $J_{2,i}$, $i = 1, 2, 3, 4$, in the second step of construction of the Cantor set with $x_2 \notin J_2$ and $J_2 \subset J_1$. By continuing this way, we obtain a decreasing sequence of closed intervals $J_{k+1} \subset J_k$, $k = 1, 2, \dots$, such that $C \cap \bigcap_{k=1}^{\infty} J_k = \emptyset$. On the other hand, $\bigcap_{k=1}^{\infty} J_k \neq \emptyset$ and thus there exists a point $x \in \bigcap_{k=1}^{\infty} J_k$. By the definition of the Cantor set, we have $x \in C$, which implies $C \cap \bigcap_{k=1}^{\infty} J_k \neq \emptyset$. This is a contradiction. ■

Moreover, C is nowhere dense and perfect (exercise). A set is called nowhere dense if its closure does not have interior points and perfect if it does not have isolated points, that is, every point of the set is a limit point of the set. We show that C is an uncountable set of measure zero.

Claim: $m^*(C) = 0$.

Reason. Since

$$m^*(C_k) = \sum_{i=1}^{2^k} m^*(J_{k,i}) = \sum_{i=1}^{2^k} \left(\frac{1}{3}\right)^k = 2^k \left(\frac{1}{3}\right)^k = \left(\frac{2}{3}\right)^k,$$

by Theorem 1.22 we have

$$m^*(C) = m^*\left(\bigcap_{k=0}^{\infty} C_k\right) = \lim_{k \rightarrow \infty} m^*(C_k) = 0.$$

This can be also seen directly from the definition of the Lebesgue measure, since C_k consists of finitely many intervals whose lengths sum up to $(\frac{2}{3})^k$. This is arbitrarily small by choosing k large enough. ■

Remark 1.84. Every real number can be represented as a decimal expansion. Instead of using base 10, we may take for example 3 as the base. In particular, every $x \in [0, 1]$ can be written as a ternary expansion

$$x = \sum_{i=1}^{\infty} \frac{\alpha_i}{3^i},$$

where $\alpha_i = 0, 1$ or 2 for every $i = 1, 2, \dots$. We denote this as $x = .\alpha_1\alpha_2\dots$. In general, this decomposition is not unique. For example,

$$\frac{1}{3} = \sum_{i=2}^{\infty} \frac{2}{3^i}$$

and $\frac{4}{9} = .11000\dots = .10222\dots$. The reason for this is that

$$\sum_{i=1}^{\infty} \frac{2}{3^i} = 1.$$

The ternary expansion is unique except for a certain type of ambiguity. A number has two different expansions if and only if it has a terminating ternary expansion, that is, only finitely many α_i 's are nonzero. Let us look at the construction of the Cantor set again. At the first stage we remove the middle third $I_{1,1}$. If $\frac{1}{3} < x < \frac{2}{3}$, then $x = .1\alpha_2\alpha_3\dots$. If $x \in [0, 1] \setminus I_{1,1}$, then $x = .0\alpha_2\alpha_3\dots$ or $x = .2\alpha_2\alpha_3\dots$. In either case the value of α_1 determines which of the three subintervals contains x . Repeating this argument show that $x \in [0, 1]$ belongs to the Cantor middle thirds set if and only if it has a ternary expansion consisting only on 0's and 2's.

The construction of a Cantor type set C can be modified so that at the k th, stage of the construction we remove 2^{k-1} centrally situated open intervals each of length l_k , $k = 1, 2, \dots$, with $l_1 + 2l_2 + \dots + 2^{k-1}l_k < 1$. If l_k , $k = 1, 2, \dots$, are chosen small enough, then

$$\sum_{k=1}^{\infty} 2^{k-1}l_k < 1.$$

In this case, we have

$$0 < 1 - \sum_{k=1}^{\infty} 2^{k-1}l_k = m^*(C) < 1.$$

and C is called a fat Cantor set. Note that C is a compact nowhere dense and perfect set of positive Lebesgue measure. Observe that $U = [0, 1] \setminus C$ is an open set with $\partial U = C$ and $m^*(\partial U) = m^*(C) > 0$.

T H E M O R A L : The boundary of an open set may have positive Lebesgue measure.

See [6, p. 83–86] and [16, p. 85–87] for more on the Cantor set.

The class of measurable functions will play a central role in the integration theory. This class is closed under usual operations and limits, but certain unexpected features occur. Measurable functions can be approximated by simple functions. Egoroff's theorem states that pointwise convergence of a sequence of measurable functions is almost uniform and Lusin's theorem states that a measurable function is almost continuous.

2

Measurable functions

2.1 Calculus with infinities

Throughout measure and integration theory we encounter $\pm\infty$. One reason for this is that we want to consider sets of infinite measure as \mathbb{R}^n with respect to the Lebesgue measure. Another reason is that we want to consider functions with singularities as $f : \mathbb{R}^n \rightarrow [0, \infty]$, $f(x) = |x|^{-\alpha}$ with $\alpha > 0$. Here we use the interpretation that $f(0) = \infty$. In addition, even if we only consider real valued functions, the limes superiors of sequences and sums of functions may be infinite at some points.

We consider the set of extended real numbers $[-\infty, \infty] = \mathbb{R} \cup \{-\infty\} \cup \{+\infty\}$. For simplicity, we write ∞ for $+\infty$. We shall use the following conventions for arithmetic operations on $[-\infty, \infty]$. For $a \in \mathbb{R}$, we define

$$a + (\pm\infty) = (\pm)\infty + a = \pm\infty$$

and $(\pm\infty) + (\pm\infty) = \pm\infty$. Subtraction is defined in a similar manner, but $(\pm\infty) + (-\infty)$ and $(\pm\infty) - (\pm\infty)$ are undefined. For multiplication, we define

$$a \cdot (\pm\infty) = (\pm\infty) \cdot a = \begin{cases} \pm\infty, & a > 0, \\ 0, & a = 0, \\ \mp\infty, & a < 0, \end{cases}$$

and $(\pm\infty) \cdot (\pm\infty) = +\infty$ and $(\pm\infty) \cdot (-\infty) = -\infty$. With these definitions the standard commutative, associative and distributive rules hold in $[-\infty, \infty]$ in the usual manner.

Cancellation properties have to be considered with some care. For example, $a + b = a + c$ implies $b = c$ only when $|a| < \infty$ and $ab = ac$ implies $b = c$ only when $0 < |a| < \infty$. A general fact is that the cancellation is safe if all terms are finite

and nonzero in the case of division. Finally we note that with this interpretation, for example, all sums of nonnegative terms $x_i \in [0, \infty]$, $i = 1, 2, \dots$, are convergent with

$$\sum_{i=1}^{\infty} x_i = \lim_{k \rightarrow \infty} \sum_{i=1}^k x_i \in [0, \infty].$$

We shall use this interpretation without further notice.

2.2 Measurable functions

A G R E E M E N T : From now on, we shall not distinguish outer measures from measures with the interpretation that an outer measure restricted to measurable sets is a measure.

Consider a function $f : X \rightarrow [-\infty, \infty]$. Recall that the preimage of a set $A \subset [-\infty, \infty]$ is

$$f^{-1}(A) = \{x \in X : f(x) \in A\}.$$

The preimage has the properties

$$\begin{aligned} f^{-1}\left(\bigcap_{i=1}^{\infty} A_i\right) &= \bigcap_{i=1}^{\infty} f^{-1}(A_i), \\ f^{-1}\left(\bigcup_{i=1}^{\infty} A_i\right) &= \bigcup_{i=1}^{\infty} f^{-1}(A_i) \end{aligned}$$

and

$$f^{-1}(A^c) = X \setminus f^{-1}(A)$$

for $A, A_i \subset [-\infty, \infty]$, $i = 1, 2, \dots$. Here $A^c = [-\infty, \infty] \setminus A$.

We begin with a definition of measurable function.

Definition 2.1. Let μ be a measure on X . The function $f : X \rightarrow [-\infty, \infty]$ is μ -measurable, if the set

$$f^{-1}((a, \infty]) = \{x \in X : f(x) > a\}$$

is μ -measurable for every $a \in \mathbb{R}$.

T H E M O R A L : As we shall see, it is important that all distribution sets are measurable in the definition of integral.

Remarks 2.2:

- (1) Every continuous function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is Lebesgue measurable.

Reason. Since f is continuous, the set $\{x \in \mathbb{R}^n : f(x) > a\}$ is open for every $a \in \mathbb{R}$. The Lebesgue measure is a Borel measure, see Lemma 1.55, and thus the set $\{x \in \mathbb{R}^n : f(x) > a\}$ is Lebesgue measurable for every $a \in \mathbb{R}$. ■

(2) The set $A \subset \mathbb{R}^n$ is Lebesgue measurable set if and only if the characteristic function

$$f : \mathbb{R}^n \rightarrow \mathbb{R}, f(x) = \chi_A(x) = \begin{cases} 1, & x \in A, \\ 0, & x \in \mathbb{R}^n \setminus A, \end{cases}$$

is a Lebesgue measurable function.

Reason.

$$\{x \in \mathbb{R}^n : f(x) > a\} = \begin{cases} \mathbb{R}^n, & a < 0, \\ A, & 0 \leq a < 1, \\ \emptyset, & a \geq 1. \end{cases}$$

■

By considering a set A which is not Lebesgue measurable, see Section 1.10, we conclude that χ_A a nonmeasurable function with respect to the Lebesgue measure.

(3) The previous remark holds for all outer measures. Moreover, a linear combination of finitely many characteristic functions of measurable sets is a measurable function. Such a function is called a simple function, see Definition 2.32.

(4) If μ is an outer measure for which all sets are μ -measurable, then all functions are μ -measurable. See Remark 1.6 (7).

(5) If the only measurable sets are \emptyset ja X , then only constant functions are measurable. See Remark 1.6 (6).

Remark 2.3. For μ -measurable subset $A \subset X$ and a function $f : A \rightarrow [-\infty, \infty]$, we consider the zero extension $\tilde{f} : X \rightarrow [-\infty, \infty]$,

$$\tilde{f}(x) = \begin{cases} f(x), & x \in A, \\ 0, & x \in X \setminus A. \end{cases}$$

Then f is μ -measurable on A if and only if \tilde{f} is μ -measurable on X .

T H E M O R A L : A function defined on a subset is measurable if and only if its zero extension to the entire space is measurable. This allows us to consider functions defined on measurable subsets.

Lemma 2.4. Let μ be a measure on X and $f : X \rightarrow [-\infty, \infty]$. Then the following claims are equivalent:

- (1) f is μ -measurable,
- (2) $\{x \in X : f(x) \geq a\}$ is μ -measurable for every $a \in \mathbb{R}$,
- (3) $\{x \in X : f(x) < a\}$ is μ -measurable for every $a \in \mathbb{R}$,
- (4) $\{x \in X : f(x) \leq a\}$ is μ -measurable for every $a \in \mathbb{R}$.

Proof. The equivalence follows from the fact that the collection of μ -measurable sets is a σ -algebra, see Lemma 1.11.

(1) \Rightarrow (2) We have

$$\{x \in X : f(x) \geq a\} = \bigcap_{i=1}^{\infty} \{x \in X : f(x) > a - \frac{1}{i}\}.$$

The sets $\{x \in X : f(x) > a - \frac{1}{i}\}$, $i = 1, 2, \dots$, are μ -measurable by the assumption and thus $\{x \in X : f(x) \geq a\}$ is μ -measurable as a countable intersection of μ -measurable sets.

(2) \Rightarrow (3) We have

$$\{x \in X : f(x) < a\} = X \setminus \{x \in X : f(x) \geq a\}.$$

The set $\{x \in X : f(x) \geq a\}$ is μ -measurable by the assumption and thus $f^{-1}(\{x \in X : f(x) < a\})$ is μ -measurable as a complement of a μ -measurable set.

(3) \Rightarrow (4) We have

$$\{x \in X : f(x) \leq a\} = \bigcap_{i=1}^{\infty} \{x \in X : f(x) < a + \frac{1}{i}\}$$

The sets $\{x \in X : f(x) < a + \frac{1}{i}\}$, $i = 1, 2, \dots$, are μ -measurable by the assumption and thus $\{x \in X : f(x) \leq a\}$ is μ -measurable as a countable intersection of μ -measurable sets.

(4) \Rightarrow (1) We have

$$\{x \in X : f(x) > a\} = X \setminus \{x \in X : f(x) \leq a\}.$$

The set $\{x \in X : f(x) \leq a\}$ is μ -measurable by the assumption and thus $f^{-1}(\{x \in X : f(x) > a\})$ is μ -measurable as a complement of a μ -measurable set. \square

As we shall see in Section 2.3, the preimage of a measurable set is not measurable, in general. However, the following characterization holds for a measurable function.

Lemma 2.5. A function $f : X \rightarrow [-\infty, \infty]$ is μ -measurable if and only if $f^{-1}(\{-\infty\})$ and $f^{-1}(\{\infty\})$ are μ -measurable and $f^{-1}(B)$ is μ -measurable for every Borel set $B \subset \mathbb{R}$.

Remark 2.6. The proof will show that we could require that $f^{-1}(B)$ is μ -measurable for every open set B . This is analogous to the fact that a function is continuous if and only if $f^{-1}(B)$ is open for every open set B .

Proof. \Rightarrow Note that

$$f^{-1}(\{-\infty\}) = \{x \in X : f(x) = -\infty\} = \bigcap_{i=1}^{\infty} \{x \in X : f(x) < -i\}$$

and

$$f^{-1}(\{\infty\}) = \{x \in X : f(x) = \infty\} = \bigcap_{i=1}^{\infty} \{x \in X : f(x) > i\}$$

are μ -measurable sets.

Let

$$\mathcal{F} = \{B \subset \mathbb{R} : B \text{ is a Borel set and } f^{-1}(B) \text{ is } \mu\text{-measurable}\}$$

Claim: \mathcal{F} is a σ -algebra.

Reason. Clearly $\emptyset \in \mathcal{F}$. If $B \in \mathcal{F}$, then $f^{-1}(\mathbb{R} \setminus B) = X \setminus f^{-1}(B)$ is μ -measurable and thus $\mathbb{R} \setminus B \in \mathcal{F}$. If $B_i \in \mathcal{F}$, $i = 1, 2, \dots$, then

$$f^{-1}\left(\bigcup_{i=1}^{\infty} B_i\right) = \bigcup_{i=1}^{\infty} f^{-1}(B_i)$$

is μ -measurable and thus $\bigcup_{i=1}^{\infty} B_i \in \mathcal{F}$. ■

Then we show that \mathcal{F} contains all open subsets of \mathbb{R} . Since every open set in \mathbb{R} is a countable union of pairwise disjoint open intervals and \mathcal{F} is a σ -algebra, it is enough to show that every open interval $(a, b) \in \mathcal{F}$. We note that

$$f^{-1}((a, b)) = f^{-1}([-\infty, b)) \cap f^{-1}((a, \infty])$$

where $f^{-1}([-\infty, b)) = \{x \in X : f(x) < b\}$ and $f^{-1}((a, \infty]) = \{x \in X : f(x) > a\}$ are μ -measurable. This implies that $f^{-1}((a, b))$ is μ -measurable. Since \mathcal{F} is a σ -algebra that contains open sets, it also contains Borel sets.

Let $B = (a, \infty)$ with $a \in \mathbb{R}$. Then

$$\{x \in X : f(x) > a\} = f^{-1}(B \cup \{\infty\}) = f^{-1}(B) \cup f^{-1}(\{\infty\})$$

is μ -measurable. □

As we shall see in Section 2.3, a composed function of two measurable functions is not measurable, in general. However, the following result holds for a composed function.

Lemma 2.7. Let μ be a measure on X . If $f : X \rightarrow \mathbb{R}$ is μ -measurable and $g : \mathbb{R} \rightarrow \mathbb{R}$ is continuous, then the composed function $g \circ f$ is measurable.

Proof. By Lemma 2.5 and Remark 2.6, it is enough to show that the preimage $(g \circ f)^{-1}(B)$ of every open set $B \subset \mathbb{R}$ is μ -measurable. Note that $(g \circ f)^{-1}(B) = f^{-1}(g^{-1}(B))$, since

$$\begin{aligned} x \in (g \circ f)^{-1}(B) &\iff (g \circ f)(x) \in B \iff g(f(x)) \in B \\ &\iff f(x) \in g^{-1}(B) \iff x \in f^{-1}(g^{-1}(B)). \end{aligned}$$

Since g is continuous, the preimage $g^{-1}(B)$ of an open set B is open. Since f is μ -measurable, the preimage $f^{-1}(g^{-1}(B))$ of an open set $g^{-1}(B)$ is μ -measurable. Thus $(g \circ f)^{-1}(B)$ is a μ -measurable set and $g \circ f$ is a μ -measurable function. □

Remark 2.8. In fact, it is enough to assume in Lemma 2.7 that g is a Borel function, that is, the preimage of every Borel set is a Borel set.

Reason. Let $A \subset \mathbb{R}$ be a Borel set and $g : X \rightarrow \mathbb{R}$ be a Borel function. Then $g^{-1}(A)$ is a Borel set. Since f is μ -measurable, Lemma 2.5 implies that the preimage $f^{-1}(g^{-1}(B))$ is μ -measurable. ■

Remark 2.9. We briefly discuss an abstract version of a definition of a measurable function.

- (1) Assume that (X, \mathcal{M}, μ) and (Y, \mathcal{N}, ν) are measure spaces and $f : X \rightarrow Y$ is a function. Then f is said to be measurable with respect to σ -algebras \mathcal{M} and \mathcal{N} if $f^{-1}(A) \in \mathcal{M}$ whenever $A \in \mathcal{N}$. In our approach we consider $Y = [-\infty, \infty]$ and \mathcal{N} equals the Borel sets in $[-\infty, \infty]$ (endowed with the order topology, see [15]).
- (2) Let (X, \mathcal{M}, μ) , (Y, \mathcal{N}, ν) and (Z, \mathcal{P}, γ) be abstract measure spaces. If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are measurable functions in the sense of (1), then the composed function $g \circ f$ is measurable. This follows directly from the abstract definition of measurability. We might be tempted to conclude that the composed function of Lebesgue measurable functions is Lebesgue measurable. This is not always the case, since the preimage of a Lebesgue measurable set is not necessarily Lebesgue measurable, see Section 2.3. This means that we cannot replace Borel sets by measurable sets in Lemma 2.5.

Lemma 2.10. If $f, g : X \rightarrow [-\infty, \infty]$ are μ -measurable functions, then

$$\{x \in X : f(x) > g(x)\}$$

is a μ -measurable set.

Proof. Let $\mathbb{Q} = \bigcup_{i=1}^{\infty} \{q_i\}$ be an enumeration of the rational numbers. If $f(x) > g(x)$, there exists $q_i \in \mathbb{Q}$ such that $f(x) > q_i > g(x)$. This implies that

$$\{x \in X : f(x) > g(x)\} = \bigcup_{i=1}^{\infty} (\{x \in X : f(x) > q_i\} \cap \{x \in X : g(x) < q_i\})$$

is a μ -measurable set. □

Remark 2.11. The sets

$$\{x \in X : f(x) \leq g(x)\} = X \setminus \{x \in X : f(x) > g(x)\}$$

and

$$\{x \in X : f(x) = g(x)\} = \{x \in X : f(x) \leq g(x)\} \cap \{x \in X : f(x) \geq g(x)\}$$

are μ -measurable as well.

Let $f : X \rightarrow [-\infty, \infty]$. The positive part of f is

$$f^+(x) = \max\{f(x), 0\} = f(x)\chi_{\{x \in X : f(x) \geq 0\}} = \begin{cases} f(x), & f(x) \geq 0, \\ 0, & f(x) < 0, \end{cases}$$

and the negative part is

$$f^-(x) = -\min\{f(x), 0\} = -f(x)\chi_{\{x \in X : f(x) \leq 0\}} = \begin{cases} -f(x), & f(x) \leq 0, \\ 0, & f(x) > 0. \end{cases}$$

Observe that $f^+, f^- \geq 0$, $f = f^+ - f^-$ and $|f| = f^+ + f^-$. Splitting a function into positive and negative parts will be a useful tool in measure theory.

Lemma 2.12. A function $f : X \rightarrow [-\infty, \infty]$ is μ -measurable if and only if f^+ and f^- are μ -measurable.

Proof. \Rightarrow Assume that f is a μ -measurable function. Then

$$\{x \in X : f^+(x) > a\} = \begin{cases} \{x \in X : f(x) > a\}, & a \geq 0, \\ X, & a < 0, \end{cases}$$

is a μ -measurable set. This implies that f^+ is a μ -measurable function. Moreover, $f^- = (-f)^+$.

\Leftarrow Assume that f^+ and f^- are μ -measurable functions. Then

$$\{x \in X : f(x) > a\} = \begin{cases} \{x \in X : f^+(x) > a\}, & a \geq 0, \\ \{x \in X : f^-(x) < -a\}, & a < 0, \end{cases}$$

is a μ -measurable set. This implies that f is a μ -measurable function. \square

Theorem 2.13. Assume that $f, g : X \rightarrow [-\infty, \infty]$ are μ -measurable functions and $a \in \mathbb{R}$. Then af , $f + g$, $\max\{f, g\}$, $\min\{f, g\}$, fg , $\frac{f}{g}$ ($g \neq 0$), are μ -measurable functions.

W A R N I N G : Since functions are extended real valued, we need to take care about the definitions of $f + g$ and fg . The sum is defined outside the bad set

$$B = \{x \in X : f(x) = \infty \text{ and } g(x) = -\infty\} \cup \{x \in X : f(x) = -\infty \text{ and } g(x) = \infty\},$$

since in B we have $\infty - \infty$ situation. We define

$$(f + g)(x) = \begin{cases} f(x) + g(x), & x \in X \setminus B, \\ a, & x \in B, \end{cases}$$

where $a \in [-\infty, \infty]$ is arbitrary.

Proof. Note that

$$(f+g)^{-1}(\{-\infty\}) = f^{-1}(\{-\infty\}) \cup g^{-1}(\{-\infty\})$$

and

$$(f+g)^{-1}(\{\infty\}) = f^{-1}(\{\infty\}) \cup g^{-1}(\{\infty\})$$

are μ -measurable. Let $a \in \mathbb{R}$. Since

$$\{x \in X : a - f(x) > \lambda\} = \{x \in X : f(x) < a - \lambda\}$$

for every $\lambda \in \mathbb{R}$, the function $a - f$ is μ -measurable. By Lemma 2.10,

$$\{x \in X : f(x) + g(x) > a\} = \{x \in X : g(x) > a - f(x)\}$$

is μ -measurable for every $a \in \mathbb{R}$. This can be also seen directly from

$$(f+g)^{-1}((-\infty, a)) = \bigcup_{r,s \in \mathbb{Q}, r+s < a} (f^{-1}((-\infty, r)) \cap g^{-1}((-\infty, s))).$$

The functions $|f|$ and g^2 are μ -measurable, see the remark below. Then we may use the formulas

$$\max\{f, g\} = \frac{1}{2}(f + g + |f - g|), \min\{f, g\} = \frac{1}{2}(f + g - |f - g|)$$

and

$$fg = \frac{1}{2}((f+g)^2 - f^2 - g^2).$$

□

W A R N I N G : f^2 measurable does not imply that f measurable.

Reason. Let $A \subset X$ be a nonmeasurable set and

$$f : X \rightarrow \mathbb{R}, f(x) = \begin{cases} 1, & x \in A, \\ -1, & x \in X \setminus A. \end{cases}$$

Then $f^2 = 1$ is measurable, but $\{x \in X : f(x) > 0\} = A$ is not a measurable set. ■

2.3 Cantor-Lebesgue function

Recall the construction of the Cantor set from Section 1.11. At the k th step, we have 2^{k-1} open pairwise disjoint open intervals $I_{k,i}$, $i = 1, \dots, 2^{k-1}$. Let $m = 1, 2, \dots$. Consider all open intervals $I_{k,i}$, with $k = 1, \dots, m$, $i = 1, \dots, 2^{k-1}$, used in the construction of the Cantor set at the steps $1, \dots, m$. Note that there are altogether $2^0 + 2^1 + 2^2 + \dots + 2^{m-1} = 2^m - 1$ intervals. Denote these intervals by $\tilde{I}_{m,i}$, $i = 1, \dots, 2^m - 1$, organized from left to right.

As in Section 1.11 we have

$$C_0 = [0, 1], \quad C_k = [0, 1] \setminus \bigcup_{i=1}^{2^k-1} \tilde{I}_{k,i}, \quad k = 1, 2, \dots,$$

and the middle thirds Cantor set is $C = \bigcap_{k=0}^{\infty} C_k$. We have seen that C is an uncountable set of Lebesgue measure zero. Define a continuous function $f_k : [0, 1] \rightarrow [0, 1]$ by $f_k(0) = 0$, $f_k(1) = 1$, $f_k(x) = \frac{i}{2^k}$, whenever $x \in \tilde{I}_{k,i}$, $i = 1, 2, \dots, 2^k - 1$ and f_k is linear on C_k , $k = 1, 2, \dots$




Figure 2.1: The construction of the Cantor-Lebesgue function.

Then $f_k \in C([0, 1])$, f is increasing and

$$|f_k(x) - f_{k+1}(x)| < \frac{1}{2^k}$$

for every $x \in [0, 1]$. Since

$$|f_k(x) - f_{k+m}(x)| \leq \sum_{j=k}^{k+m-1} \frac{1}{2^j} < \frac{1}{2^{k-1}}$$

for every $x \in [0, 1]$, (f_k) is a Cauchy sequence in the space $(C([0, 1]), \|\cdot\|_{\infty})$, where

$$\|f\|_{\infty} = \sup_{x \in [0, 1]} |f(x)|.$$

This is a complete space and thus there exists $f \in C([0, 1])$ such that $\|f_k - f\|_{\infty} \rightarrow 0$ as $k \rightarrow \infty$. In other words, $f_k \rightarrow f$ uniformly in $[0, 1]$ as $k \rightarrow \infty$. The function f is called the Cantor-Lebesgue function. We collect properties of the Cantor-Lebesgue function below.

- (1) $f : [0, 1] \rightarrow [0, 1]$ is continuous, $f(0) = 0$ and $f(1) = 1$.
- (2) f is nondecreasing and is constant on each interval in the complement of the Cantor set.

(3) $f : [0, 1] \rightarrow [0, 1]$ is onto, that is, $f([0, 1]) = [0, 1]$. In fact $f(C) = [0, 1]$, that is, for every $y \in [0, 1]$ there exists $x \in C$ with $f(x) = y$.

(4) f maps the complement of the Cantor set to a countable set. Thus f maps the the Cantor set C with $m^*(C) = 0$ to a set $f(C)$ with $m^*(f(C)) = 1$.

(5) f is locally constant and thus differentiable in the complement of the Cantor set. Thus f is differentiable almost everywhere and its derivative is zero outside the Cantor set. However,

$$\int_{[0,1]} f'(x) dx = 0 \neq 1 = f(1) - f(0)$$

and therefore the fundamental theorem of calculus does not hold.

(6) f is not differentiable at any point in the Cantor set.

See [2, p. 67], [6, p. 86–101], [10, p. 38], [11, p. 140–141], [16, p. 87–90] and [15] for more on the Cantor-Lebesgue function.

Remark 2.14. Recall from Section 1.11 that

$$C = \left\{ \sum_{i=1}^{\infty} \frac{a_i}{3^i} : a_i \in \{0, 2\}, i = 1, 2, \dots \right\}.$$

It can be shown that if

$$x = \sum_{i=1}^{\infty} \frac{a_i}{3^i} \quad \text{with} \quad a_i \in \{0, 2\}, i = 1, 2,$$

then

$$f(x) = \sum_{i=1}^{\infty} \frac{b_i}{2^i} \quad \text{with} \quad b_i = \frac{a_i}{2}.$$

In this sense, the Cantor-Lebesgue function converts the base three expansions to base two expansions.

Let $g : [0, 1] \rightarrow [0, 2]$, $g(x) = x + f(x)$. Then $g(0) = 0$, $g(1) = 2$, $g \in C([0, 1])$ is strictly increasing and $g([0, 1]) = [0, 2]$. This implies that g is a homeomorphism, that is, g is a continuous function from $[0, 1]$ onto $[0, 2]$ with a continuous inverse function. Since

$$C = \bigcap_{k=0}^{\infty} C_k = \bigcap_{k=1}^{\infty} \left([0, 1] \setminus \bigcup_{i=1}^{2^k-1} \tilde{I}_{k,i} \right) = [0, 1] \setminus \bigcup_{k=1}^{\infty} \bigcup_{i=1}^{2^k-1} \tilde{I}_{k,i},$$

where $\tilde{I}_{k,i}$ are pairwise disjoint open intervals,

$$\begin{aligned}
 m^*(g(C)) &= m^*\left(g([0,1]) \setminus \bigcup_{k=1}^{\infty} \bigcup_{i=1}^{2^k-1} g(\tilde{I}_{k,i})\right) \\
 &= m^*([0,2]) - m^*\left(\bigcup_{k=1}^{\infty} \bigcup_{i=1}^{2^k-1} g(\tilde{I}_{k,i})\right) \quad (g(\tilde{I}_{k,i}) \text{ is an interval}) \\
 &= 2 - \lim_{k \rightarrow \infty} \sum_{i=1}^{2^k-1} m^*(g(\tilde{I}_{k,i})) \quad (g(\tilde{I}_{k,i}) \text{ are pairwise disjoint}) \\
 &= 2 - \lim_{k \rightarrow \infty} \sum_{i=1}^{2^k-1} m^*(\tilde{I}_{k,i}) \quad (g(x) = x + a_{k,i} \quad \forall x \in \tilde{I}_{k,i}) \\
 &= 2 - m^*([0,1] \setminus C) = 2 - 1 = 1. \quad (m^*(C) = 0)
 \end{aligned}$$

Thus g maps the zero measure Cantor set C to $g(C)$ set of measure one. Since $m(g(C)) > 0$, by Remark 1.82, there exists $B \subset g(C)$, which is nonmeasurable with respect to the one-dimensional Lebesgue measure. Let $A = g^{-1}(B)$. Then $A \subset C$ and $m(A) = 0$. This implies that A is Lebesgue measurable. We collect a few observations related to the Cantor-Lebesgue function below.

(1) The homeomorphism g maps the Cantor set C with $m^*(C) = 0$ to a set $g(C)$ with $m^*(g(C)) > 0$.

T H E M O R A L : Sets of Lebesgue measure zero are not necessarily mapped to sets of Lebesgue measure zero in continuous mappings.

(2) The homeomorphism g maps a measurable set A to a nonmeasurable set B .

T H E M O R A L : Lebesgue measurable sets are not necessarily preserved in continuous mappings.

(3) A is a Lebesgue measurable set that is not a Borel set. Assume for the contradiction, that A is a Borel set. Then $B = g(A)$ is a Borel set, since a homeomorphism maps Borel sets to Borel sets (exercise). However, B is not a Lebesgue measurable set, which implies that it is not a Borel set.

T H E M O R A L : There exists a Lebesgue measurable set that is not a Borel set, compare with Corollary 1.74.

(4) Since $A \subset C$, we conclude that the Cantor set has a subset that is not a Borel set. The set A is Lebesgue measurable subset of the Borel set C with $m^*(C) = 0$, but A is not a Borel set.

T H E M O R A L : The restriction of the Lebesgue measure to Borel sets is not a complete measure, see Definition 1.15.

(5) $\chi_A \circ g^{-1} = \chi_B$, where the function χ_B is nonmeasurable, but the functions χ_A and g^{-1} are measurable functions, since A is a measurable set and g^{-1} is continuous.

T H E M O R A L : A composed function of two Lebesgue measurable functions is not necessarily Lebesgue measurable, compare with Lemma 2.7.

(6) g^{-1} is a Lebesgue measurable function for which $(g^{-1})^{-1}(A) = B$.

T H E M O R A L : The preimage of a measurable set under a measurable function is not necessarily a measurable set, compare with Lemma 2.5.

2.4 Lipschitz mappings on \mathbb{R}^n

The Cantor-Lebesgue function showed that Lebesgue measurability of a set is not necessarily preserved in continuous mappings. In this section we study certain conditions for a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, which guarantee that f maps Lebesgue measurable sets to Lebesgue measurable sets.

Definition 2.15. A mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is said to be Lipschitz continuous, if there exists a constant L such that

$$|f(x) - f(y)| \leq L|x - y|$$

for every $x, y \in \mathbb{R}^n$.

Remark 2.16. A mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is of the form $f(x) = (f_1(x), \dots, f_n(x))$, where $x = (x_1, \dots, x_n)$ and the coordinate functions $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for $i = 1, \dots, n$. Such a mapping f is Lipschitz continuous if and only if all coordinate functions f_i , $i = 1, \dots, n$, satisfy a Lipschitz condition

$$|f_i(x) - f_i(y)| \leq L_i|x - y|$$

for every $x, y \in \mathbb{R}^n$ with some constant L_i .

Examples 2.17:

(1) Every linear mapping $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is Lipschitz continuous.

Reason. Let A be the $n \times n$ -matrix representing L . Then

$$|L(x) - L(y)| = |Ax - Ay| = |A(x - y)| \leq \|A\||x - y|$$

for every $x, y \in \mathbb{R}^n$, where $\|A\| = \max\{|a_{ij}| : i, j = 1, \dots, n\}$. ■

(2) Every mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $f = (f_1, \dots, f_n)$, whose coordinate functions f_i , $i = 1, \dots, n$, have bounded first partial derivatives in \mathbb{R}^n , is Lipschitz continuous.

Reason. By the fundamental theorem of calculus,

$$f_i(x) - f_i(y) = \int_0^1 \frac{\partial}{\partial t} (f_i((1-t)x + ty)) dt = \int_0^1 \nabla f_i((1-t)x + ty) \cdot (y - x) dt.$$

This implies

$$|f_i(x) - f_i(y)| \leq \int_0^1 |\nabla f_i((1-t)x + ty)| |x - y| dt \leq \sup_{z \in \mathbb{R}^n} |\nabla f_i(z)| |x - y|$$

for every $x, y \in \mathbb{R}^n$. ■

Lemma 2.18. Assume that $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a Lipschitz continuous mapping. Then $m^*(f(A)) = 0$ whenever $m^*(A) = 0$.

T H E M O R A L : A Lipschitz mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ maps sets of Lebesgue measure zero to sets of Lebesgue measure zero.

Proof. Assume that $m^*(A) = 0$ and let $\varepsilon > 0$. Then (exercise) there are balls $\{B(x_i, r_i)\}_{i=1}^\infty$ such that

$$A \subset \bigcup_{i=1}^\infty B(x_i, r_i) \quad \text{and} \quad \sum_{i=1}^\infty m^*(B(x_i, r_i)) < \varepsilon.$$

By the Lipschitz condition,

$$|f(x_i) - f(y)| \leq L|x_i - y| < Lr_i,$$

for every $y \in B(x_i, r_i)$ and thus $f(y) \in B(f(x_i), Lr_i)$. This implies

$$f(A) \subset f\left(\bigcup_{i=1}^\infty B(x_i, r_i)\right) = \bigcup_{i=1}^\infty f(B(x_i, r_i)) \subset \bigcup_{i=1}^\infty B(f(x_i), Lr_i).$$

By monotonicity, countable subadditivity translation invariance and the scaling property of the Lebesgue measure we have

$$\begin{aligned} m^*(f(A)) &\leq m^*\left(\bigcup_{i=1}^\infty B(f(x_i), Lr_i)\right) \leq \sum_{i=1}^\infty m^*(B(f(x_i), Lr_i)) \\ &= L^n \sum_{i=1}^\infty m^*(B(x_i, r_i)) < L^n \varepsilon. \end{aligned}$$

This implies that $m^*(f(A)) = 0$. □

Theorem 2.19. Assume that $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a continuous function which maps sets of Lebesgue measure zero to sets of Lebesgue measure zero. Then f maps Lebesgue measurable sets to Lebesgue measurable sets.

T H E M O R A L : In particular, a Lipschitz mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ maps Lebesgue measurable sets to Lebesgue measurable sets.

Proof. Let $A \subset \mathbb{R}^n$ be a Lebesgue measurable set. Consider $A_k = A \cap B(0, k)$ with $m^*(A_k) < \infty$ for every $k = 1, 2, \dots$. By Theorem 1.73 there exist compact sets $K_i \subset A_k$, $i = 1, 2, \dots$, such that

$$m^*(A_k \setminus \bigcup_{i=1}^\infty K_i) = 0.$$

Since the set A_k can be written as

$$A_k = \left(\bigcup_{i=1}^\infty K_i\right) \cup \left(A_k \setminus \bigcup_{i=1}^\infty K_i\right),$$




Figure 2.2: The image of a set of measure zero.

we have

$$f(A_k) = \bigcup_{i=1}^{\infty} f(K_i) \cup f(A_k \setminus \bigcup_{i=1}^{\infty} K_i).$$

Since a continuous function maps compact sets to compact sets and compact sets are Lebesgue measurable, the countable union $\bigcup_{i=1}^{\infty} f(K_i)$ is a Lebesgue measurable set. On the other hand, by Lemma 2.18 the function f maps sets of measure zero to sets of measure zero. This implies that $f(A_k \setminus \bigcup_{i=1}^{\infty} K_i)$ is of measure zero and thus Lebesgue measurable. The set $f(A_k)$ is measurable as a union of two measurable sets. Finally

$$f(A) = f\left(\bigcup_{k=1}^{\infty} A_k\right) = \bigcup_{k=1}^{\infty} f(A_k)$$

is Lebesgue measurable as a countable union of Lebesgue measurable sets. \square

Remark 2.20. If $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a Lipschitz mapping with constant L , then there exists a constant c , depending only on L and n , such that

$$m^*(f(A)) \leq c m^*(A)$$

for every set $A \subset \mathbb{R}^n$ (exercise).

W A R N I N G : It is important that the source and the target dimensions are same in the results above. There is a measurable subset A of \mathbb{R}^n with respect to the n -dimensional Lebesgue measure such that the projection to the first coordinate

axis is not Lebesgue measurable with respect to the one-dimensional Lebesgue measure. Observe that the projection is a Lipschitz continuous mapping from \mathbb{R}^n to \mathbb{R} with the Lipschitz constant one.

Reason. Let B be a nonmeasurable subset of \mathbb{R} . Then $B \times \{0\}$ is a Lebesgue measurable set in \mathbb{R}^2 , since the two-dimensional Lebesgue measure of $B \times \{0\}$ is zero, but the projection B is not Lebesgue measurable with respect to the one-dimensional Lebesgue measure. ■

2.5 Limits of measurable functions

Next we show that measurability is preserved under limit operations on a sequence (f_i) of μ -measurable functions $f_i : X \rightarrow [-\infty, \infty]$, $i = 1, 2, \dots$. We begin with a useful remark related to preimages in the definition of measurable function.

Remarks 2.21:

(1) Note that

$$\begin{aligned} \sup_i f_i(x) > a &\iff \exists i \text{ such that } f_i(x) > a, \\ \sup_i f_i(x) < a &\iff \exists k \text{ such that } f_i(x) < a - \frac{1}{k} \text{ for } \forall i, \\ \sup_i f_i(x) = \infty &\iff \forall k \exists i \text{ such that } f_i(x) > k. \end{aligned}$$

Thus

$$\begin{aligned} \{x \in X : \sup_i f_i(x) > a\} &= \bigcup_{i=1}^{\infty} \{x \in X : f_i(x) > a\}, \\ \{x \in X : \sup_i f_i(x) < a\} &= \bigcup_{k=1}^{\infty} \bigcap_{i=1}^{\infty} \{x \in X : f_i(x) < a - \frac{1}{k}\}, \\ \{x \in X : \sup_i f_i(x) = \infty\} &= \bigcap_{k=1}^{\infty} \bigcup_{i=1}^{\infty} \{x \in X : f_i(x) > k\}. \end{aligned}$$

Similarly

$$\begin{aligned} \inf_i f_i(x) < a &\iff \exists i \text{ such that } f_i(x) < a, \\ \inf_i f_i(x) > a &\iff \exists k \text{ such that } f_i(x) > a + \frac{1}{k} \text{ for } \forall i, \\ \inf_i f_i(x) = -\infty &\iff \forall k \exists i \text{ such that } f_i(x) < -k. \end{aligned}$$

Thus

$$\begin{aligned} \{x \in X : \inf_i f_i(x) < a\} &= \bigcup_{i=1}^{\infty} \{x \in X : f_i(x) < a\}, \\ \{x \in X : \sup_i f_i(x) < a\} &= \bigcup_{k=1}^{\infty} \bigcap_{i=1}^{\infty} \{x \in X : f_i(x) > a + \frac{1}{k}\}, \\ \{x \in X : \inf_i f_i(x) = -\infty\} &= \bigcap_{k=1}^{\infty} \bigcup_{i=1}^{\infty} \{x \in X : f_i(x) < -k\}. \end{aligned}$$

All sets above are measurable if $f_i : X \rightarrow [-\infty, \infty]$, $i = 1, 2, \dots$, are μ -measurable functions.

(2) There is an advantage in considering strict inequalities above. For example, $\sup_i f_i(x) > a$ if and only if $f_i(x) > a$ for some i . On the other hand, it is not true that $\sup_i f_i(x) \geq a$ if and only if $f_i(x) \geq a$ for some i . For example, consider a strictly increasing sequence (f_i) such that $f_i(x) < a$ for every $i = 1, 2, \dots$ and $\lim_{i \rightarrow \infty} f_i(x) = a$. Then $\sup_i f_i(x) \geq a$, but $f_i(x) < a$ for every $i = 1, 2, \dots$

(3) Recall that

$$\limsup_{i \rightarrow \infty} f_i(x) = \inf_{j \geq 1} (\sup_{i \geq j} f_i(x))$$

and

$$\liminf_{i \rightarrow \infty} f_i(x) = \sup_{j \geq 1} (\inf_{i \geq j} f_i(x)).$$

Theorem 2.22. Assume that $f_i : X \rightarrow [-\infty, \infty]$, $i = 1, 2, \dots$, are μ -measurable functions. Then

$$\sup_i f_i, \quad \inf_i f_i, \quad \limsup_{i \rightarrow \infty} f_i \quad \text{and} \quad \liminf_{i \rightarrow \infty} f_i$$

are μ -measurable functions.

Proof. Since

$$\{x \in X : \sup_i f_i(x) > a\} = \bigcup_{i=1}^{\infty} \{x \in X : f_i(x) > a\}$$

for every $a \in \mathbb{R}$, the function $\sup_i f_i$ is μ -measurable. The measurability of $\inf_i f_i$ follows from

$$\inf_i f_i(x) = -\sup_i (-f_i(x))$$

or from

$$\{x \in X : \inf_i f_i(x) < a\} = \bigcup_{i=1}^{\infty} \{x \in X : f_i(x) < a\}$$

for every $a \in \mathbb{R}$. The claims that $\limsup_{i \rightarrow \infty} f_i$ and $\liminf_{i \rightarrow \infty} f_i$ are μ -measurable functions follow immediately. \square

Theorem 2.23. Assume that $f_i : X \rightarrow [-\infty, \infty]$, $i = 1, 2, \dots$, are μ -measurable functions such that the sequence $(f_i(x))$ converges for every $x \in X$ as $i \rightarrow \infty$. Then

$$f = \lim_{i \rightarrow \infty} f_i$$

is a μ -measurable function.

T H E M O R A L : Measurability is preserved in taking limits. This is a very important property of a measurable function.

Proof. This follows from the previous theorem, since

$$f = \limsup_{i \rightarrow \infty} f_i = \liminf_{i \rightarrow \infty} f_i$$

□

Remarks 2.24:

(1) Let $x \in X$. Since the sequence $(f_i(x))$ converges in \mathbb{R} if and only if it is a Cauchy sequence, we have

$$\begin{aligned} \exists \lim_{i \rightarrow \infty} f_i(x) \in \mathbb{R} &\iff \forall \varepsilon > 0 \ \exists k \in \mathbb{N} \text{ such that } |f_i(x) - f_j(x)| < \varepsilon \ \forall i, j \geq k \\ &\iff \forall \varepsilon > 0 \ \exists k \in \mathbb{N} \text{ such that } |f_{k+i}(x) - f_k(x)| < \varepsilon \ \forall i \\ &\iff \forall m \ \exists k \in \mathbb{N} \text{ such that } |f_{k+i}(x) - f_k(x)| < \frac{1}{m} \ \forall i. \end{aligned}$$

That is

$$\{x \in X : \exists \lim_{i \rightarrow \infty} f_i(x) \in \mathbb{R}\} = \bigcap_{m=1}^{\infty} \bigcup_{k=1}^{\infty} \bigcap_{i=1}^{\infty} \{x \in X : |f_{k+i}(x) - f_k(x)| < \frac{1}{m}\}.$$

(2) By de Morgan's law, we have

$$\begin{aligned} \{x \in X : \nexists \lim_{i \rightarrow \infty} f_i(x) \in \mathbb{R}\} &= X \setminus \{x \in X : \exists \lim_{i \rightarrow \infty} f_i(x) \in \mathbb{R}\} \\ &= X \setminus \bigcap_{m=1}^{\infty} \bigcup_{k=1}^{\infty} \bigcap_{i=1}^{\infty} \{x \in X : |f_{k+i}(x) - f_k(x)| < \frac{1}{m}\} \\ &= \bigcup_{m=1}^{\infty} \left(X \setminus \bigcup_{k=1}^{\infty} \bigcap_{i=1}^{\infty} \{x \in X : |f_{k+i}(x) - f_k(x)| < \frac{1}{m}\} \right) \\ &= \bigcup_{m=1}^{\infty} \bigcap_{k=1}^{\infty} \left(X \setminus \bigcap_{i=1}^{\infty} \{x \in X : |f_{k+i}(x) - f_k(x)| < \frac{1}{m}\} \right) \\ &= \bigcup_{m=1}^{\infty} \bigcap_{k=1}^{\infty} \bigcup_{i=1}^{\infty} (X \setminus \{x \in X : |f_{k+i}(x) - f_k(x)| < \frac{1}{m}\}) \\ &= \bigcup_{m=1}^{\infty} \bigcap_{k=1}^{\infty} \bigcup_{i=1}^{\infty} \{x \in X : |f_{k+i}(x) - f_k(x)| \geq \frac{1}{m}\}. \end{aligned}$$

(3) Observe that, for extended real valued functions, the set

$$\{x \in X : |f_{k+i}(x) - f_k(x)| < \frac{1}{m}\}$$

consists of points $x \in X$ at which $|f_{k+i}(x) - f_k(x)|$ exists and is less than $\frac{1}{m}$. It may be an empty set. For example, if $f_{k+i}(x) = f_k(x) = \infty$ for every $x \in X$, then $|f_{k+i}(x) - f_k(x)|$ does not exist for any $x \in X$. However, if $(f_i(x))$ converges in \mathbb{R} , there exists k such that $f_i(x) \in \mathbb{R}$ for every $i \geq k$ and

$$\{x \in X : |f_{k+i}(x) - f_k(x)| < \frac{1}{m}\} \neq \emptyset$$

for every i .

(4) We note that

$$\lim_{i \rightarrow \infty} f_i(x) = \infty \iff \forall m \in \mathbb{N} \ \exists k \in \mathbb{N} \text{ such that } f_i(x) > m \ \forall i \geq k$$

This implies

$$\{x \in X : \lim_{i \rightarrow \infty} f_i(x) = \infty\} = \bigcap_{m=1}^{\infty} \bigcup_{k=1}^{\infty} \bigcap_{i=1}^{\infty} \{x \in X : f_{k+i}(x) > m\}.$$

Similarly

$$\{x \in X : \lim_{i \rightarrow \infty} f_i(x) = -\infty\} = \bigcap_{m=1}^{\infty} \bigcup_{k=1}^{\infty} \bigcap_{i=1}^{\infty} \{x \in X : f_{k+i}(x) < -m\}.$$

All sets above are measurable if $f_i : X \rightarrow [-\infty, \infty]$, $i = 1, 2, \dots$, are μ -measurable functions. The set where the sequence does not converge is also measurable, since it is the complement of the union of the sets above.

Example 2.25. Assume that $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable. Then f is continuous and thus Lebesgue measurable. Moreover, the difference quotients

$$g_i(x) = \frac{f(x + \frac{1}{i}) - f(x)}{\frac{1}{i}}, \quad i = 1, 2, \dots,$$

are continuous and thus Lebesgue measurable. Hence

$$f' = \lim_{i \rightarrow \infty} g_i$$

is a Lebesgue measurable function. Note that f' is not necessarily continuous.

Reason. The function $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 \sin(\frac{1}{x}), & x \neq 0, \\ 0, & x = 0, \end{cases}$$

is differentiable everywhere, but f' is not continuous at $x = 0$. ■

2.6 Almost everywhere

Sets of measure zero are negligible sets in the measure theory. In other words, an outer measure does not see sets of measure zero. Sets of measure zero are measurable with respect to an outer measure by Remark 1.6 (4), that is, an outer measure is complete, see Definition 1.15. Thus sets of measure zero do not affect measurability of a set (exercise). Measure theory is very flexible, but the price we have to pay is that we obtain information only up to sets of measure zero by measure theoretical tools.

Definition 2.26. Let μ be an outer measure on X . A property is said to hold μ -almost everywhere in X , if it holds in $X \setminus A$ for a set $A \subset X$ with $\mu(A) = 0$. It is sometimes denoted that the property holds μ -a.e.

Remark 2.27. Almost everywhere is called “almost surely” in probability theory.

Examples 2.28:

- (1) The function $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \chi_{(0,\infty)}(x)$, is continuous almost everywhere, because the set of discontinuity $\{0\}$ has Lebesgue measure zero.
- (2) The function $x \mapsto |x|$ is differentiable almost everywhere, because the set of non-differentiable points $\{0\}$ has Lebesgue measure zero.
- (3) Many useful functions such as

$$f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = \frac{\sin(x)}{x} \quad \text{and} \quad f : \mathbb{R}^n \rightarrow \mathbb{R}, f(x) = |x|^{-\alpha}, \alpha > 0,$$

are defined only almost everywhere with respect to the Lebesgue measure.

- (4) Let $\alpha > 0$. The function $f : \mathbb{R}^n \rightarrow [-\infty, \infty]$,

$$f(x) = \begin{cases} |x|^{-\alpha}, & x \neq 0, \\ \infty, & x = 0, \end{cases}$$

is finite almost everywhere, because the infinity set $f^{-1}(\{\infty\}) = \{0\}$ has Lebesgue measure zero.

Lemma 2.29. Assume that $f : X \rightarrow [-\infty, \infty]$ is a μ -measurable function. If $g : X \rightarrow [-\infty, \infty]$ is a function with $f = g$ μ -almost everywhere in X , then g is a μ -measurable function.

T H E M O R A L : Sets of measure zero do not affect measurability of a function. In case $f = g$ μ -almost everywhere, we do not usually distinguish f from g . Measure theoretically they are the same function. To be very formal, we could define an equivalence relation

$$f \sim g \iff f = g \quad \mu\text{-almost everywhere.}$$

Proof. Let $A = \{x \in X : f(x) \neq g(x)\}$ and $a \in \mathbb{R}$. By assumption $\mu(A) = 0$ and thus A is a μ -measurable set. Then

$$\begin{aligned} \{x \in X : g(x) > a\} &= \{x \in A : g(x) > a\} \cup \{x \in X \setminus A : g(x) > a\} \\ &= \{x \in A : g(x) > a\} \cup \{x \in X \setminus A : f(x) > a\}, \end{aligned}$$

since $g(x) = f(x)$ for every $x \in X \setminus A$. We claim that both sets on the right-hand side are μ -measurable, which implies that $\{x \in X : g(x) > a\}$ is a μ -measurable set and, consequently, that g is a μ -measurable function. Since

$$0 \leq \mu(\{x \in A : g(x) > a\}) \leq \mu(A) = 0,$$

we have $\mu(\{x \in A : g(x) > a\}) = 0$ and thus $\{x \in A : g(x) > a\}$ is a μ -measurable set. On the other hand, since f is a μ -measurable function and $X \setminus A$ is a μ -measurable set, we conclude that

$$\{x \in X \setminus A : f(x) > a\} = \{x \in X : f(x) > a\} \cap (X \setminus A)$$

is a μ -measurable set. This completes the proof. \square

Remark 2.30. All properties of measurable functions can be relaxed to conditions that hold almost everywhere. For example, if $f_i : X \rightarrow [-\infty, \infty]$, $i = 1, 2, \dots$, are μ -measurable functions and

$$f = \lim_{i \rightarrow \infty} f_i$$

μ -almost everywhere, then f is a μ -measurable function. Moreover, if the functions f and g are defined almost everywhere, the functions $f + g$ and fg are defined only in the intersection of the domains of f and g . Since the union of two sets of measure zero is a set of measure zero the functions are defined almost everywhere.

Remark 2.31. We discuss property that holds almost everywhere on a measure space (X, \mathcal{M}, μ) , see Definition 1.13. A property of points of X is said to hold μ -almost everywhere in X , if there exists a set $A \in \mathcal{M}$ with $\mu(A) = 0$, such that A contains every point at which the property does to hold. Consider a property that holds μ -almost everywhere, and let B be the set of points in X at which it does not hold. Then it is not necessary that $B \in \mathcal{M}$, but that there exists a set $A \in \mathcal{M}$ with $B \subset A$ and $\mu(A) = 0$. If μ is a complete measure, then $B \in \mathcal{M}$, see Definition 1.15. Let (X, \mathcal{M}, μ) be a measure space that is not complete, let $A \in \mathcal{M}$ be a set with $\mu(A) = 0$ and let $B \subset A$ be a set with $B \notin \mathcal{M}$. Then $f = 0$ and $g = \chi_B$ satisfy $f = g$ in $X \setminus A$ and thus $f = g$ μ -almost everywhere in X . However, f is a measurable function, but g is not. Thus Lemma 2.29 does not hold for measures that are not complete. In addition, the sequence $f_i = 0$ converges to g μ -almost everywhere as $i \rightarrow \infty$, so that the limit of measurable functions that converge μ -almost everywhere is not necessarily μ -measurable. Recall that all outer measures are complete by Remark 1.6 (4) and these problems do not occur.

2.7 Approximation by simple functions

Next we consider the approximation of a measurable function with simple functions, which are the basic blocks in the definition of the integral.

Definition 2.32. A function $f : X \rightarrow \mathbb{R}$ is simple, if its range is a finite set $\{a_1, \dots, a_n\}$, $n \in \mathbb{N}$, and the preimages

$$f^{-1}(\{a_i\}) = \{x \in X : f(x) = a_i\}$$

are μ -measurable sets.

T H E M O R A L : A simple function is a linear combination of finitely many characteristic functions of pairwise disjoint measurable sets, since it can be written as a finite sum

$$f = \sum_{i=1}^n a_i \chi_{A_i},$$

where $A_i = f^{-1}(\{a_i\})$. Remark 2.2 (3) and Theorem 2.13 imply that a simple function is μ -measurable.

W A R N I N G : A simple function assumes only finitely many values, but the sets $A_i = f^{-1}(\{a_i\})$ may not be geometrically simple.

Reason. The function $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \chi_{\mathbb{Q}}(x)$ is simple with respect to the one dimensional Lebesgue measure, but it is discontinuous at every point. In particular, it is possible that a measurable function is discontinuous at every point and thus it does not have any regularity in this sense. ■




Figure 2.3: A simple function.

We discuss approximation properties of nonnegative measurable functions.

Theorem 2.33. Let $f : X \rightarrow [0, \infty]$ be a function. Then f is a μ -measurable function if and only if there exists an increasing sequence (f_i) of simple functions f_i , $i = 1, 2, \dots$, such that

$$f(x) = \lim_{i \rightarrow \infty} f_i(x)$$

for every $x \in X$.

T H E M O R A L : Every nonnegative measurable function can be approximated by an increasing sequence of simple functions. Thus simple functions are basic building blocks in measure and integration theory.

Proof. \Rightarrow For every $i = 1, 2, \dots$ partition $[0, i)$ into $i2^i$ dyadic intervals

$$I_{i,k} = \left[\frac{k-1}{2^i}, \frac{k}{2^i} \right), \quad k = 1, \dots, i2^i$$

of length $\frac{1}{2^i}$. The intervals $I_{i,k}$ form a partition of $[0, i)$ into pairwise disjoint intervals and

$$[0, \infty] = [i, \infty) \cup \bigcup_{k=1}^{i2^i} I_{i,k}.$$

Note that every interval $I_{i,k}$ in the i th generation is a union of two pairwise disjoint intervals in the $(i+1)$ th generation.

Let

$$A_{i,k} = f^{-1}(I_{i,k}) = \left\{ x \in X : \frac{k-1}{2^i} \leq f(x) < \frac{k}{2^i} \right\}, \quad k = 1, \dots, i2^i,$$

and

$$A_i = f^{-1}([i, \infty]) = \{x \in X : f(x) \geq i\}.$$

Lemma 2.4 implies that the sets $A_{i,k}$, $k = 1, \dots, i2^i$, and A_i are μ -measurable and they form a pairwise disjoint partition of X , that is,

$$X = A_i \cup \bigcup_{k=1}^{i2^i} A_{i,k}.$$

The approximating simple function is defined as

$$f_i(x) = \sum_{k=1}^{i2^i} \frac{k-1}{2^i} \chi_{A_{i,k}}(x) + i \chi_{A_i}(x).$$

Note that this function takes constant value $\frac{k-1}{2^i}$ on $A_{i,k}$, $k = 1, \dots, i2^i$, and i on A_i . By the construction, we have $0 \leq f_i(x) \leq f_{i+1}(x) \leq f(x)$ for every $x \in X$. This shows that the sequence (f_i) is increasing.

Recall that an increasing sequence converges in $[0, \infty]$. We claim that

$$\lim_{i \rightarrow \infty} f_i(x) = f(x)$$

for every $x \in X$. To this end, we observe that

$$|f(x) - f_i(x)| \leq \frac{1}{2^i}, \quad \text{if } x \in \bigcup_{k=1}^{i2^i} A_{i,k} = \{x \in X : f(x) < i\}$$

and

$$f_i(x) = i, \quad \text{if } x \in A_i = \{x \in X : f(x) \geq i\}.$$

Let $\varepsilon > 0$. If $f(x) < \infty$, there exists $i_\varepsilon \in \mathbb{N}$ such that $x \in \bigcup_{k=1}^{i_\varepsilon 2^i} A_{i,k} = \{x \in X : f(x) < i\}$ for every $i \geq i_\varepsilon$. It follows that

$$|f(x) - f_i(x)| \leq \frac{1}{2^i} \leq \varepsilon \quad \text{for every } i \geq \max\left\{i_\varepsilon, \frac{\log \varepsilon}{\log 2}\right\}.$$




Figure 2.4: Approximation by simple functions.

This implies that $f_i(x) \rightarrow f(x)$ for every $x \in \{x \in X : f(x) < \infty\}$ as $i \rightarrow \infty$. On the other hand, if $f(x) = \infty$, then $x \in \bigcap_{i=1}^{\infty} A_i$. Thus $f_i(x) \geq i$ for every $i = 1, 2, \dots$ and $f_i(x) \rightarrow \infty$ as $i \rightarrow \infty$. Thus $f_i(x) \rightarrow f(x)$ for every $x \in X$ as $i \rightarrow \infty$.

Follows from the fact that a pointwise limit of measurable functions is measurable, see Theorem 2.23. \square

Remark 2.34. As the proof above shows, the approximation by simple functions is based on a subdivision of the range instead of the domain, as in the case of step functions. The approximation procedure is compatible with the definition of a measurable function.

Next we consider sign-changing functions.

Corollary 2.35. The function $f : X \rightarrow [-\infty, \infty]$ is a μ -measurable function if and only if there exists a sequence (f_i) of simple functions $f_i, i = 1, 2, \dots$, such that

$$f(x) = \lim_{i \rightarrow \infty} f_i(x)$$

for every $x \in X$.

THE MORAL : A function is measurable if and only if it can be approximated pointwise by simple functions.

Proof. We use the decomposition $f = f^+ - f^-$. By Theorem 2.33 there are simple functions g_i and h_i , $i = 1, 2, \dots$, such that

$$f^+ = \lim_{i \rightarrow \infty} g_i \quad \text{and} \quad f^- = \lim_{i \rightarrow \infty} h_i.$$

The functions $f_i = g_i - h_i$ do as an approximation. \square

Remarks 2.36:

- (1) The sequence $(|f_i|)$ is increasing, that is, $|f_i| \leq |f_{i+1}| \leq |f|$ for every $i = 1, 2, \dots$, because $|f_i| = g_i + h_i$ and the sequences (g_i) and (h_i) are increasing.
- (2) If the limit function f is bounded, then the simple functions will converge uniformly to f in X .
- (3) This approximation holds for every function f , but in that case the simple functions may not be measurable.

2.8 Modes of convergence

Let us recall two classical modes for a sequence of functions $f_i : X \rightarrow \mathbb{R}$, $i = 1, 2, \dots$, to converge to a function $f : X \rightarrow \mathbb{R}$.

- (1) A sequence (f_i) converges pointwise to f , if

$$|f_i(x) - f(x)| \xrightarrow{i \rightarrow \infty} 0$$

for every $x \in X$. This means that for every $\varepsilon > 0$, there exists i_ε such that

$$|f_i(x) - f(x)| < \varepsilon$$

whenever $i \geq i_\varepsilon$. Note that i_ε depends on x and ε .

- (2) A sequence (f_i) converges uniformly to f in X , if

$$\sup_{x \in X} |f_i(x) - f(x)| \xrightarrow{i \rightarrow \infty} 0.$$

This means that for every $\varepsilon > 0$, there exists i_ε such that

$$|f_i(x) - f(x)| < \varepsilon$$

for every $x \in X$ whenever $i \geq i_\varepsilon$. In this case i_ε does not depend on x .

Example 2.37. A uniform convergence implies pointwise convergence, but the converse is not true. For example, let $f_i : \mathbb{R} \rightarrow \mathbb{R}$,

$$f_i(x) = \frac{x}{i}, \quad i = 1, 2, \dots$$

Then $f_i(x) \rightarrow 0$ for every $x \in \mathbb{R}$, but f_i does not converge uniformly to f in \mathbb{R} .

Reason. Let $\varepsilon > 0$, $x \in \mathbb{R}$ and $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = 0$. Then

$$|f_i(x) - f(x)| = |f_i(x)| = \frac{|x|}{i} < \varepsilon$$

whenever $i \geq i_\varepsilon$, where i_ε is the smallest positive integer that is larger or equal to $\frac{|x|}{\varepsilon}$. This shows that $f_i(x) \rightarrow 0$ for every $x \in \mathbb{R}$. Note that i_ε depends on x and ε . The sequence f_i does not converge uniformly to f in \mathbb{R} , since

$$\sup_{x \in \mathbb{R}} |f_i(x) - f(x)| = \infty$$

for every $i = 1, 2, \dots$. ■

T H E M O R A L : A uniform limit of continuous functions is continuous, that is, continuity is preserved under uniform convergence. In contrast, continuity is not preserved under pointwise convergence.

There are also other modes of convergence that are relevant in measure theory. For simplicity we will discuss only real-valued functions. The discussion can be extended to the case when $f_i : X \rightarrow [-\infty, \infty]$ and $f : X \rightarrow [-\infty, \infty]$ are μ -measurable functions with $|f_i| < \infty$ and $|f| < \infty$ μ -almost everywhere in X for every $i = 1, 2, \dots$. In this case we consider a set $A \subset X$ with $\mu(A) = 0$ such that $|f_i|$, $i = 1, 2, \dots$, and $|f|$ are finite in the complement of A and replace f_i and f by $g_i = f_i \chi_{X \setminus A}$ and $g = \chi_{X \setminus A} g$. This allows us to avoid expressions as $|f_i(x) - f(x)|$ when $|f_i(x)|$ or $|f(x)|$ is infinite.

Definition 2.38 (Convergence almost everywhere). A sequence (f_i) converges to f almost everywhere in X , if $f_i(x) \rightarrow f(x)$ for μ -almost every $x \in X$ as $i \rightarrow \infty$. In other words, there exists a set $A \subset X$ with $\mu(X \setminus A) = 0$ such that $f_i(x) \rightarrow f(x)$ for every $x \in A$ as $i \rightarrow \infty$.

T H E M O R A L : Almost everywhere convergence is pointwise convergence outside a set of measure zero.

Remark 2.39. $f_i \rightarrow f$ almost everywhere in X if and only if for every $\varepsilon > 0$ there exists a μ -measurable set $A \subset X$ such that $\mu(X \setminus A) < \varepsilon$ and $f_i \rightarrow f$ pointwise in A as $i \rightarrow \infty$.

Reason. \Rightarrow If $f_i \rightarrow f$ almost everywhere in X , there exists a set A with $\mu(A) = 0$ such that $f_i(x) \rightarrow f(x)$ for every $x \in X \setminus A$ as $i \rightarrow \infty$. The set A satisfies the required properties.

\Leftarrow Assume that for every $j = 1, 2, \dots$ there exists a μ -measurable set $A_j \subset X$ such that $\mu(X \setminus A_j) < \frac{1}{j}$ and $f_i(x) \rightarrow f(x)$ for every $x \in A_j$ as $i \rightarrow \infty$. Let $A = \bigcup_{j=1}^{\infty} A_j$. Then

$$\mu(X \setminus A) = \mu\left(X \setminus \bigcup_{j=1}^{\infty} A_j\right) = \mu\left(\bigcap_{j=1}^{\infty}(X \setminus A_j)\right) \leq \mu(X \setminus A_j) < \frac{1}{j}$$

for every $j = 1, 2, \dots$. This implies that $\mu(X \setminus A) = 0$. Let $x \in A = \bigcup_{j=1}^{\infty} A_j$. Then $x \in A_j$ for some j and $f_i(x) \rightarrow f(x)$ as $i \rightarrow \infty$. This shows that $f_i(x) \rightarrow f(x)$ for every $x \in A$ as $i \rightarrow \infty$. \blacksquare

Definition 2.40 (Almost uniform convergence). A sequence (f_i) converges almost uniformly to f in X , if for every $\varepsilon > 0$ there exists a μ -measurable set $A \subset X$ such that $\mu(X \setminus A) < \varepsilon$ and $f_i \rightarrow f$ uniformly in A as $i \rightarrow \infty$, that is,

$$\sup_{x \in A} |f_i(x) - f(x)| \xrightarrow{i \rightarrow \infty} 0.$$

THEOREM: Almost uniform convergence is uniform convergence outside a set of arbitrarily small measure.

Remark 2.41. If $f_i \rightarrow f$ almost uniformly in X , then $f_i \rightarrow f$ almost everywhere in X , see Remark 2.39.

Example 2.42. Let $f_i : [0, 1] \rightarrow \mathbb{R}$, $f_i(x) = x^i$, $i = 1, 2, \dots$, and

$$f(x) = \begin{cases} 0, & 0 \leq x < 1, \\ 1, & x = 1. \end{cases}$$

Then $f_i(x) \rightarrow f(x)$ for every $x \in [0, 1]$, but the sequence (f_i) does not converge to f uniformly in $[0, 1]$.

Reason. Let $\varepsilon > 0$ and $x \in [0, 1]$. Then

$$|f_i(x) - f(x)| = |f_i(x)| = x^i < \varepsilon$$

whenever $i \geq i_\varepsilon$, where i_ε is the smallest positive integer that is larger or equal to $\frac{\log \varepsilon}{\log x}$. For $x = 1$ we have $|f_i(x) - f(x)| = 0 < \varepsilon$. This shows that $f_i(x) \rightarrow 0$ for every $x \in [0, 1]$. Note that i_ε depends on x and ε . The sequence (f_i) does not converge uniformly to f in $[0, 1]$, since

$$\sup_{x \in [0, 1]} |f_i(x) - f(x)| = 1$$

for every $i = 1, 2, \dots$. \blacksquare

However, $f_i \rightarrow f$ almost uniformly in $[0, 1]$, since $f_i \rightarrow f$ uniformly in every $[0, 1 - \varepsilon]$ with $0 < \varepsilon < 1$. This example also shows that almost uniform convergence does not imply uniform convergence outside a set of measure zero.

Definition 2.43 (Convergence in measure). A sequence (f_i) converges to f in measure in X , if

$$\lim_{i \rightarrow \infty} \mu(\{x \in X : |f_i(x) - f(x)| \geq \varepsilon\}) = 0$$

for every $\varepsilon > 0$.

T H E M O R A L : It is instructive to compare almost uniform convergence with convergence in measure. If $f_i \rightarrow f$ in measure on X , then for every $\eta, \varepsilon > 0$ there exists a set $A = A_{\eta, \varepsilon, i}$ with $\mu(X \setminus A) < \eta$ and $i_{\eta, \varepsilon}$ such that $|f_i(x) - f(x)| < \varepsilon$ for every $x \in A$ whenever $i \geq i_{\eta, \varepsilon}$. Note that the sets A_i may vary with i , as in the case of a sliding sequence of functions below. If $f_i \rightarrow f$ almost uniformly in X , then for every $\eta, \varepsilon > 0$ there exist a set $A = A_{\eta, \varepsilon} \subset X$ such that $\mu(X \setminus A) < \eta$ and $i_{\eta, \varepsilon}$ such that $|f_i(x) - f(x)| < \varepsilon$ for every $x \in A$ whenever $i \geq i_{\varepsilon}$. Almost uniform convergence requires that a single set A will do for all sufficiently large indices, that is, the set A does not depend on i .

Remark 2.44. If $f_i \rightarrow f$ almost uniformly in X , then $f_i \rightarrow f$ in measure in X .

Reason. For $j = 1, 2, \dots$ there exists a μ -measurable set A_j such that $\mu(X \setminus A_j) < \frac{1}{j}$ and $f_i \rightarrow f$ uniformly in A_j . Let $\varepsilon > 0$. There exists i_{ε} such that

$$\sup_{x \in A_j} |f_i(x) - f(x)| < \varepsilon,$$

whenever $i \geq i_{\varepsilon}$. This implies

$$\mu(\{x \in X : |f_i(x) - f(x)| \geq \varepsilon\}) \leq \mu(X \setminus A_j) < \frac{1}{j},$$

whenever $i \geq i_{\varepsilon}$. It follows that

$$\limsup_{i \rightarrow \infty} \mu(\{x \in X : |f_i(x) - f(x)| \geq \varepsilon\}) \leq \frac{1}{j}.$$

By letting $j \rightarrow \infty$, we obtain

$$\lim_{i \rightarrow \infty} \mu(\{x \in X : |f_i(x) - f(x)| \geq \varepsilon\}) = 0.$$

■

Remark 2.45. The limit function is unique under convergence in measure, that is, if (f_i) converges in measure to f and (f_i) converges in measure to g , then $f = g$ μ -almost everywhere in X .

Reason. Let $\varepsilon > 0$. Then

$$\begin{aligned} |f(x) - g(x)| &= |f(x) - f_i(x) + f_i(x) - g(x)| \\ &\leq |f(x) - f_i(x)| + |f_i(x) - g(x)|. \end{aligned}$$

This implies that if $|f(x) - g(x)| \geq \varepsilon$, then $|f(x) - f_i(x)| \geq \frac{\varepsilon}{2}$ or $|f_i(x) - g(x)| \geq \frac{\varepsilon}{2}$. To see this assume that $|f(x) - f_i(x)| < \frac{\varepsilon}{2}$ and $|f_i(x) - g(x)| < \frac{\varepsilon}{2}$. Then the display above implies that $|f(x) - g(x)| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$. Thus we have

$$\begin{aligned} \{x \in X : |f(x) - g(x)| \geq \varepsilon\} \\ \subset \{x \in X : |f_i(x) - f(x)| \geq \frac{\varepsilon}{2}\} \cup \{x \in X : |f_i(x) - g(x)| \geq \frac{\varepsilon}{2}\} \end{aligned}$$

and consequently

$$\begin{aligned}\mu(\{x \in X : |f(x) - g(x)| \geq \varepsilon\}) \\ \leq \mu(\{x \in X : |f_i(x) - f(x)| \geq \frac{\varepsilon}{2}\}) + \mu(\{x \in X : |f_i(x) - g(x)| \geq \frac{\varepsilon}{2}\}).\end{aligned}$$

By letting $i \rightarrow \infty$, we have

$$\mu(\{x \in X : |f(x) - g(x)| \geq \varepsilon\}) = 0.$$

The claim follows, since

$$\{x \in X : f(x) \neq g(x)\} = \bigcup_{i=1}^{\infty} \{x \in X : |f(x) - g(x)| \geq \frac{1}{i}\}$$

and thus

$$\mu(\{x \in X : f(x) \neq g(x)\}) \leq \sum_{i=1}^{\infty} \mu(\{x \in X : |f(x) - g(x)| \geq \frac{1}{i}\}) = 0. \quad \blacksquare$$

Next we give examples which distinguish between the modes of convergence. In the following moving bump examples we have $X = \mathbb{R}$ with the Lebesgue measure.

Examples 2.46:

(1) (Escape to horizontal infinity) Let $f_i : \mathbb{R} \rightarrow \mathbb{R}$,

$$f_i(x) = \chi_{[i, i+1]}(x), \quad i = 1, 2, \dots$$

Then $f_i \rightarrow 0$ everywhere and thus almost everywhere in \mathbb{R} , but not uniformly, almost uniformly or in measure.

(2) (Escape to width infinity) Let $f_i : \mathbb{R} \rightarrow \mathbb{R}$,

$$f_i(x) = \frac{1}{i} \chi_{[0, i]}(x), \quad i = 1, 2, \dots$$

Then $f_i \rightarrow 0$ uniformly in \mathbb{R} .

(3) (Escape to vertical infinity) Let $f_i : \mathbb{R} \rightarrow \mathbb{R}$,

$$f_i(x) = i \chi_{[\frac{1}{i}, \frac{2}{i}]}(x), \quad i = 1, 2, \dots$$

Then $f_i \rightarrow 0$ pointwise, almost uniformly and in measure, but not uniformly in \mathbb{R} .

(4) (A sliding sequence of functions) Let $f_i : [0, 1] \rightarrow \mathbb{R}$, $i = 1, 2, \dots$, be defined by

$$f_{2^k+j}(x) = k \chi_{[\frac{j}{2^k}, \frac{j+1}{2^k}]}(x), \quad k = 0, 1, 2, \dots, \quad j = 0, 1, \dots, 2^k - 1.$$

Then

$$\limsup_{i \rightarrow \infty} f_i(x) = \infty \quad \text{and} \quad \liminf_{i \rightarrow \infty} f_i(x) = 0$$

for every $x \in [0, 1]$ and thus the pointwise limit does not exist at any point. However,

$$m(\{x \in [0, 1] : f_{2^k+j}(x) \geq \varepsilon\}) = m\left(\left[\frac{j}{2^k}, \frac{j+1}{2^k}\right]\right) = \frac{1}{2^k} \xrightarrow{k \rightarrow \infty} 0.$$

This shows that $f_i \rightarrow 0$ in measure on $[0, 1]$. Note that there are several converging subsequences. For example, $f_{2^i}(x) \rightarrow 0$ for every $x \neq 0$, although the original sequence diverges everywhere.




Figure 2.5: A sliding sequence of functions.

The next result shows that, for a sequence that converges in measure, a converging subsequence, as in the sliding sequence of functions above, always exists.

Theorem 2.47. Assume that $f_i \rightarrow f$ in measure. There exists a subsequence (f_{i_k}) such that $f_{i_k} \rightarrow f$ μ -almost everywhere.

Proof. Choose i_1 such that

$$\mu(\{x \in X : |f_{i_1}(x) - f(x)| \geq 1\}) < \frac{1}{2}.$$

Assume then that i_1, \dots, i_k have been chosen. Choose $i_{k+1} > i_k$ such that

$$\mu(\{x \in X : |f_{i_{k+1}}(x) - f(x)| \geq \frac{1}{k+1}\}) < \frac{1}{2^{k+1}}.$$

Let

$$A_j = \bigcup_{k=j}^{\infty} \{x \in X : |f_{i_k}(x) - f(x)| \geq \frac{1}{k}\}, \quad j = 1, 2, \dots$$

Clearly $A_{j+1} \subset A_j$ and denote $A = \bigcap_{j=1}^{\infty} A_j$. For every $j = 1, 2, \dots$, we have

$$\mu(A) \leq \mu(A_j) \leq \sum_{k=j}^{\infty} \frac{1}{2^k} = \frac{2}{2^j}$$

and by letting $j \rightarrow \infty$ we conclude $\mu(A) = 0$. By de Morgan's law

$$\begin{aligned} X \setminus A &= X \setminus \bigcap_{j=1}^{\infty} A_j = \bigcup_{j=1}^{\infty} (X \setminus A_j) \\ &= \bigcup_{j=1}^{\infty} \left(X \setminus \bigcup_{k=j}^{\infty} \{x \in X : |f_{i_k}(x) - f(x)| \geq \frac{1}{k}\} \right) \\ &= \bigcup_{j=1}^{\infty} \bigcap_{k=j}^{\infty} (X \setminus \{x \in X : |f_{i_k}(x) - f(x)| \geq \frac{1}{k}\}) \\ &= \bigcup_{j=1}^{\infty} \bigcap_{k=j}^{\infty} \{x \in X : |f_{i_k}(x) - f(x)| < \frac{1}{k}\}. \end{aligned}$$

For every $x \in X \setminus A$ there exists j such that

$$|f_{i_k}(x) - f(x)| < \frac{1}{k} \quad \text{for every } k \geq j.$$

This implies that $f_{i_k}(x) \rightarrow f(x)$ for every $x \in X \setminus A$ as $k \rightarrow \infty$. \square

Remark 2.48. The following assertions are valid for almost everywhere convergence, almost uniform convergence and convergence in measure.

- (1) The limit function f is μ -measurable. See Theorem 2.23 and the discussion in Section 2.6.
- (2) The limit function f is unique up to a set of μ -measure zero.
- (3) Convergence is not affected by changing f_i or f on a set of μ -measure zero.

Remark 2.49. Convergence almost everywhere is called “convergence almost surely” in probability theory and convergence in measure is called “convergence in probability”.

2.9 Egoroff's and Lusin's theorems

The next result gives the main motivation for almost uniform convergence.

Theorem 2.50 (Egoroff's theorem). Assume that $\mu(X) < \infty$. Let $f_i : X \rightarrow [-\infty, \infty]$ be μ -measurable functions with $|f_i| < \infty$ μ -almost everywhere for every $i = 1, 2, \dots$ such that $f_i \rightarrow f$ almost everywhere in X as $i \rightarrow \infty$ and $|f| < \infty$ μ -almost everywhere. Then $f_i \rightarrow f$ almost uniformly in X as $i \rightarrow \infty$.

T H E M O R A L : Almost uniform convergence and almost everywhere convergence are equivalent in a space with finite measure.




Figure 2.6: Comparison of modes of convergence.

Proof. Let $\varepsilon > 0$ and

$$A_{j,k} = \bigcup_{i=j}^{\infty} \left\{ x \in X : |f_i(x) - f(x)| > \frac{1}{2^k} \right\}, \quad j, k = 1, 2, \dots$$

As in Remark 2.24 we have

$$\begin{aligned} \exists \lim_{i \rightarrow \infty} f_i(x) = f(x) \in \mathbb{R} &\iff \forall \varepsilon > 0 \ \exists i_{\varepsilon} \in \mathbb{N} \text{ such that } |f_i(x) - f(x)| < \varepsilon \ \forall i \geq i_{\varepsilon} \\ &\iff \forall k \in \mathbb{N} \ \exists i_k \in \mathbb{N} \text{ such that } |f_i(x) - f(x)| \leq \frac{1}{2^k} \ \forall i \geq i_k. \end{aligned}$$

By a repeated application of de Morgan's law, we have

$$\begin{aligned} \{x \in X : \exists \lim_{i \rightarrow \infty} f_i(x) = f(x) \in \mathbb{R}\} &= \bigcap_{k=1}^{\infty} \bigcup_{j=1}^{\infty} \bigcap_{i=j}^{\infty} \left\{ x \in X : |f_i(x) - f(x)| \leq \frac{1}{2^k} \right\} \\ &= \bigcap_{k=1}^{\infty} \bigcup_{j=1}^{\infty} \bigcap_{i=j}^{\infty} \left(X \setminus \left\{ x \in X : |f_i(x) - f(x)| > \frac{1}{2^k} \right\} \right) \\ &= \bigcap_{k=1}^{\infty} \bigcup_{j=1}^{\infty} \left(X \setminus \bigcup_{i=j}^{\infty} \left\{ x \in X : |f_i(x) - f(x)| > \frac{1}{2^k} \right\} \right) \\ &= \bigcap_{k=1}^{\infty} \bigcup_{j=1}^{\infty} (X \setminus A_{j,k}) = \bigcap_{k=1}^{\infty} (X \setminus \bigcap_{j=1}^{\infty} A_{j,k}) \\ &= X \setminus \bigcup_{k=1}^{\infty} \bigcap_{j=1}^{\infty} A_{j,k}. \end{aligned}$$

Since $f_i \rightarrow f \in \mathbb{R}$ almost everywhere in X , we conclude that

$$\mu\left(\bigcup_{k=1}^{\infty} \bigcap_{j=1}^{\infty} A_{j,k}\right) = 0.$$

Since $\bigcap_{j=1}^{\infty} A_{j,k} \subset \bigcup_{k=1}^{\infty} \bigcap_{j=1}^{\infty} A_{j,k}$ for every $k = 1, 2, \dots$, we have

$$\mu\left(\bigcap_{j=1}^{\infty} A_{j,k}\right) = 0$$

for every $k = 1, 2, \dots$. Since $A_{j+1,k} \subset A_{j,k}$ are μ -measurable sets for every $j, k = 1, 2, \dots$ and $\mu(X) < \infty$, Theorem 1.22 (2) implies that

$$\lim_{j \rightarrow \infty} \mu(A_{j,k}) = \mu\left(\bigcap_{j=1}^{\infty} A_{j,k}\right) = 0$$

for every $k = 1, 2, \dots$. Thus for every $k = 1, 2, \dots$ there exists j_k such that

$$\mu(A_{j_k,k}) < \frac{\varepsilon}{2^{k+1}}.$$

Let $A = X \setminus \bigcup_{k=1}^{\infty} A_{j_k,k}$. Then we have

$$\mu(X \setminus A) = \mu\left(\bigcup_{k=1}^{\infty} A_{j_k,k}\right) \leq \sum_{k=1}^{\infty} \mu(A_{j_k,k}) \leq \sum_{k=1}^{\infty} \frac{\varepsilon}{2^{k+1}} < \varepsilon.$$

By a repeated application de Morgan's law, we have

$$\begin{aligned} A &= X \setminus \bigcup_{k=1}^{\infty} A_{j_k,k} = X \setminus \bigcup_{k=1}^{\infty} \bigcup_{i=j_k}^{\infty} \left\{x \in X : |f_i(x) - f(x)| > \frac{1}{2^k}\right\} \\ &= \bigcap_{k=1}^{\infty} \left(X \setminus \bigcup_{i=j_k}^{\infty} \left\{x \in X : |f_i(x) - f(x)| > \frac{1}{2^k}\right\} \right) \\ &= \bigcap_{k=1}^{\infty} \bigcap_{i=j_k}^{\infty} \left(X \setminus \left\{x \in X : |f_i(x) - f(x)| > \frac{1}{2^k}\right\} \right) \\ &= \bigcap_{k=1}^{\infty} \bigcap_{i=j_k}^{\infty} \left\{x \in X : |f_i(x) - f(x)| \leq \frac{1}{2^k}\right\}. \end{aligned}$$

For every $k = 1, 2, \dots$ and $i \geq j_k$ we have

$$|f_i(x) - f(x)| \leq \frac{1}{2^k}$$

for every $x \in A$. This implies that $f_i \rightarrow f$ uniformly in A with $\mu(X \setminus A) < \varepsilon$. \square

Remarks 2.51:

- (1) Egoroff's theorem does not hold true without the assumption that $|f| < \infty$ μ -almost everywhere. For example, if $|f_i| < \infty$ everywhere, but $|f| = \infty$ on a set of positive measure, then $|f_i - f| = \infty$ on a set of positive measure.
- (2) Egoroff's theorem does not hold true without the assumption $\mu(X) < \infty$. For example, let $f_i : \mathbb{R} \rightarrow \mathbb{R}$, $f_i(x) = \frac{x}{i}$, $i = 1, 2, \dots$. Then $f_i(x) \rightarrow 0$ for every $x \in \mathbb{R}$, but f_i does not converge almost uniformly to f in \mathbb{R} .

Reason. We show that for some $\varepsilon > 0$ there does not exist Lebesgue measurable set $A \subset \mathbb{R}$ with $m(\mathbb{R} \setminus A) < \varepsilon$ such that (f_i) converges uniformly in A . Let $0 < \varepsilon < \infty$ and let $A \subset \mathbb{R}$ be a Lebesgue measurable set with $m(\mathbb{R} \setminus A) < \varepsilon$. Then $m(A) = \infty$ and thus A is unbounded. Since A is unbounded, there exist $x_k \in A$, $k = 1, 2, \dots$, such that $|x_k| \rightarrow \infty$ as $k \rightarrow \infty$. Thus

$$\sup_k |f_i(x_k) - f(x_k)| = \sup_k |f_i(x_k)| = \sup_k \frac{|x_k|}{i} = \infty$$

for every $i = 1, 2, \dots$. This shows that (f_i) does not converge uniformly to f in the set $\{x_k : k = 1, 2, \dots\}$ and, consequently, (f_i) does not converge uniformly to f in A containing the set $\{x_k : k = 1, 2, \dots\}$. ■

(3) If $\mu(X) = \infty$, we can apply Egoroff's theorem for μ -measurable subsets $A \subset X$ with $\mu(A) < \infty$. As far as \mathbb{R}^n is concerned, a sequence (f_i) is said to converge locally uniformly to f , if $f_i \rightarrow f$ uniformly on every bounded set $A \subset \mathbb{R}^n$. Equivalently, we could require that for every point $x \in \mathbb{R}^n$ there is a ball $B(x, r)$, with $r > 0$, such that $f_i \rightarrow f$ uniformly in $B(x, r)$. Let us rephrase Egoroff's theorem for the Lebesgue measure, or a more general Radon measure, on \mathbb{R}^n . Let (f_i) be a sequence measurable functions with $|f_i| < \infty$ almost everywhere for every $i = 1, 2, \dots$ such that $f_i \rightarrow f$ almost everywhere in \mathbb{R}^n as $i \rightarrow \infty$ and $|f| < \infty$ almost everywhere. Then for every $\varepsilon > 0$ there exists a measurable set $A \subset \mathbb{R}^n$ such that the measure of A is at most ε and $f_i \rightarrow f$ locally uniformly in $\mathbb{R}^n \setminus A$.

Remark 2.52. Relations of measurable sets and functions to standard open sets and continuous functions are summarized in Littlewood's three principles.

- (1) Every measurable set is almost open (Theorem 1.48 and Theorem 1.73).
- (2) Pointwise convergence is almost uniform (Egoroff's theorem 2.50).
- (3) A measurable function is almost continuous (Lusin's theorem 2.53).

Here the word “almost” has to be understood measure theoretically.

The following result is related to Littlewood's third principle. We shall prove it only in the case $X = \mathbb{R}^n$, but the result also holds in more general metric spaces with the same proof.

Theorem 2.53 (Lusin's theorem). Let μ be a Borel regular outer measure on \mathbb{R}^n , $A \subset \mathbb{R}^n$ a μ -measurable set such that $\mu(A) < \infty$ and $f : \mathbb{R}^n \rightarrow [-\infty, \infty]$ be a μ -measurable function such that $|f| < \infty$ μ -almost everywhere. For every $\varepsilon > 0$ there exists a compact set $K \subset A$ such that $\mu(A \setminus K) < \varepsilon$ and that the restricted function $f|_K$ is a continuous function.

T H E M O R A L : A measurable function can be measure theoretically approximated by a continuous function.

Remarks 2.54:

- (1) The assumption $\mu(A) < \infty$ can be removed if the compact set in the claim is replaced with a closed set.
- (2) Lusin's theorem gives a characterization for measurable functions.

Reason. Assume that for every $i = 1, 2, \dots$, there exists a compact set $K_i \subset A$ such that $\mu(A \setminus K_i) < \frac{1}{i}$ and $f|_{K_i}$ is continuous. Let $B = \bigcup_{i=1}^{\infty} K_i$ and $N = A \setminus B$. Then

$$\begin{aligned} 0 &\leq \mu(N) = \mu(A \setminus B) = \mu\left(A \setminus \bigcup_{i=1}^{\infty} K_i\right) \\ &= \mu\left(\bigcap_{i=1}^{\infty}(A \setminus K_i)\right) \leq \mu(A \setminus K_i) < \frac{1}{i} \end{aligned}$$

for every $i = 1, 2, \dots$. Thus $\mu(N) = 0$. We have

$$\{x \in A : f(x) > a\} = \{x \in B : f(x) > a\} \cup \{x \in A \setminus B : f(x) > a\}$$

for every $a \in \mathbb{R}$. The set $\{x \in B : f(x) > a\}$ is μ -measurable, since f is continuous in B and $\{x \in A \setminus B : f(x) > a\}$ is μ -measurable, since it is a set of measure zero. This implies that f is μ -measurable in A . \blacksquare

Proof. By excluding the set $\{x \in \mathbb{R}^n : |f(x)| = \infty\}$, which is of measure zero, we may assume that $f : \mathbb{R}^n \rightarrow \mathbb{R}$. For every $i = 1, 2, \dots$, let $B_{i,j}$, $j = 1, 2, \dots$, be pairwise disjoint Borel sets such that

$$\bigcup_{j=1}^{\infty} B_{i,j} = \mathbb{R} \quad \text{and} \quad \text{diam}(B_{i,j}) < \frac{1}{i}.$$

Let $A_{i,j} = A \cap f^{-1}(B_{i,j})$. The sets $A_{i,j}$ are μ -measurable by Lemma 2.5. Moreover

$$\begin{aligned} A &= A \cap f^{-1}(\mathbb{R}) = A \cap f^{-1}\left(\bigcup_{j=1}^{\infty} B_{i,j}\right) \\ &= \bigcup_{j=1}^{\infty} (A \cap f^{-1}(B_{i,j})) = \bigcup_{j=1}^{\infty} A_{i,j}, \quad i = 1, 2, \dots \end{aligned}$$

Since $\mu(A) < \infty$, $\nu = \mu|_A$ is a Radon measure by Lemma 1.46. By Corollary 1.51, there exists a compact set $K_{i,j} \subset A_{i,j}$ such that

$$\nu(A_{i,j} \setminus K_{i,j}) < \frac{\varepsilon}{2^{i+j}}.$$

Then

$$\begin{aligned} \mu\left(A \setminus \bigcup_{j=1}^{\infty} K_{i,j}\right) &= \nu\left(A \setminus \bigcup_{j=1}^{\infty} K_{i,j}\right) = \nu\left(\bigcup_{j=1}^{\infty} A_{i,j} \setminus \bigcup_{j=1}^{\infty} K_{i,j}\right) \\ &= \nu\left(\bigcup_{j=1}^{\infty}\left(A_{i,j} \setminus \bigcup_{j=1}^{\infty} K_{i,j}\right)\right) \leq \nu\left(\bigcup_{j=1}^{\infty}(A_{i,j} \setminus K_{i,j})\right) \\ &\leq \sum_{j=1}^{\infty} \nu(A_{i,j} \setminus K_{i,j}) < \frac{\varepsilon}{2^i}. \end{aligned}$$

Since $\mu(A) < \infty$, by Theorem 1.22 (2) we have

$$\begin{aligned}\lim_{k \rightarrow \infty} \mu\left(A \setminus \bigcup_{j=1}^k K_{i,j}\right) &= \lim_{k \rightarrow \infty} \mu\left(\bigcap_{j=1}^k (A \setminus K_{i,j})\right) \\ &= \mu\left(\bigcap_{j=1}^{\infty} (A \setminus K_{i,j})\right) = \mu\left(A \setminus \bigcup_{j=1}^{\infty} K_{i,j}\right) < \frac{\varepsilon}{2^i}.\end{aligned}$$

Thus there exists an index k_i such that

$$\mu\left(A \setminus \bigcup_{j=1}^{k_i} K_{i,j}\right) < \frac{\varepsilon}{2^i}$$

As a union of finitely many compact sets, the set $K_i = \bigcup_{j=1}^{k_i} K_{i,j}$ is compact. For every i, j , we choose a point $\alpha_{i,j} \in B_{i,j}$. Then we define a function $g_i : K_i \rightarrow \mathbb{R}$ by

$$g_i(x) = \alpha_{i,j}, \quad \text{when } x \in K_{i,j}, \quad j = 1, \dots, k_i.$$

Since $K_{i,1}, \dots, K_{i,k_i}$ are pairwise disjoint compact sets, we have

$$\text{dist}(K_{i,j}, K_{i,l}) > 0 \quad \text{when } j \neq l.$$

This implies that g_i is continuous in K_i and

$$|f(x) - g_i(x)| < \frac{1}{i} \quad \text{for every } x \in K_i,$$

since $f(K_{i,j}) \subset f(A_{i,j}) \subset B_{i,j}$ and $\text{diam}(B_{i,j}) < \frac{1}{i}$. The set $K = \bigcap_{i=1}^{\infty} K_i$ is compact and

$$\begin{aligned}\mu(A \setminus K) &= \mu\left(A \setminus \bigcap_{i=1}^{\infty} K_i\right) = \mu\left(\bigcup_{i=1}^{\infty} (A \setminus K_i)\right) \\ &\leq \sum_{i=1}^{\infty} \mu(A \setminus K_i) < \varepsilon \sum_{i=1}^{\infty} \frac{1}{2^i} = \varepsilon.\end{aligned}$$

Since

$$|f(x) - g_i(x)| < \frac{1}{i} \quad \text{for every } x \in K, \quad i = 1, 2, \dots,$$

we see that $g_i \rightarrow f$ uniformly in K . The function f is continuous in K as a uniform limit of continuous functions. \square

W A R N I N G : Note carefully, that $f|_K$ denotes the restriction of f to K . Theorem 2.53 states that f is continuous viewed as a function defined only on the set K . This does not immediately imply that f defined as a function on A is continuous at the points in K .

Reason. $f : [0, 1] \rightarrow \mathbb{R}$, $f(x) = \chi_{\mathbb{Q}}(x)$ is discontinuous at every point of $[0, 1]$. However, $f|_{[0,1] \cap \mathbb{Q}} = 1$ and $f|_{[0,1] \setminus \mathbb{Q}} = 0$ are continuous functions. It is an exercise to construct the compact set in Lusin's theorem for this function. \blacksquare

Keeping this example in mind, we are now ready to prove a stronger result.

Corollary 2.55. Let μ be a Borel regular outer measure on \mathbb{R}^n , $A \subset \mathbb{R}^n$ a μ -measurable set such that $\mu(A) < \infty$ and $f : \mathbb{R}^n \rightarrow [-\infty, \infty]$ be a μ -measurable function such that $|f| < \infty$ μ -almost everywhere. Then for every $\varepsilon > 0$ there exists a continuous function $\bar{f} : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$\mu(\{x \in A : \bar{f}(x) \neq f(x)\}) < \varepsilon.$$

W A R N I N G : The corollary does not imply that there is a continuous function $\bar{f} : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $\bar{f}(x) = f(x)$ μ -almost everywhere, see Example 3.34.

Proof. Let $\varepsilon > 0$. By Lusin's theorem 2.53, there exists a compact set $K \subset A$ such that $\mu(A \setminus K) < \varepsilon$ and $f|_K$ is continuous. By Tieze's extension theorem there exists a continuous function $\bar{f} : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $\bar{f}(x) = f(x)$ for every $x \in K$. We refer to [2] for Tieze's theorem. Then

$$\mu(\{x \in A : \bar{f}(x) \neq f(x)\}) \leq \mu(A \setminus K) < \varepsilon,$$

which implies the claim. \square

Remarks 2.56:

- (1) Tieze's extension theorem holds in metric spaces. Let F be a closed subset of a metric space X and suppose that $f : F \rightarrow \mathbb{R}$ is a continuous function. Then f can be extended to a continuous function $\bar{f} : X \rightarrow \mathbb{R}$ defined everywhere on X . Moreover, if $|f(x)| \leq M$ for every $x \in F$, then $|\bar{f}(x)| \leq M$ for every $x \in X$. See [2].
- (2) It is essential in Tieze's extension theorem that the set F is closed.

Reason. The function $f : (0, 1] \rightarrow \mathbb{R}$, $f(x) = \sin \frac{1}{x}$ is a continuous function on $(0, 1]$, but it cannot be extended to a continuous function on $[0, 1]$. \blacksquare

Example 2.57. Let $A \subset \mathbb{R}^n$ be a Lebesgue measurable set with $m(A) < \infty$ and $f = \chi_A$. By Theorem 1.73, for every $\varepsilon > 0$, there exists a compact set $K \subset A$ and an open set $G \supset A$ such that $m(A \setminus K) < \frac{\varepsilon}{2}$ and $m(G \setminus A) < \frac{\varepsilon}{2}$. As in Remark 1.32, let

$$\bar{f}(x) = \frac{\text{dist}(x, \mathbb{R}^n \setminus G)}{\text{dist}(x, \mathbb{R}^n \setminus G) + \text{dist}(x, K)}.$$

Then \bar{f} is a continuous function in \mathbb{R}^n and

$$m(\{x \in \mathbb{R}^n : \bar{f}(x) \neq f(x)\}) \leq m(G \setminus A) + m(A \setminus K) < \varepsilon.$$

In this special case, the function in the previous corollary can be constructed explicitly.

Integral is first defined for nonnegative simple functions, then for nonnegative measurable functions and finally for signed functions. Integral has all basic properties one might expect and it behaves well with respect to limits, as the monotone convergence theorem, Fatou's lemma and the dominated convergence theorem show.

3

Integration

3.1 Integral of a nonnegative simple function

Let A be a μ -measurable set. It is natural to define integral of the characteristic function of A as

$$\int_X \chi_A d\mu = \mu(A).$$

The same approach can be applied for simple functions. Recall that a function $f : X \rightarrow \mathbb{R}$ is simple, if its range is a finite set $\{a_1, \dots, a_n\}$, $n \in \mathbb{N}$, and the preimages

$$f^{-1}(\{a_i\}) = \{x \in X : f(x) = a_i\}$$

are μ -measurable sets, see Definition 2.32. A simple function is a linear combination of finitely many characteristic functions of μ -measurable sets, since it can be written as a finite sum

$$f = \sum_{i=1}^n a_i \chi_{A_i}, \quad n \in \mathbb{N},$$

where $A_i = f^{-1}(\{a_i\})$. Remark 2.2 (3) and Theorem 2.13 imply that a simple function is μ -measurable. This is called the canonical representation of a simple function. Observe that the sets A_i are disjoint and thus for each $x \in X$ there is only one nonzero term in the sum above.

Definition 3.1. Let μ be a measure on X and let $f = \sum_{i=1}^n a_i \chi_{A_i}$ be the canonical representation of a nonnegative simple function. Then

$$\int_X f d\mu = \int_X f(x) d\mu(x) = \sum_{i=1}^n a_i \mu(A_i).$$

If for some i we have $a_i = 0$ and $\mu(A_i) = \infty$, we define $a_i \mu(A_i) = 0$.

T H E M O R A L : The definition of integral of a simple functions is based on a subdivision of the range instead of the domain, as in the case of step functions. This is compatible with the definition of measurable function.

Example 3.2. The function $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \chi_{\mathbb{Q}}(x)$ is simple with respect to the one dimensional Lebesgue measure and $\int_{\mathbb{R}} f(x) dx = 0$.

Remarks 3.3:

- (1) For a nonnegative simple function f , we have $0 \leq \int_X f d\mu \leq \infty$.
- (2) If f is a simple function and A is μ -measurable subset of X , then $f\chi_A$ is a simple function.
- (3) (Compatibility with the measure) If A is a μ -measurable subset of X , then $\int_X \chi_A d\mu = \mu(A)$.
- (4) The representation of a simple function is not in general unique in the sense that there may be several ways to write a function as a finite linear combination of characteristic functions of pairwise disjoint measurable sets. For example, $\chi_X = \chi_{X \setminus A} + \chi_A$ for every μ -measurable set $A \subset X$. However, the definition of integral of a nonnegative simple function is independent of the representation of the function.

Reason. Let $f = \sum_{i=1}^n a_i \chi_{A_i}$ be the canonical representation of a nonnegative simple function f and let $f = \sum_{j=1}^m b_j \chi_{B_j}$ be another representation, where b_j are nonnegative real numbers and B_j pairwise disjoint μ -measurable subsets of X with $\bigcup_{j=1}^m B_j = X$. Additivity of μ on pairwise disjoint μ -measurable sets and the fact that $a_i = b_j$ if $A_i \cap B_j \neq \emptyset$ imply

$$\begin{aligned} \sum_{i=1}^n a_i \mu(A_i) &= \sum_{i=1}^n \sum_{j=1}^m a_i \mu(A_i \cap B_j) \\ &= \sum_{i=1}^n \sum_{j=1}^m b_j \mu(A_i \cap B_j) \\ &= \sum_{j=1}^m \sum_{i=1}^n b_j \mu(A_i \cap B_j) = \sum_{j=1}^m b_j \mu(B_j). \end{aligned} \quad \blacksquare$$

- (5) If A is a μ -measurable subset of X , then we define

$$\int_A f d\mu = \int_X f \chi_A d\mu.$$

If $f = \sum_{i=1}^n a_i \chi_{A_i}$ is the canonical representation of a nonnegative simple function, then

$$\int_A f d\mu = \sum_{i=1}^n a_i \mu(A_i \cap A).$$

Observe that the sum on the right-hand side is not necessarily the canonical form of $f\chi_A$. However, the integral of a nonnegative simple function is independent of the representation.

(6) If f and g are nonnegative simple functions such that $f = g$ μ -almost everywhere, then $\int_X f d\mu = \int_X g d\mu$. Note that the converse is not true.
 (7) $\int_X f d\mu = 0$ if and only if $f = 0$ μ -almost everywhere.

Lemma 3.4. Assume that f and g are nonnegative simple functions on X .

(1) (Monotonicity in sets) If A and B are μ -measurable sets with $A \subset B$, then $\int_A f d\mu \leq \int_B f d\mu$.
 (2) (Homogeneity) $\int_X af d\mu = a \int_X f d\mu$, $a \geq 0$.
 (3) (Linearity) $\int_X (f+g) d\mu = \int_X f d\mu + \int_X g d\mu$.
 (4) (Monotonicity in functions) $f \leq g$ implies $\int_X f d\mu \leq \int_X g d\mu$.

Proof. Claims (1) and (2) are clear. To prove (3), let

$$\int_X f d\mu = \sum_{i=1}^n a_i \mu(A_i) \quad \text{and} \quad \int_X g d\mu = \sum_{j=1}^m b_j \mu(B_j)$$

be the canonical representations of f and g . We have $X = \bigcup_{i=1}^n A_i = \bigcup_{j=1}^m B_j$. Then $f+g$ is a nonnegative simple function. The sets

$$C_{i,j} = A_i \cap B_j, \quad i = 1, \dots, n, j = 1, \dots, m,$$

are pairwise disjoint and $X = \bigcup_{i=1}^n \bigcup_{j=1}^m C_{i,j}$ and each of the functions f and g are constant on each set $C_{i,j}$. Thus

$$\begin{aligned} \int_X (f+g) d\mu &= \sum_{i=1}^n \sum_{j=1}^m (a_i + b_j) \mu(C_{i,j}) \\ &= \sum_{i=1}^n \sum_{j=1}^m a_i \mu(A_i \cap B_j) + \sum_{i=1}^n \sum_{j=1}^m b_j \mu(A_i \cap B_j) \\ &= \sum_{i=1}^n a_i \mu(A_i) + \sum_{j=1}^m b_j \mu(B_j) \\ &= \int_X f d\mu + \int_X g d\mu. \end{aligned}$$

To prove (4) we note that on the sets $C_{i,j} = A_i \cap B_j$ we have $f = a_i \leq b_j = g$ and thus

$$\int_X f d\mu = \sum_{i=1}^n \sum_{j=1}^m a_i \mu(C_{i,j}) \leq \sum_{i=1}^n \sum_{j=1}^m b_j \mu(C_{i,j}) = \int_X g d\mu. \quad \square$$

Remark 3.5. Since the sum in the representation of a nonnegative simple function consists of finitely many terms, it is clear that integral inherits the properties of measure. For example, if A_i , $i = 1, 2, \dots$ are μ -measurable sets, then

$$\int_{\bigcup_{i=1}^{\infty} A_i} f d\mu \leq \sum_{i=1}^{\infty} \int_{A_i} f d\mu$$

and

$$\int_{\bigcup_{i=1}^{\infty} A_i} f d\mu = \sum_{i=1}^{\infty} \int_{A_i} f d\mu$$

if the sets A_i , $i = 1, 2, \dots$, are pairwise disjoint. Moreover, if $A_i \supset A_{i+1}$ for every i and $\int_{A_1} f d\mu < \infty$, we have

$$\int_{\bigcap_{i=1}^{\infty} A_i} f d\mu = \lim_{i \rightarrow \infty} \int_{A_i} f d\mu$$

and finally if $A_i \subset A_{i+1}$ for every i , then

$$\int_{\bigcup_{i=1}^{\infty} A_i} f d\mu = \lim_{i \rightarrow \infty} \int_{A_i} f d\mu.$$

Reason. We only prove the last claim, since the other cases are similar (exercise).

Let

$$f = \sum_{j=1}^n a_j \chi_{\{x \in X : f(x) = a_j\}}.$$

By Theorem 1.22 (1), we have

$$\begin{aligned} \int_{\bigcup_{i=1}^{\infty} A_i} f d\mu &= \sum_{j=1}^n a_j \mu\left(\left\{x \in \bigcup_{i=1}^{\infty} A_i : f(x) = a_j\right\}\right) \\ &= \sum_{j=1}^n a_j \lim_{i \rightarrow \infty} \mu(\{x \in A_i : f(x) = a_j\}) \\ &= \lim_{i \rightarrow \infty} \sum_{j=1}^n a_j \mu(\{x \in A_i : f(x) = a_j\}) = \lim_{i \rightarrow \infty} \int_{A_i} f d\mu. \end{aligned} \quad \blacksquare$$

3.2 Integral of a nonnegative measurable function

Integral of an arbitrary nonnegative measurable function is defined through an approximation by simple functions.

Definition 3.6. Let $f : X \rightarrow [0, \infty]$ be a nonnegative μ -measurable function. Integral of f with respect to μ is

$$\int_X f d\mu = \sup \left\{ \int_X g d\mu : g \text{ is simple and } 0 \leq g(x) \leq f(x) \text{ for every } x \in X \right\}.$$

A nonnegative function is integrable, if

$$\int_X f d\mu < \infty.$$

T H E M O R A L : Integral is defined for all nonnegative measurable functions. Observe, that integral may be infinite.

Remarks 3.7:

(1) As before, if A is a μ -measurable subset of X , then we let

$$\int_A f d\mu = \int_X \chi_A f d\mu.$$

Thus by taking the zero extension, we may assume that the function is defined on the whole space.

(2) The definition is consistent with the one for nonnegative simple functions.
 (3) If $\mu(X) = 0$, then $\int_X f d\mu = 0$ for every f .

We collect a few basic properties of integral of a nonnegative function below.

Lemma 3.8. Let $f, g : X \rightarrow [0, \infty]$ be μ -measurable functions.

(1) (Monotonicity in sets) If A and B are μ -measurable sets with $A \subset B$, then

$$\int_A f d\mu \leq \int_B f d\mu.$$

(2) (Homogeneity) $\int_X af d\mu = a \int_X f d\mu$, $a \geq 0$.
 (3) (Linearity) $\int_X (f+g) d\mu = \int_X f d\mu + \int_X g d\mu$.
 (4) (Monotonicity in functions) $f \leq g$ implies $\int_X f d\mu \leq \int_X g d\mu$.
 (5) (Tchebyshev's inequality)

$$\mu(\{x \in X : f(x) > a\}) \leq \frac{1}{a} \int_X f d\mu$$

for every $a > 0$.

W A R N I N G : Some of the claims do not necessarily hold true for a sign changing function. However, we may consider the absolute value of a function instead. We shall return to this later.

Proof. (1) Follows immediately from the corresponding property for nonnegative simple functions.

(2) If $a = 0$, then

$$\int_X (0f) d\mu = \int_X 0 d\mu = 0 = 0 \int_X f d\mu.$$

Let then $a > 0$. If g is simple and $0 \leq g \leq f$, then ag is a nonnegative simple function with $ag \leq af$. It follows that

$$a \int_X g d\mu = \int_X ag d\mu \leq \int_X af d\mu.$$

Taking supremum over all such functions g implies that

$$a \int_X f d\mu \leq \int_X af d\mu.$$

The reverse inequality follows by applying the inequality above, since

$$\int_X af d\mu = a \left(\frac{1}{a} \int_X af d\mu \right) \leq a \int_X \frac{1}{a} (af) d\mu = a \int_X f d\mu.$$

(3) Exercise, see also Remark 3.14 (2) below.

(4) Let h be a simple function with $0 \leq h(x) \leq f(x)$ for every $x \in X$. Then $0 \leq h(x) \leq g(x)$ for every $x \in X$ and thus $\int_X h d\mu \leq \int_X g d\mu$. By taking supremum over all such functions h we have $\int_X f d\mu \leq \int_X g d\mu$.

(5) Since $f \geq a \chi_{\{x \in X : f(x) > a\}}$, we have

$$a\mu(\{x \in X : f(x) > a\}) = \int_X a \chi_{\{x \in X : f(x) > a\}} d\mu \leq \int_X f d\mu.$$

□

Lemma 3.9. Let $f : X \rightarrow [0, \infty]$ be a μ -measurable function.

- (1) (Vanishing) $\int_X f d\mu = 0$ if and only if $f = 0$ μ -almost everywhere.
- (2) (Finiteness) $\int_X f d\mu < \infty$ implies $f < \infty$ μ -almost everywhere.

W A R N I N G : The claim (1) is not necessarily true for a sign changing function. The converse of claim (2) is not true: $f < \infty$ μ -almost everywhere does not imply that $\int_X f d\mu < \infty$.

Proof. (1) \implies Let

$$A_i = \{x \in X : f(x) > \frac{1}{i}\}, \quad i = 1, 2, \dots$$

By Tchebyshev's inequality in Lemma 3.8 (5), we have

$$0 \leq \mu(A_i) \leq i \int_X f d\mu = 0$$

which implies that $\mu(A_i) = 0$ for every $i = 1, 2, \dots$. Thus

$$\mu(\{x \in X : f(x) > 0\}) = \mu\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty} \mu(A_i) = 0.$$

\implies Since $\mu(\{x \in X : f(x) > 0\}) = 0$, we have

$$0 = \int_{\{x \in X : f(x) > 0\}} \infty d\mu = \int_X \infty \chi_{\{x \in X : f(x) > 0\}} d\mu \geq \int_X f d\mu \geq 0.$$

Thus $\int_X f d\mu = 0$. Another way to prove this claim is to use the definition of integral directly (exercise).

(2) By Tchebyshev's inequality in Lemma 3.8 (5), we have

$$\mu(\{x \in X : f(x) = \infty\}) \leq \mu(\{x \in X : f(x) > i\}) \leq \frac{1}{i} \int_X f d\mu \xrightarrow{i \rightarrow \infty} 0,$$

since $\int_X f d\mu < \infty$. □

Lemma 3.10. Let $f, g : X \rightarrow [0, \infty]$ be μ -measurable functions. If $f = g$ μ -almost everywhere then

$$\int_X f d\mu = \int_X g d\mu.$$

T H E M O R A L : A redefinition of a function on a set of measure zero does not affect the integral.

W A R N I N G : The converse of the claim does not hold: $\int_X f d\mu = \int_X g d\mu$ does not in general imply that $f = g$ μ -almost everywhere. However, if $\int_A f d\mu = \int_A g d\mu$ for every μ -measurable set A , then $f = g$ μ -almost everywhere (exercise).s

Proof. Let $N = \{x \in X : f(x) \neq g(x)\}$. Then $\mu(N) = 0$ and thus

$$\int_N f d\mu = 0 = \int_N g d\mu.$$

It follows that

$$\begin{aligned} \int_X f d\mu &= \int_{X \setminus N} f d\mu + \int_N f d\mu = \int_{X \setminus N} f d\mu \\ &= \int_{X \setminus N} g d\mu = \int_{X \setminus N} g d\mu + \int_N g d\mu = \int_X g d\mu. \end{aligned} \quad \square$$

3.3 Monotone convergence theorem

Assume that $f_i : X \rightarrow [0, \infty]$, $i = 1, 2, \dots$, are μ -measurable functions and that $f_i \rightarrow f$ either everywhere or μ -almost everywhere as $i \rightarrow \infty$. Then f is a μ -measurable function, see Section 2.6. Next we discuss the question whether

$$\int_X \lim_{i \rightarrow \infty} f_i d\mu = \lim_{i \rightarrow \infty} \int_X f_i d\mu.$$

In other words, is it possible to switch the order of limit and integral? We begin with moving bump examples that we have already discussed in Example 2.46.

Examples 3.11:

(1) (Escape to horizontal infinity) Let $f_i : \mathbb{R} \rightarrow \mathbb{R}$,

$$f_i(x) = \chi_{[i, i+1]}(x), \quad i = 1, 2, \dots$$

Then $\lim_{i \rightarrow \infty} f_i(x) = 0$ as $i \rightarrow \infty$ for every $x \in \mathbb{R}$, but

$$\int_{\mathbb{R}} \lim_{i \rightarrow \infty} f_i dm = \int_{\mathbb{R}} 0 dm = 0 < 1 = m([i, i+1]) = \lim_{i \rightarrow \infty} \int_{\mathbb{R}} f_i dm.$$

(2) (Escape to width infinity) Let $f_i : \mathbb{R} \rightarrow \mathbb{R}$,

$$f_i(x) = \frac{1}{i} \chi_{[0, i]}(x), \quad i = 1, 2, \dots$$

Then $f_i \rightarrow 0$ uniformly in \mathbb{R} , but

$$\int_{\mathbb{R}} \lim_{i \rightarrow \infty} f_i dm = \int_{[0, \infty)} 0 dm = 0 < 1 = \frac{1}{i} m([0, i]) = \lim_{i \rightarrow \infty} \int_{\mathbb{R}} f_i dm.$$

(3) (Escape to vertical infinity) Let $f_i : \mathbb{R} \rightarrow \mathbb{R}$,

$$f_i(x) = i \chi_{[\frac{1}{i}, \frac{2}{i}]}(x), \quad i = 1, 2, \dots$$

Then $\lim_{i \rightarrow \infty} f_i(x) = 0$ for every $x \in \mathbb{R}$, but

$$\int_{\mathbb{R}} \lim_{i \rightarrow \infty} f_i dm = \int_{\mathbb{R}} 0 dm = 0 < 1 = im([\frac{1}{i}, \frac{2}{i}]) = \lim_{i \rightarrow \infty} \int_{\mathbb{R}} f_i dm.$$

(4) Let $f_i : \mathbb{R} \rightarrow \mathbb{R}$, $f_i(x) = \frac{1}{i}$, $i = 1, 2, \dots$ Then $\lim_{i \rightarrow \infty} f_i(x) = 0$ for every $x \in \mathbb{R}$, but

$$\int_{\mathbb{R}} \lim_{i \rightarrow \infty} f_i dm = 0 < \infty = \lim_{i \rightarrow \infty} \int_{\mathbb{R}} f_i dm.$$

This example shows that the following monotone convergence theorem does not hold for decreasing sequences of functions.

The next convergence result will be very useful.

Theorem 3.12 (Monotone convergence theorem). If $f_i : X \rightarrow [0, \infty]$ are μ -measurable functions such that $f_i \leq f_{i+1}$, $i = 1, 2, \dots$, then

$$\int_X \lim_{i \rightarrow \infty} f_i d\mu = \lim_{i \rightarrow \infty} \int_X f_i d\mu$$

T H E M O R A L : The order of taking limit and integral can be switched for an increasing sequence of nonnegative measurable functions. This tells that mass is preserved in a pointwise increasing limit of nonnegative functions.

Remarks 3.13:

- (1) It is enough to assume that $f_i \leq f_{i+1}$ almost everywhere.
- (2) The increasing limits

$$\lim_{i \rightarrow \infty} f_i \quad \text{and} \quad \lim_{i \rightarrow \infty} \int_X f_i d\mu$$

exist, but may be infinite.

- (3) In the special case when $f_i = \chi_{A_i}$, where A_i is μ -measurable and $A_i \subset A_{i+1}$, the monotone convergence theorem reduces to the upwards monotone convergence result for measures, see Theorem 1.22 (1).

Proof. Let $f = \lim_{i \rightarrow \infty} f_i$. Theorem 2.23 implies that the function f is μ -measurable. By monotonicity and the definition of integral, we have

$$\int_X f_i d\mu \leq \int_X f_{i+1} d\mu \leq \int_X f d\mu$$

for every $i = 1, 2, \dots$ This implies that the limit exists and

$$\lim_{i \rightarrow \infty} \int_X f_i d\mu \leq \int_X f d\mu.$$

To prove the reverse inequality, let g be a nonnegative simple function with $g \leq f$. Let $0 < t < 1$ and

$$A_i = \{x \in X : f_i(x) \geq tg(x)\}, \quad i = 1, 2, \dots$$

By Lemma 2.10 and Lemma 2.4, the sets A_i are μ -measurable and $A_i \subset A_{i+1}$, $i = 1, 2, \dots$.

Claim: $\bigcup_{i=1}^{\infty} A_i = X$.

Reason. \square Since $A_i \subset X$, $i = 1, 2, \dots$, we have $\bigcup_{i=1}^{\infty} A_i \subset X$.

\square For every $x \in X$, either $f(x) \leq tg(x)$ or $f(x) > tg(x)$. If $f(x) \leq tg(x)$, then $f(x) \leq tg(x) \leq tf(x)$ and, since $0 < t < 1$ and $0 \leq f(x) \leq g(x) < \infty$, we have $f(x) = 0$. In this case $x \in A_i$ for every $i = 1, 2, \dots$. In the other hand, if $f(x) > tg(x)$, then $f(x) = \lim_{i \rightarrow \infty} f_i(x) > tg(x)$. Thus there exists i such that $f_i(x) > tg(x)$ and consequently $x \in A_i$. This shows that $x \in \bigcup_{i=1}^{\infty} A_i$ for every $x \in X$. \blacksquare

Thus

$$\begin{aligned} \int_X f_i d\mu &\geq \int_{A_i} f_i d\mu \geq \int_{A_i} tg d\mu = t \int_{A_i} g d\mu \\ &\xrightarrow{i \rightarrow \infty} t \int_{\bigcup_{i=1}^{\infty} A_i} g d\mu = t \int_X g d\mu. \end{aligned}$$

Here we used the measure properties of the integral of nonnegative simple functions, see Remark 3.5. It follows that

$$\lim_{i \rightarrow \infty} \int_X f_i d\mu \geq t \int_X g d\mu.$$

By taking the supremum over all nonnegative simple functions $g \leq f$ we have

$$\lim_{i \rightarrow \infty} \int_X f_i d\mu \geq t \int_X f d\mu$$

and the claim follows by letting $t \rightarrow 1$. \square

Remarks 3.14:

(1) By Theorem 2.33 for every nonnegative μ -measurable function f there exists an increasing sequence f_i , $i = 1, 2, \dots$, of simple functions such that

$$f(x) = \lim_{i \rightarrow \infty} f_i(x)$$

for every $x \in X$. By the monotone convergence theorem we have

$$\int_X f d\mu = \lim_{i \rightarrow \infty} \int_X f_i d\mu.$$

Conversely, if f_i , $i = 1, 2, \dots$, are nonnegative simple functions such that $f_i \leq f_{i+1}$ and $f = \lim_{i \rightarrow \infty} f_i$, then

$$\int_X f d\mu = \lim_{i \rightarrow \infty} \int_X f_i d\mu.$$

Moreover, this limit is independent of the approximating sequence.

(2) Let $f, g : X \rightarrow [0, \infty]$ be μ -measurable functions. Let (f_i) be an increasing sequence of nonnegative simple functions $f_i, i = 1, 2, \dots$, such that $f_i(x) \rightarrow f(x)$ for every $x \in X$ as $i \rightarrow \infty$ and let (g_i) be an increasing sequence of nonnegative simple functions $g_i, i = 1, 2, \dots$, such that $g_i(x) \rightarrow g(x)$ for every $x \in X$ as $i \rightarrow \infty$, see Theorem 2.33. Then

$$f(x) + g(x) = \lim_{i \rightarrow \infty} (f_i(x) + g_i(x))$$

and the monotone convergence theorem implies

$$\begin{aligned} \int_X (f + g) d\mu &= \int_X \lim_{i \rightarrow \infty} (f_i + g_i) d\mu \\ &= \lim_{i \rightarrow \infty} \int_X (f_i + g_i) d\mu \\ &= \lim_{i \rightarrow \infty} \int_X f_i d\mu + \lim_{i \rightarrow \infty} \int_X g_i d\mu \\ &= \int_X f d\mu + \int_X g d\mu. \end{aligned}$$

This shows the approximation by simple functions can be used to prove properties of the integral, compare to Lemma 3.8.

Example 3.15. The monotone convergence theorem can be used to compute limits of certain nonnegative integrals.

(1) Consider

$$\lim_{i \rightarrow \infty} \int_0^1 \frac{i}{1 + i\sqrt{x}} dx.$$

Let $f_i : [0, 1] \rightarrow \mathbb{R}$,

$$f_i(x) = \frac{i}{1 + i\sqrt{x}}$$

for every $i = 1, 2, \dots$. Note that $0 \leq f_i(x) \leq f_{i+1}(x)$ for every $0 \leq x \leq 1$ and $i = 1, 2, \dots$. The monotone convergence theorem implies

$$\begin{aligned} \lim_{i \rightarrow \infty} \int_0^1 \frac{i}{1 + i\sqrt{x}} dx &= \lim_{i \rightarrow \infty} \int_0^1 f_i(x) dx = \int_0^1 \lim_{i \rightarrow \infty} f_i(x) dx \\ &= \int_0^1 \lim_{i \rightarrow \infty} \frac{i}{1 + i\sqrt{x}} dx = \int_0^1 \lim_{i \rightarrow \infty} \frac{1}{\frac{1}{i} + \sqrt{x}} dx \\ &= \int_0^1 \frac{1}{\sqrt{x}} dx = 2. \end{aligned}$$

(2) Consider

$$\lim_{x \rightarrow 0^+} \int_0^\infty \frac{e^{-xt}}{1 + t^2} dt.$$

Note that $\lim_{x \rightarrow 0^+} f(x)$ exists if and only if $\lim_{i \rightarrow \infty} f(x_i)$ exists for all decreasing sequences (x_i) with $x_i \searrow 0$ and is independent of the sequence. Let (x_i) be such a sequence and let

$$f_i(t) = \frac{e^{-x_i t}}{1 + t^2}$$

for every $t \geq 0$ and $i = 1, 2, \dots$. Since (x_i) is a decreasing sequence, we have $0 \leq f_i(t) \leq f_{i+1}(t)$ for every $t \geq 0$ and $i = 1, 2, \dots$. We can thus apply the monotone convergence theorem, use the fact that $\lim_{i \rightarrow \infty} x_i = 0$ and continuity of elementary functions to obtain

$$\begin{aligned} \lim_{i \rightarrow \infty} \int_0^\infty \frac{e^{-x_i t}}{1+t^2} dt &= \lim_{i \rightarrow \infty} \int_0^\infty f_i(t) dt = \int_0^\infty \lim_{i \rightarrow \infty} f_i(t) dt \\ &= \int_0^\infty \lim_{i \rightarrow \infty} \frac{e^{-x_i t}}{1+t^2} dt = \int_0^\infty \frac{e^0}{1+t^2} dt \\ &= \int_0^\infty \frac{1}{1+t^2} dt = \arctan t \Big|_{t=0}^\infty = \frac{\pi}{2}. \end{aligned}$$

In the examples above we assumed that the familiar rules for computing integrals hold. This is a consequence of the fact that Lebesgue integral is equal to Riemann integral for bounded continuous functions, see Section 3.9.

Corollary 3.16. Let $f_i : X \rightarrow [0, \infty]$, $i = 1, 2, \dots$, be nonnegative μ -measurable functions. Then

$$\int_X \sum_{i=1}^\infty f_i d\mu = \sum_{i=1}^\infty \int_X f_i d\mu.$$

T H E M O R A L : The order of taking sum and integral can be switched for a sequence of nonnegative measurable functions. In other words, a series of nonnegative measurable functions can be integrated termwise.

Proof. Let $s_n = f_1 + \dots + f_n = \sum_{i=1}^n f_i$ be the n th partial sum and

$$f = \lim_{n \rightarrow \infty} s_n = \sum_{i=1}^\infty f_i.$$

The functions s_n , $n = 1, 2, \dots$, form an increasing sequence of nonnegative μ -measurable functions. By the monotone convergence theorem

$$\begin{aligned} \int_X f d\mu &= \int_X \lim_{n \rightarrow \infty} s_n d\mu = \lim_{n \rightarrow \infty} \int_X s_n d\mu \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \int_X f_i d\mu = \sum_{i=1}^\infty \int_X f_i d\mu. \end{aligned} \quad \square$$

Remark 3.17. Let f be a nonnegative μ -measurable function on X . Let

$$v(A) = \int_A f d\mu$$

for any μ -measurable set A . Then v is a measure.

Reason. It is clear that v is nonnegative and that $v(\emptyset) = 0$. We show that v is countably additive on pairwise disjoint μ -measurable sets. Let A_i , $i = 1, 2, \dots$, be

pairwise disjoint μ -measurable sets and let $f_i = f\chi_{A_i}$. By Corollary 3.16, we have

$$\begin{aligned}\sum_{i=1}^{\infty} \nu(A_i) &= \sum_{i=1}^{\infty} \int_X f_i d\mu = \int_X \sum_{i=1}^{\infty} f_i d\mu = \int_X \sum_{i=1}^{\infty} f\chi_{A_i} d\mu \\ &= \int_X f \sum_{i=1}^{\infty} \chi_{A_i} d\mu = \int_X f \chi_{\bigcup_{i=1}^{\infty} A_i} d\mu \\ &= \int_{\bigcup_{i=1}^{\infty} A_i} f d\mu = \nu\left(\bigcup_{i=1}^{\infty} A_i\right).\end{aligned}\blacksquare$$

This provides a useful method of constructing measures related to a nonnegative density function f .

T H E M O R A L : An integral can be computed by partitioning a set into a countably many pairwise disjoint measurable sets or representing a set as a countable union (or intersection) of measurable sets.

3.4 Fatou's lemma

The next convergence result holds without monotonicity assumptions.

Theorem 3.18 (Fatou's lemma). If $f_i : X \rightarrow [0, \infty]$, $i = 1, 2, \dots$, are μ -measurable functions, then

$$\int_X \liminf_{i \rightarrow \infty} f_i d\mu \leq \liminf_{i \rightarrow \infty} \int_X f_i d\mu.$$

T H E M O R A L : Fatou's lemma tells that mass can be destroyed but not created in a pointwise limit of nonnegative functions as the moving bump examples show.

Remarks 3.19:

- (1) The power of Fatou's lemma is that there are no assumptions on the convergences. In particular, the limits

$$\lim_{i \rightarrow \infty} f_i \quad \text{and} \quad \lim_{i \rightarrow \infty} \int_X f_i d\mu$$

do not necessarily have to exist, but the corresponding limes inferiors exist for nonnegative functions.

- (2) The moving bump examples show that a strict inequality may occur in Fatou's lemma. By considering the moving bump example with a negative sign, we observe that the nonnegativity assumption is necessary.

Proof. Recall that

$$\liminf_{i \rightarrow \infty} f_i(x) = \sup_{j \geq 1} (\inf_{i \geq j} f_i(x)) = \lim_{j \rightarrow \infty} (\inf_{i \geq j} f_i(x)) = \lim_{j \rightarrow \infty} g_j(x),$$

where $g_j = \inf_{i \geq j} f_i$. The functions g_j , $j = 1, 2, \dots$, form an increasing sequence of μ -measurable functions. By the monotone convergence theorem

$$\begin{aligned} \int_X \liminf_{i \rightarrow \infty} f_i d\mu &= \int_X \lim_{j \rightarrow \infty} g_j d\mu \\ &= \lim_{j \rightarrow \infty} \int_X g_j d\mu \\ &\leq \liminf_{i \rightarrow \infty} \int_X f_i d\mu, \end{aligned}$$

where the last inequality follows from the fact that $g_i \leq f_i$. \square

3.5 Integral of a signed function

The integral of a signed function will be defined by considering the positive and negative parts of the function. Recall that $f^+ = \max\{f, 0\} \geq 0$ and $f^- = -\min\{f, 0\} \geq 0$ and $f = f^+ - f^-$. By Lemma 2.12 a function $f : X \rightarrow [-\infty, \infty]$ is μ -measurable if and only if f^+ and f^- are μ -measurable.

Definition 3.20. Let $f : X \rightarrow [-\infty, \infty]$ be a μ -measurable function. If either $\int_X f^- d\mu < \infty$ or $\int_X f^+ d\mu < \infty$, then the integral of f in X is defined as

$$\int_X f d\mu = \int_X f^+ d\mu - \int_X f^- d\mu.$$

Moreover, the function f is integrable in X , if both $\int_X f^- d\mu < \infty$ and $\int_X f^+ d\mu < \infty$. In this case we denote $f \in L^1(X; \mu)$.

T H E M O R A L : The integral can be defined if either positive or negative parts have a finite integral. For an integrable function both have finite integrals.

Remark 3.21. (Triangle inequality) A function f is integrable if and only if $|f|$ is integrable, that is, $\int_X |f| d\mu < \infty$. In this case, we have

$$\left| \int_X f d\mu \right| \leq \int_X |f| d\mu.$$

T H E M O R A L : $f \in L^1(X; \mu) \iff \int_X |f| d\mu < \infty$.

Reason. \Rightarrow Assume that f is integrable in X . Since $|f| = f^+ + f^-$ and integral is linear on nonnegative functions, we have

$$\int_X |f| d\mu = \int_X f^+ d\mu + \int_X f^- d\mu < \infty.$$

It follows that $|f|$ is integrable in X .

\Leftarrow Assume that $|f|$ is integrable in X . Since $0 \leq f^+ \leq |f|$ and $0 \leq f^- \leq |f|$ we have

$$\int_X f^+ d\mu \leq \int_X |f| d\mu < \infty \quad \text{and} \quad \int_X f^- d\mu \leq \int_X |f| d\mu < \infty.$$

It follows that f is integrable in X .

Moreover,

$$\begin{aligned} \left| \int_X f d\mu \right| &= \left| \int_X f^+ d\mu - \int_X f^- d\mu \right| \leq \left| \int_X f^+ d\mu \right| + \left| \int_X f^- d\mu \right| \\ &= \int_X f^+ d\mu + \int_X f^- d\mu = \int_X (f^+ + f^-) d\mu = \int_X |f| d\mu. \end{aligned} \quad \blacksquare$$

Remarks 3.22:

- (1) If $f \in L^1(X; \mu)$, then $|f| \in L^1(X; \mu)$ and by Lemma 3.9 we have $|f| < \infty$ μ -almost everywhere in X .
- (2) (Majorant principle) Let $f : X \rightarrow [-\infty, \infty]$ be a μ -measurable function. If there exists a nonnegative integrable function g such that $|f| \leq g$ μ -almost everywhere, then f is integrable.

Reason. $\int_X |f| d\mu \leq \int_X g d\mu < \infty$. ■

- (3) A measure space (X, \mathcal{M}, μ) with $\mu(X) = 1$ is called a probability or sample space, μ a probability measure and sets belonging to \mathcal{M} events. A probability measure is often denoted by P . In probability theory a measurable function is called a random variable, denoted for example by X . The integral is called the expectation or mean of X and it is written as

$$E(X) = \int X(\omega) dP(\omega).$$

Next we give some examples of integrals.

Examples 3.23:

- (1) Let $X = \mathbb{R}^n$ and μ be the Lebesgue measure. We shall discuss properties of the Lebesgue measure in detail later.
- (2) Let $X = \mathbb{N}$ and μ be the counting measure. Then all functions are μ -measurable. Observe that a function $f : \mathbb{N} \rightarrow \mathbb{R}$ is a sequence of real numbers with $x_i = f(i)$, $i = 1, 2, \dots$. Then

$$\int_X f d\mu = \sum_{i=1}^{\infty} f(i) = \sum_{i=1}^{\infty} x_i$$

and $f \in L^1(X; \mu)$ if and only if

$$\int_X |f| d\mu = \sum_{i=1}^{\infty} |f(i)| < \infty.$$

In other words, the integral is the sum of the series and integrability means that the series converges absolutely.

- (3) Let $x_0 \in X$ be a fixed point and recall that the Dirac measure at x_0 is defined as

$$\mu(A) = \begin{cases} 1, & x_0 \in A, \\ 0, & x_0 \notin A. \end{cases}$$

Then all functions are μ -measurable. Moreover,

$$\int_X f d\mu = f(x_0)$$

and $f \in L^1(X; \mu)$ if and only if

$$\int_X |f| d\mu = |f(x_0)| < \infty.$$

Lemma 3.24. Let $f, g : X \rightarrow [-\infty, \infty]$ be integrable functions.

- (1) (Homogeneity) $\int_X af d\mu = a \int_X f d\mu$, $a \in \mathbb{R}$.
- (2) (Linearity) $\int_X (f+g) d\mu = \int_X f d\mu + \int_X g d\mu$.
- (3) (Monotonicity in functions) $f \leq g$ implies $\int_X f d\mu \leq \int_X g d\mu$.
- (4) (Vanishing) $\mu(X) = 0$ implies $\int_X f d\mu = 0$.
- (5) (Almost everywhere equivalence) If $f = g$ μ -almost everywhere in X , then $\int_X f d\mu = \int_X g d\mu$.

W A R N I N G : If A and B are μ -measurable sets with $A \subset B$, then it does not in general follow that

$$\int_A f d\mu \leq \int_B f d\mu.$$

Thus monotonicity in sets does not necessarily hold for sign-changing functions.

Remark 3.25. Since $f, g \in L^1(X; \mu)$, we have $|f| < \infty$ and $|g| < \infty$ μ -almost everywhere in X . Thus $f+g$ is defined μ -almost everywhere in X .

Proof. (1) If $a \geq 0$, then $(af)^+ = af^+$ and $(af)^- = af^-$. This implies

$$\int_X (af)^+ d\mu = a \int_X f^+ d\mu \quad \text{and} \quad \int_X (af)^- d\mu = a \int_X f^- d\mu$$

The claim follows from this. If $a < 0$, then $(af)^+ = (-a)f^-$ and $(af)^- = (-a)f^+$ and the claim follows as above.

(2) Let $h = f+g$. Then h is defined almost everywhere and measurable. The pointwise inequality $|h| \leq |f| + |g|$ implies

$$\int_X |h| d\mu \leq \int_X |f| d\mu + \int_X |g| d\mu < \infty$$

and thus h is integrable. Note that in general $h^+ \neq f^+ + g^+$, but

$$h^+ - h^- = h = f + g = f^+ - f^- + g^+ - g^-$$

implies

$$h^+ + f^- + g^- = h^- + f^+ + g^+.$$

Both sides are nonnegative integrable functions. It follows that

$$\int_X h^+ d\mu + \int_X f^- d\mu + \int_X g^- d\mu = \int_X h^- d\mu + \int_X f^+ d\mu + \int_X g^+ d\mu$$

and since all integrals are finite we arrive at

$$\begin{aligned} \int_X h d\mu &= \int_X h^+ d\mu - \int_X h^- d\mu \\ &= \int_X f^+ d\mu - \int_X f^- d\mu + \int_X g^+ d\mu - \int_X g^- d\mu \\ &= \int_X f d\mu + \int_X g d\mu. \end{aligned}$$

(3) (1) and (2) imply that $g - f \geq 0$ is integrable and

$$\int_X g d\mu = \int_X f d\mu + \int_X (g - f) d\mu \geq \int_X f d\mu.$$

(4) $\mu(X) = 0$ implies $\int_X f^+ d\mu = 0$ and $\int_X f^- d\mu = 0$ and consequently $\int_X f d\mu = 0$.

(5) If $f = g$ μ -almost everywhere in X , then $f^+ = g^+$ and $f^- = g^-$ μ -almost everywhere in X . This implies that

$$\int_X f^+ d\mu = \int_X g^+ d\mu \quad \text{and} \quad \int_X f^- d\mu = \int_X g^- d\mu,$$

from which the claim follows. \square

3.6 Dominated convergence theorem

Now we are ready to state the principal convergence theorem in the theory of integration. The power of the theorem is that it applies to sign changing functions and there is no assumption on monotonicity, compare with the monotone convergence theorem and Fatou's lemma, see Theorem 3.12 and Theorem 3.18.

Theorem 3.26 (Dominated convergence theorem). Let $f_i : X \rightarrow [-\infty, \infty]$, $i = 1, 2, \dots$, be μ -measurable functions such that $f_i \rightarrow f$ μ -almost everywhere as $i \rightarrow \infty$. If there exists an integrable function g such that $|f_i| \leq g$ μ -almost everywhere for every $i = 1, 2, \dots$, then f is integrable and

$$\int_X f d\mu = \lim_{i \rightarrow \infty} \int_X f_i d\mu.$$

T H E M O R A L : The order of taking limits and integral can be switched if there exists a dominating function g with $\int_X |g| d\mu < \infty$. Observe that the same g has to do for all functions $|f_i|$. In other words, mass is preserved under dominated convergence. The integrable dominating function shuts down the loss of mass.

Remark 3.27. As the moving bump examples show, see Examples 3.11, the assumption on an integrable dominating function is necessary. Indeed, an integrable dominating function does not exist in the moving bump examples.

Proof. Consider the set

$$\begin{aligned} N = & \{x \in X : \liminf_{i \rightarrow \infty} f_i(x) \neq f(x)\} \cup \{x \in X : \limsup_{i \rightarrow \infty} f_i(x) \neq f(x)\} \\ & \cup \bigcup_{i=1}^{\infty} \{x \in X : |f_i(x)| > g(x)\} \cup \{x \in X : g(x) = \infty\}, \end{aligned}$$

where the limit $\lim_{i \rightarrow \infty} f_i(x)$ does not exist or $|f_i(x)| > g(x)$ for some i . Then $\mu(N) = 0$ and

$$|f(x)| = \lim_{i \rightarrow \infty} |f_i(x)| \leq g(x) < \infty$$

for every $x \in X \setminus N$. It follows that

$$\int_X |f| d\mu = \int_{X \setminus N} |f| d\mu \leq \int_{X \setminus N} g d\mu = \int_X g d\mu < \infty$$

and thus $f \in L^1(X; \mu)$. Since $|f_i(x)| \leq g(x)$ for every $x \in X \setminus N$, as above, we conclude that $f_i \in L^1(X; \mu)$ for every $i = 1, 2, \dots$

Let

$$g_i(x) = \begin{cases} |f_i(x) - f(x)|, & x \in X \setminus N, \\ 0, & x \in N, \end{cases}$$

and $h = |f| + g$. Then

$$\int_X h d\mu = \int_X (|f| + g) d\mu = \int_X |f| d\mu + \int_X g d\mu < \infty.$$

Thus $h \in L^1(X; \mu)$ and

$$\begin{aligned} h(x) - g_i(x) &= |f(x)| + g(x) - |f_i(x) - f(x)| \\ &\geq |f(x)| + g(x) - (|f_i(x)| + |f(x)|) \\ &= g(x) - |f_i(x)| \geq 0 \end{aligned}$$

for every $x \in X \setminus N$.

Since $g_i(x) \rightarrow 0$ for every $x \in X$ as $i \rightarrow \infty$, Fatou's lemma, see Theorem 3.18, implies

$$\begin{aligned} \int_X h d\mu &= \int_X \liminf_{i \rightarrow \infty} (h - g_i) d\mu \\ &\leq \liminf_{i \rightarrow \infty} \int_X (h - g_i) d\mu \\ &= \int_X h d\mu - \limsup_{i \rightarrow \infty} \int_X g_i d\mu. \end{aligned}$$

Since $\int_X h d\mu < \infty$, we conclude that

$$\limsup_{i \rightarrow \infty} \int_X g_i d\mu \leq 0.$$

Since $g_i \geq 0$, we have

$$0 = \lim_{i \rightarrow \infty} \int_X g_i d\mu = \lim_{i \rightarrow \infty} \int_X |f_i - f| d\mu.$$

It follows that

$$\left| \int_X f_i d\mu - \int_X f d\mu \right| = \left| \int_X (f_i - f) d\mu \right| \leq \int_X |f_i - f| d\mu \rightarrow 0$$

as $i \rightarrow \infty$. □

Remarks 3.28:

(1) The proof shows that

$$\lim_{i \rightarrow \infty} \int_X |f_i - f| d\mu = 0.$$

This is also clear from the dominated convergence theorem, since $|f_i - f| \rightarrow 0$ μ -almost everywhere and $|f_i - f| \leq |f_i| + |f| \leq 2g$. Thus

$$\lim_{i \rightarrow \infty} \int_X |f_i - f| d\mu = \int_X \lim_{i \rightarrow \infty} |f_i - f| d\mu = 0.$$

In other words, the dominated convergence theorem upgrades pointwise convergence to L^1 convergence.

(2) The result is interesting and useful already for the characteristic functions of measurable sets.

(3) Assume that $\mu(X) < \infty$ and $f_i : X \rightarrow [-\infty, \infty]$, $i = 1, 2, \dots$, are μ -measurable functions such that $f_i \rightarrow f$ μ -almost everywhere as $i \rightarrow \infty$. If there exists $M < \infty$ such that $|f_i| \leq M$ μ -almost everywhere for every $i = 1, 2, \dots$, then f is integrable and

$$\int_X f d\mu = \lim_{i \rightarrow \infty} \int_X f_i d\mu.$$

Reason. The constant function $g = M$ is integrable in X , since $\mu(X) < \infty$. ■

(4) Assume that $\mu(X) < \infty$ and $f_i : X \rightarrow [-\infty, \infty]$, $i = 1, 2, \dots$, are integrable functions on X such that $f_i \rightarrow f$ uniformly in X as $i \rightarrow \infty$. Then f is integrable and

$$\int_X f d\mu = \lim_{i \rightarrow \infty} \int_X f_i d\mu.$$

We leave this as an exercise.

(5) We deduced the dominated convergence theorem from Fatou's lemma and Fatou's lemma from the monotone convergence theorem. This can be done in other order as well.

Remark 3.29. We have the following useful results for a function $f \in L^1(X; \mu)$. Compare these properties to the corresponding properties for nonnegative measurable functions.

(1) (Countable additivity) If A_i , $i = 1, 2, \dots$, are pairwise disjoint μ -measurable sets, then

$$\int_{\bigcup_{i=1}^{\infty} A_i} f d\mu = \sum_{i=1}^{\infty} \int_{A_i} f d\mu.$$

Reason. Let $s_n = \sum_{i=1}^n f \chi_{A_i}$, $n = 1, 2, \dots$, and denote $A = \bigcup_{i=1}^{\infty} A_i$. Then $s_n \rightarrow f \chi_A$ everywhere in X as $n \rightarrow \infty$. By the triangle inequality

$$|s_n| = \left| \sum_{i=1}^n f \chi_{A_i} \right| \leq \sum_{i=1}^n |f| \chi_{A_i} \leq \sum_{i=1}^{\infty} |f| \chi_{A_i} = |f| \chi_{\bigcup_{i=1}^{\infty} A_i} \leq |f|$$

for every $n = 1, 2, \dots$, where $f \in L^1(X; \mu)$. By the dominated convergence theorem

$$\begin{aligned} \int_{\bigcup_{i=1}^{\infty} A_i} f d\mu &= \int_X f \chi_A d\mu = \int_X \lim_{n \rightarrow \infty} s_n d\mu \\ &= \lim_{n \rightarrow \infty} \int_X s_n d\mu = \lim_{n \rightarrow \infty} \sum_{i=1}^n \int_{A_i} f d\mu = \sum_{i=1}^{\infty} \int_{A_i} f d\mu. \end{aligned}$$

The last equality follows from the fact that the series $\sum_{i=1}^{\infty} \int_{A_i} f d\mu$ converges absolutely, from which it follows that it converges. To conclude this, we note that

$$\sum_{i=1}^n \left| \int_{A_i} f d\mu \right| \leq \sum_{i=1}^n \int_{A_i} |f| d\mu \leq \int_A |f| d\mu < \infty$$

for every $n = 1, 2, \dots$ and consequently

$$\sum_{i=1}^{\infty} \left| \int_{A_i} f d\mu \right| = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left| \int_{A_i} f d\mu \right| < \infty. \quad \blacksquare$$

(2) (Downwards monotone convergence) If A_i is μ -measurable, $A_i \supset A_{i+1}$, $i = 1, 2, \dots$, then

$$\int_{\bigcap_{i=1}^{\infty} A_i} f d\mu = \lim_{i \rightarrow \infty} \int_{A_i} f d\mu.$$

Reason. Let $f_i = f \chi_{A_i}$ and denote $A = \bigcap_{i=1}^{\infty} A_i$. Then $|f_i| \leq |f|$, for every $i = 1, 2, \dots$, $f \in L^1(X; \mu)$ and $f_i \rightarrow f \chi_A$ everywhere in X as $i \rightarrow \infty$. By the dominated convergence theorem

$$\begin{aligned} \int_{\bigcap_{i=1}^{\infty} A_i} f d\mu &= \int_X f \chi_A d\mu = \int_X \lim_{i \rightarrow \infty} f_i d\mu \\ &= \lim_{i \rightarrow \infty} \int_X f_i d\mu = \lim_{i \rightarrow \infty} \int_{A_i} f d\mu. \end{aligned} \quad \blacksquare$$

(3) (Upwards monotone convergence) If $A_i, i = 1, 2, \dots$, are μ -measurable and $A_i \subset A_{i+1}, i = 1, 2, \dots$, then

$$\int_{\bigcup_{i=1}^{\infty} A_i} f d\mu = \lim_{i \rightarrow \infty} \int_{A_i} f d\mu.$$

Reason. Let $f_i = f \chi_{A_i}$ and denote $A = \bigcup_{i=1}^{\infty} A_i$. Then $|f_i| \leq |f|$, for every $i = 1, 2, \dots$, $f \in L^1(X)$ and $f_i \rightarrow f \chi_A$ everywhere in X as $i \rightarrow \infty$. By the dominated convergence theorem

$$\begin{aligned} \int_{\bigcup_{i=1}^{\infty} A_i} f d\mu &= \int_X f \chi_A d\mu = \int_X \lim_{i \rightarrow \infty} f_i d\mu \\ &= \lim_{i \rightarrow \infty} \int_X f_i d\mu = \lim_{i \rightarrow \infty} \int_{A_i} f d\mu. \end{aligned} \quad \blacksquare$$

(4) (Infinite series) Let $f_i \in L^1(X; \mu)$, $i = 1, 2, \dots$, be such that $\sum_{i=1}^{\infty} \|f_i\|_{L^1(X; \mu)} < \infty$. Then

$$\int_X \sum_{i=1}^{\infty} f_i d\mu = \sum_{i=1}^{\infty} \int_X f_i d\mu.$$

Reason. By Corollary 3.16 we have

$$\int_X \sum_{i=1}^{\infty} |f_i| d\mu = \sum_{i=1}^{\infty} \int_X |f_i| d\mu < \infty.$$

Lemma 3.9 implies that $\sum_{i=1}^{\infty} |f_i| < \infty$ μ -almost everywhere in X . This shows that the series $\sum_{i=1}^{\infty} f_i$ converges absolutely μ -almost everywhere in X and thus it converges μ -almost everywhere in X . Theorem 2.23 implies that

$$f = \sum_{i=1}^{\infty} f_i = \lim_{n \rightarrow \infty} \sum_{i=1}^n f_i$$

is μ -measurable. Let $s_n = \sum_{i=1}^n f_i$, $n = 1, 2, \dots$. Then

$$|s_n| = \left| \sum_{i=1}^n f_i \right| \leq \sum_{i=1}^n |f_i| \leq \sum_{i=1}^{\infty} |f_i|,$$

μ -almost everywhere in X for every $n = 1, 2, \dots$. Let $g = \sum_{i=1}^{\infty} |f_i|$. Then $g \in L^1(X; \mu)$ and $|s_n| \leq g$, $n = 1, 2, \dots$, μ -almost everywhere in X . It follows that

$$|f| = \lim_{n \rightarrow \infty} |s_n| \leq g$$

μ -almost everywhere in X and thus $f \in L^1(X; \mu)$. By the dominated convergence theorem we have

$$\begin{aligned} \int_X \sum_{i=1}^{\infty} f_i d\mu &= \int_X \lim_{n \rightarrow \infty} \sum_{i=1}^n f_i d\mu = \lim_{n \rightarrow \infty} \int_X \sum_{i=1}^n f_i d\mu \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \int_X f_i d\mu = \sum_{i=1}^{\infty} \int_X f_i d\mu. \end{aligned} \quad \blacksquare$$

The dominated convergence theorem can be applied to compute limits of certain integrals.

Examples 3.30:

(1) Consider

$$\lim_{i \rightarrow \infty} \int_0^1 ix^{-\frac{3}{2}} \sin\left(\frac{x}{i}\right) dx.$$

Let

$$f_i(x) = ix^{-\frac{3}{2}} \sin\left(\frac{x}{i}\right) = \frac{i}{x} \sin\left(\frac{x}{i}\right) x^{-\frac{1}{2}}$$

for every $0 \leq x \leq 1$ and $i = 1, 2, \dots$. Since $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$, we have

$$\lim_{i \rightarrow \infty} \frac{i}{x} \sin\left(\frac{x}{i}\right) = 1,$$

and thus $\lim_{i \rightarrow \infty} f_i(x) = x^{-\frac{1}{2}}$ for every $0 \leq x \leq 1$. Since $|\sin t| \leq t$ for every $t \geq 0$, we have

$$\left| \frac{i}{x} \sin\left(\frac{x}{i}\right) \right| \leq \left| \frac{i}{x} \cdot \frac{x}{i} \right| \leq 1$$

and thus

$$|f_i(x)| = \left| \frac{i}{x} \sin\left(\frac{x}{i}\right) x^{-\frac{1}{2}} \right| \leq x^{-\frac{1}{2}}$$

for every $i = 1, 2, \dots$ and every $0 \leq x \leq 1$. Before we can use the dominated convergence theorem, we need to show that the function $g : [0, 1] \rightarrow \mathbb{R}$, $g(x) = x^{-\frac{1}{2}}$ is integrable, but this is clear since

$$\int_0^1 g(x) dx = 2 < \infty.$$

The dominated convergence theorem implies

$$\begin{aligned} \lim_{i \rightarrow \infty} \int_0^1 ix^{-\frac{3}{2}} \sin\left(\frac{x}{i}\right) dx &= \lim_{i \rightarrow \infty} \int_0^1 f_i(x) dx \\ &= \int_0^1 \lim_{i \rightarrow \infty} f_i(x) dx = \int_0^1 g(x) dx = 2. \end{aligned}$$

(2) Assume $f \in L^1(\mathbb{R}^n)$ with respect to the Lebesgue measure. Then

$$\lim_{i \rightarrow \infty} \int_{\mathbb{R}^n} f(x) e^{-\frac{|x|^2}{i}} dx = \int_{\mathbb{R}^n} f(x) dx.$$

Reason. Since

$$\left| f(x) e^{-\frac{|x|^2}{i}} \right| \leq |f(x)|$$

for every $x \in \mathbb{R}^n$ and $i = 1, 2, \dots$, the function $|f|$ will do as an integrable majorant in the dominated convergence theorem. Thus

$$\lim_{i \rightarrow \infty} \int_{\mathbb{R}^n} f(x) e^{-\frac{|x|^2}{i}} dx = \int_{\mathbb{R}^n} f(x) \underbrace{\lim_{i \rightarrow \infty} e^{-\frac{|x|^2}{i}}}_{=1} dx = \int_{\mathbb{R}^n} f(x) dx.$$

■

We conclude this section with two useful results, which are related to integrals depending on a parameter. Assume that μ is a measure on X and let $I \subset \mathbb{R}$ be an interval. Let $f : X \times I \rightarrow [-\infty, \infty]$, $f = f(x, t)$, be such that for every $t \in I$ the function $x \mapsto f(x, t)$ is integrable. For each $t \in I$, we consider the integral of the function over X and denote

$$F(t) = \int_X f(x, t) d\mu(x).$$

We are interested in the regularity of F . First we discuss continuity.

Theorem 3.31 (Continuity). Assume that

- (1) for every $t \in I$, the function $x \mapsto f(x, t)$ is integrable in X ,
- (2) the function $t \mapsto f(x, t)$ is continuous for every $x \in X$ at $t_0 \in I$ and
- (3) there exists $g \in L^1(X; \mu)$ such that $|f(x, t)| \leq g(x)$ for every $(x, t) \in X \times I$.

Then F is continuous at t_0 .

Proof. This is a direct consequence of the dominated convergence theorem, since g will do as an integrable majorant and

$$\begin{aligned} \lim_{t \rightarrow t_0} F(t) &= \lim_{t \rightarrow t_0} \int_X f(x, t) d\mu(x) \\ &= \int_X \lim_{t \rightarrow t_0} f(x, t) d\mu(x) \\ &= \int_X f(x, t_0) d\mu(x) = F(t_0). \end{aligned} \quad \square$$

T H E M O R A L : Under these assumptions we can take limit under the integral sign. In other words, we can switch the order of taking limit and integral.

Then we discuss differentiability.

Theorem 3.32 (Differentiability). Assume that

- (1) for every $t \in I$, the function $x \mapsto f(x, t)$ is integrable in X ,
- (2) the function $t \mapsto f(x, t)$ is differentiable for every $x \in X$ at every point $t \in I$ and
- (3) there exists $h \in L^1(X; \mu)$ such that $\left| \frac{\partial f}{\partial t}(x, t) \right| \leq h(x)$ for every $(x, t) \in X \times I$.

Then F is differentiable at every point $t \in I$ and

$$F'(t) = \frac{\partial}{\partial t} \left(\int_X f(x, t) d\mu(x) \right) = \int_X \frac{\partial}{\partial t} f(x, t) d\mu(x).$$

T H E M O R A L : Under these assumptions we can differentiate under the integral sign. In other words, we can switch the order of taking derivative and integral.

Proof. Let $t \in I$ be fixed. For $|h|$ small consider the difference quotient

$$\frac{F(t+h) - F(t)}{h} = \int_X \frac{f(x, t+h) - f(x, t)}{h} d\mu(x)$$

Since f is differentiable, we have

$$\lim_{h \rightarrow 0} \frac{f(x, t+h) - f(x, t)}{h} = \frac{\partial}{\partial t} f(x, t).$$

By the mean value theorem of differential calculus

$$\left| \frac{f(x, t+h) - f(x, t)}{h} \right| = \left| \frac{\partial}{\partial t} f(x, t') \right| \leq h(x)$$

for some $t' \in (t, t+h)$. Thus by the dominated convergence theorem

$$\begin{aligned} F'(t) &= \lim_{h \rightarrow 0} \frac{F(t+h) - F(t)}{h} \\ &= \lim_{h \rightarrow 0} \int_X \frac{f(x, t+h) - f(x, t)}{h} d\mu(x) \\ &= \int_X \lim_{h \rightarrow 0} \frac{f(x, t+h) - f(x, t)}{h} d\mu(x) \\ &= \int_X \frac{\partial}{\partial t} f(x, t) d\mu(x). \end{aligned} \quad \square$$

This kind of arguments are frequently used for the Lebesgue integral and partial derivatives in real analysis.

3.7 Lebesgue integral

Lebesgue integrable functions

Let $f : \mathbb{R}^n \rightarrow [-\infty, \infty]$ be a Lebesgue measurable function. The Lebesgue integral of f is denoted as

$$\int_{\mathbb{R}^n} f dm = \int_{\mathbb{R}^n} f(x) dm(x) = \int_{\mathbb{R}^n} f(x) dx = \int_{\mathbb{R}^n} f dx,$$

whenever the integral is defined. For a Lebesgue measurable subset A of \mathbb{R}^n , we define

$$\int_A f dx = \int_{\mathbb{R}^n} f \chi_A dx.$$

Example 3.33. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $f(x) = |x|^{-\alpha}$, $\alpha > 0$. The function becomes unbounded in any neighbourhood of the origin. The function f is not defined at the origin, but we may set $f(0) = 0$.

Let $A = B(0, 1)$ and $A_i = B(0, 2^{-i+1}) \setminus B(0, 2^{-i})$, $i = 1, 2, \dots$. The sets A_i are Lebesgue measurable, pairwise disjoint and $B(0, 1) \setminus \{0\} = \bigcup_{i=1}^{\infty} A_i$. By Corollary




Figure 3.1: An exhaustion of $B(0,1)$ by annuli.

3.16, we have

$$\begin{aligned}
 \int_{B(0,1)} \frac{1}{|x|^\alpha} dx &= \int_{\mathbb{R}^n} \frac{1}{|x|^\alpha} \chi_{\bigcup_{i=1}^{\infty} A_i}(x) dx = \int_{\mathbb{R}^n} \frac{1}{|x|^\alpha} \sum_{i=1}^{\infty} \chi_{A_i}(x) dx \\
 &= \sum_{i=1}^{\infty} \int_{\mathbb{R}^n} \frac{1}{|x|^\alpha} \chi_{A_i}(x) dx = \sum_{i=1}^{\infty} \int_{A_i} \frac{1}{|x|^\alpha} dx \\
 &\leq \sum_{i=1}^{\infty} \int_{A_i} 2^{i\alpha} dx = \sum_{i=1}^{\infty} 2^{i\alpha} m(A_i) \\
 &\leq \sum_{i=1}^{\infty} 2^{i\alpha} m(B(0, 2^{-i+1})) = \sum_{i=1}^{\infty} 2^{i\alpha} 2^{n(-i+1)} m(B(0, 1)) \\
 &= 2^n m(B(0, 1)) \sum_{i=1}^{\infty} 2^{i\alpha - in} < \infty, \quad \alpha < n.
 \end{aligned}$$

On the other hand,

$$\begin{aligned}
 \int_{B(0,1)} \frac{1}{|x|^\alpha} dx &= \sum_{i=1}^{\infty} \int_{A_i} \frac{1}{|x|^\alpha} dx \geq \sum_{i=1}^{\infty} \int_{A_i} 2^{(i-1)\alpha} dx = \sum_{i=1}^{\infty} 2^{(i-1)\alpha} m(A_i) \\
 &= (2^n - 1) 2^{-\alpha} m(B(0, 1)) \sum_{i=1}^{\infty} 2^{i\alpha - in} = \infty, \quad \alpha \geq n.
 \end{aligned}$$

The last equality follows from

$$\begin{aligned}
 m(A_i) &= m(B(0, 2^{-i+1})) - m(B(0, 2^{-i})) \\
 &= (2^{(-i+1)n} - 2^{-in}) m(B(0, 1)) \\
 &= 2^{-in} (2^n - 1) m(B(0, 1)).
 \end{aligned}$$

Thus

$$\int_{B(0,1)} \frac{1}{|x|^\alpha} dx < \infty \iff \alpha < n.$$

A similar reasoning with the sets $A_i = B(0, 2^i) \setminus B(0, 2^{i-1})$, $i = 1, 2, \dots$, shows that

$$\int_{\mathbb{R}^n \setminus B(0,1)} \frac{1}{|x|^\alpha} dx < \infty \iff \alpha > n.$$




Figure 3.2: An exhaustion of $\mathbb{R}^n \setminus B(0,1)$ by annuli.

We shall show in Example 3.38 how to compute these integrals by a change of variables and spherical coordinates.

Example 3.34. Let (q_i) be a countable and dense subset of \mathbb{R}^n . For example, we may consider \mathbb{Q}^n which is the set of points in \mathbb{R}^n with rational coordinates. Let $\phi : \mathbb{R}^n \rightarrow [0, \infty]$,

$$\phi(x) = |x|^{-\alpha} \chi_{B(0,1)}(x) = \begin{cases} |x|^{-\alpha} & |x| < 1, \\ 0, & |x| \geq 1, \end{cases}$$

with $0 < \alpha < n$. Let $f : \mathbb{R}^n \rightarrow [0, \infty]$,

$$f(x) = \sum_{i=1}^{\infty} \frac{1}{2^i} \phi(x - q_i).$$

By Corollary 3.16, translation invariance of Lebesgue integral (discussed later)

and Example 3.33, we have

$$\begin{aligned}
 \int_{\mathbb{R}^n} f(x) dx &= \int_{\mathbb{R}^n} \sum_{i=1}^{\infty} \frac{1}{2^i} \phi(x - q_i) dx \\
 &= \sum_{i=1}^{\infty} \frac{1}{2^i} \int_{\mathbb{R}^n} \phi(x - q_i) dx \\
 &= \sum_{i=1}^{\infty} \frac{1}{2^i} \int_{\mathbb{R}^n} \phi(x) dx \\
 &= \int_{B(0,1)} \phi(x) dx < \infty.
 \end{aligned}$$

Thus $f \in L^1(\mathbb{R}^n)$. Note that f has a singularity at every q_i , $i = 1, 2, \dots$, that is,

$$\lim_{x \rightarrow q_i} f(x) = \infty \quad \text{for every } i = 1, 2, \dots$$

However, since f is integrable we have $f(x) < \infty$ for almost every $x \in \mathbb{R}^n$. In other words, the series

$$f(x) = \sum_{i=1}^{\infty} \frac{1}{2^i} \phi(x - q_i)$$

converges for almost every $x \in \mathbb{R}^n$. This function is discontinuous at every point and it cannot be redefined on a set of measure zero so that it becomes continuous, compare to Lusin's theorem 2.53. Moreover, the function is unbounded in any neighborhood of any point.

L^1 space

Let $f : \mathbb{R}^n \rightarrow [-\infty, \infty]$ be an integrable function in \mathbb{R}^n with respect to the Lebesgue measure, denoted by $f \in L^1(\mathbb{R}^n)$. The number

$$\|f\|_{L^1(\mathbb{R}^n)} = \int_{\mathbb{R}^n} |f| dx < \infty.$$

is called the L^1 -norm of f . This has the usual properties

- (1) $0 \leq \|f\|_{L^1(\mathbb{R}^n)} < \infty$,
- (2) $\|f\|_{L^1(\mathbb{R}^n)} = 0 \iff f = 0$ almost everywhere,
- (3) $\|af\|_{L^1(\mathbb{R}^n)} = |a| \|f\|_{L^1(\mathbb{R}^n)}$, $a \in \mathbb{R}$, and
- (4) $\|f + g\|_{L^1(\mathbb{R}^n)} \leq \|f\|_{L^1(\mathbb{R}^n)} + \|g\|_{L^1(\mathbb{R}^n)}$.

The last triangle inequality in L^1 follows from the pointwise triangle inequality $|f(x) + g(x)| \leq |f(x)| + |g(x)|$. However, there are slight problems with the vector space properties of $L^1(\mathbb{R}^n)$, since the sum function $f + g$ may be $\infty - \infty$ and is not necessarily defined at every point. However, by Lemma 3.9 (2) integrable functions are finite almost everywhere and this is not a serious problem. Moreover, $\|f\|_{L^1(\mathbb{R}^n)} = 0$ implies that $f = 0$ almost everywhere, but not necessarily everywhere,

see Lemma 3.9 (1). We can overcome this problem by considering equivalence classes of functions that coincide almost everywhere.

We also recall the following useful properties which also hold for more general measures. Let $f \in L^1(\mathbb{R}^n)$. Then the following claims are true:

(1) (Finiteness) If $f \in L^1(\mathbb{R}^n)$, then $|f| < \infty$ almost everywhere in \mathbb{R}^n . The converse does not hold as the example above shows.

(2) (Vanishing) If $\int_{\mathbb{R}^n} |f| dx = 0$, then $f = 0$ almost everywhere in \mathbb{R}^n .

(3) (Horizontal truncation) Approximation by integrals over bounded sets

$$\int_{\mathbb{R}^n} |f| dx = \int_{\mathbb{R}^n} \lim_{i \rightarrow \infty} \chi_{B(0,i)} |f| dx = \lim_{i \rightarrow \infty} \int_{B(0,i)} |f| dx.$$

Here we used the monotone convergence theorem or the dominated convergence theorem if $f \in L^1(\mathbb{R}^n)$.

(4) (Vertical truncation) Approximation by integrals of bounded functions

$$\int_{\mathbb{R}^n} |f| dx = \int_{\mathbb{R}^n} \lim_{i \rightarrow \infty} \min\{|f|, i\} dx = \lim_{i \rightarrow \infty} \int_{\mathbb{R}^n} \min\{|f|, i\} dx.$$

Here we again used the monotone convergence theorem or the dominated convergence theorem if $f \in L^1(\mathbb{R}^n)$.

L^1 convergence

We say that $f_i \rightarrow f$ in $L^1(\mathbb{R}^n)$, if

$$\|f_i - f\|_{L^1(\mathbb{R}^n)} \xrightarrow{i \rightarrow \infty} 0.$$

This is yet another mode of convergence.

Remarks 3.35:

(1) If $f_i \rightarrow f$ in $L^1(\mathbb{R}^n)$, then $f_i \rightarrow f$ in measure.

Reason. By Tchebyschev's inequality (Lemma 3.8 (5))

$$m(\{x \in \mathbb{R}^n : |f_i(x) - f(x)| > \varepsilon\}) \leq \frac{1}{\varepsilon} \int_{\mathbb{R}^n} |f_i(x) - f(x)| dx \xrightarrow{i \rightarrow \infty} 0$$

for every $\varepsilon > 0$. ■

Theorem 2.47 implies that if $f_i \rightarrow f$ in $L^1(\mathbb{R}^n)$, then there exists a subsequence such that $f_{i_k} \rightarrow f$ μ -almost everywhere. An example of a sliding sequence of functions, see Example 2.46 (4), shows that the claim is not true for the original sequence.

(2) The Riesz-Fischer theorem states that L^1 is a Banach space, that is, every Cauchy sequence converges. We shall prove this result in the real analysis course.

Invariance properties

The invariance properties of the Lebesgue measure in Section 1.8 imply the following results:

(1) (Translation invariance)

$$\int_{\mathbb{R}^n} f(x + x_0) dx = \int_{\mathbb{R}^n} f(x) dx$$

for any $x_0 \in \mathbb{R}^n$. This means that the Lebesgue integral is invariant in translations.

Reason. We shall check this first with $f = \chi_A$, where A is Lebesgue measurable. Then $\chi_A(x + x_0) = \chi_{A - x_0}(x)$ and the claim follows from

$$\begin{aligned} \int_{\mathbb{R}^n} f(x + x_0) dx &= \int_{\mathbb{R}^n} \chi_A(x + x_0) dx = \int_{\mathbb{R}^n} \chi_{A - x_0}(x) dx \\ &= m(A - x_0) = m(A) \\ &= \int_{\mathbb{R}^n} \chi_A(x) dx = \int_{\mathbb{R}^n} f(x) dx. \end{aligned}$$

By linearity, the result holds for nonnegative simple functions. For nonnegative Lebesgue measurable functions the claim follows from the monotone convergence theorem by approximating with an increasing sequence of simple functions, see Theorem 2.33 and Theorem 3.12. The general case follows from this. \blacksquare

(2) (Reflection invariance)

$$\int_{\mathbb{R}^n} f(x) dx = \int_{\mathbb{R}^n} f(-x) dx.$$

(3) (Scaling property)

$$\int_{\mathbb{R}^n} f(x) dx = |\delta|^n \int_{\mathbb{R}^n} f(\delta x) dx$$

for any $\delta \neq 0$. This shows that the Lebesgue integral behaves as expected in dilations (exercise).

(4) (Linear change of variables) Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a general invertible linear mapping. Then

$$\int_{\mathbb{R}^n} f(Lx) dx = \frac{1}{|\det L|} \int_{\mathbb{R}^n} f(x) dx,$$

or equivalently,

$$\int_{\mathbb{R}^n} f(L^{-1}x) dx = |\det L| \int_{\mathbb{R}^n} f(x) dx.$$

Reason. Let $A \subset \mathbb{R}^n$ be a Lebesgue measurable set and $f = \chi_A$. Then $\chi_A \circ L = \chi_{L^{-1}(A)}$ is a Lebesgue measurable function and

$$\begin{aligned} \int_{\mathbb{R}^n} f(Lx) dx &= \int_{\mathbb{R}^n} \chi_A(Lx) dx = \int_{\mathbb{R}^n} (\chi_A \circ L)(x) dx = \int_{\mathbb{R}^n} \chi_{L^{-1}(A)}(x) dx \\ &= m(L^{-1}(A)) = |\det L^{-1}| m(A) \\ &= \frac{1}{|\det L|} \int_{\mathbb{R}^n} \chi_A(x) dx = \frac{1}{|\det L|} \int_{\mathbb{R}^n} f(x) dx. \end{aligned}$$

By taking linear combinations, we conclude the result for simple functions and the general case follows from the fact that a measurable function can be approximated by simple functions and the definition of the integral, see [6] pages 170–171 and 65–80 or [16] pages 612–619. ■

This is a change of variables formula for linear mappings, which is compatible with the corresponding property $m(L(A)) = |\det L| m(A)$ of the Lebesgue measure, see Section 1.8.

Reason.

$$\begin{aligned} m(L(A)) &= \int_{L(A)} 1 dx = \int_{\mathbb{R}^n} \chi_{L(A)}(x) dx \\ &= \int_{\mathbb{R}^n} (\chi_A \circ L^{-1})(x) dx = \int_{\mathbb{R}^n} \chi_A(L^{-1}x) dx \\ &= |\det L| \int_{\mathbb{R}^n} \chi_A(x) dx = |\det L| m(A). \end{aligned}$$

Moreover, A is a Borel set if and only if $L(A)$ is a Borel set, since L is a homeomorphism. ■

(5) (Nonlinear change of variables) Let $U \subset \mathbb{R}^n$ be an open set and suppose that $\Phi : U \rightarrow \mathbb{R}^n$, $\Phi = (\phi_1, \dots, \phi_n)$ is a C^1 diffeomorphism. We denote by $D\Phi$ the derivative matrix with entries $D_j \phi_i$, $i, j = 1, \dots, n$. The mapping Φ is called C^1 diffeomorphism if it is injective and $D\Phi(x)$ is invertible at every $x \in U$. In this case the inverse function theorem guarantees that the inverse map $\Phi^{-1} : \Phi(U) \rightarrow U$ is also a C^1 diffeomorphism. This means that all component functions ϕ_i , $i = 1, \dots, n$ have continuous first order partial derivatives and

$$D\Phi^{-1}(y) = (D\Phi(\Phi^{-1}(y)))^{-1}$$

for every $y \in \Phi(U)$. If f is a Lebesgue measurable function on $\Phi(U)$, then $f \circ \Phi$ is a Lebesgue measurable function on U . If f is nonnegative or integrable on $\Phi(U)$, then

$$\int_{\Phi(U)} f(y) dy = \int_U f(\Phi(x)) |\det D\Phi(x)| dx.$$

Moreover, if $A \subset U$ is a Lebesgue measurable set, then $\Phi(A)$ is a Lebesgue measurable set and

$$m(\Phi(A)) = \int_A |\det D\Phi| dx.$$




Figure 3.3: A linear change of variables.

This is a change of variables formula for differentiable mappings, see [6, p. 494–503] or [16, p. 649–660]. See also [4] Chapter 3. Formally it can be seen as the substitution $y = \Phi(x)$. This means that we replace $f(y)$ by $f(\Phi(x))$, $\Phi(U)$ by U and dy by $|\det D\Phi(x)| dx$. Observe, that if Φ is a linear mapping, that is there exists a matrix A with $\Phi(x) = Ax$, then $D\Phi = A$, and this is compatible with the change of variables formula for linear mappings.

Example 3.36. Probably the most important nonlinear coordinate systems in \mathbb{R}^2 are the polar coordinates $(x_1 = r \cos \theta_1, x_2 = r \sin \theta_1)$ and in \mathbb{R}^3 are the spherical coordinates $(x_1 = r \cos \theta_1, x_2 = r \sin \theta_1 \cos \theta_2, x_3 = r \sin \theta_1 \sin \theta_2)$. Let us consider the spherical coordinates in \mathbb{R}^n . Let

$$U = (0, \infty) \times (0, \pi)^{n-2} \times (0, 2\pi) \subset \mathbb{R}^n, \quad n \geq 2.$$

Denote the coordinates of a point in U by $r, \theta_1, \dots, \theta_{n-2}, \theta_{n-1}$, respectively. We define $\Phi : U \rightarrow \mathbb{R}^n$ by the spherical coordinate formulas as follows. If $x = \Phi(r, \theta)$, then

$$x_i = r \sin \theta_1 \cdots \sin \theta_{i-1} \cos \theta_i, \quad i = 1, \dots, n,$$

where $\theta_n = 0$ so that $x_n = r \sin \theta_1 \cdots \sin \theta_{n-1}$. Then ϕ is a bijection from U onto the




Figure 3.4: A diffeomorphic change of variables.

open set $\mathbb{R}^n \setminus (\mathbb{R}^{n-2} \times [0, \infty) \times \{0\})$. The change of variables formula implies that

$$\begin{aligned} & \int_{\mathbb{R}^n} f(x) dx \\ &= \int_0^r \int_0^\pi \dots \int_0^\pi \int_0^{2\pi} f(\Phi(r, \theta)) r^{n-1} (\sin \theta_1)^{n-2} \dots (\sin \theta_{n-3})^2 \sin \theta_{n-2} d\theta_{n-1} \dots d\theta_1 dr. \end{aligned}$$

It can be shown that

$$\omega_{n-1} = \int_0^\pi (\sin \theta_1)^{n-2} d\theta_{n-1} \dots \int_0^\pi (\sin \theta_{n-3})^2 d\theta_{n-3} \int_0^\pi \sin \theta_{n-2} d\theta_{n-2} \int_0^{2\pi} d\theta_{n-1},$$

where

$$\omega_{n-1} = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})}$$

is the $(n-1)$ -dimensional volume of the unit sphere $\partial B(0, 1) = \{x \in \mathbb{R}^n : |x| = 1\}$.

Here

$$\Gamma(a) = \int_0^\infty x^{a-1} e^{-x} dx, \quad 0 < a < \infty,$$

is the gamma function. The gamma function has the properties $\Gamma(1) = 1$ and $\Gamma(a+1) = a\Gamma(a)$. It follows that $\Gamma(k+1) = k!$ for a nonnegative integer k .

Suppose that $f : \mathbb{R}^n \rightarrow [0, \infty]$ is radial. Thus f depends only on $|x|$ and it can be expressed as $f(|x|)$, where f is a function defined on $[0, \infty)$. Then

$$\int_{\mathbb{R}^n} f(|x|) dx = \omega_{n-1} \int_0^\infty f(r) r^{n-1} dr. \quad (3.37)$$

see [6] pages 503–504 or [16] pages 661–673.

Let us show how to use this formula to compute the volume of a ball $B(x, r) = \{y \in \mathbb{R}^n : |y - x| < r\}$, $x \in \mathbb{R}^n$ and $r > 0$. Denote $\Omega_n = m(B(0, 1))$. By the translation and scaling invariance and (3.37), we have

$$\begin{aligned} r^n \Omega_n &= r^n m(B(0, 1)) = m(B(x, r)) = m(B(0, r)) \\ &= \int_{\mathbb{R}^n} \chi_{B(0,r)}(y) dy = \int_{\mathbb{R}^n} \chi_{(0,r)}(|y|) dy \\ &= \omega_{n-1} \int_0^r \rho^{n-1} d\rho = \omega_{n-1} \frac{r^n}{n}. \end{aligned}$$

In particular, it follows that $\omega_{n-1} = n \Omega_n$ and

$$m(B(x, r)) = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \frac{r^n}{n} = \frac{\pi^{n/2}}{\Gamma(\frac{n}{2} + 1)} r^n.$$

Example 3.38. Let $r > 0$. Then by property (3) above and Example 3.33,

$$\begin{aligned} \int_{\mathbb{R}^n \setminus B(0,r)} \frac{1}{|x|^\alpha} dx &= \int_{\mathbb{R}^n} \frac{1}{|x|^\alpha} \chi_{\mathbb{R}^n \setminus B(0,r)}(x) dx \\ &= r^n \int_{\mathbb{R}^n} \frac{1}{|rx|^\alpha} \chi_{\mathbb{R}^n \setminus B(0,r)}(rx) dx \\ &= r^{n-\alpha} \int_{\mathbb{R}^n} \frac{1}{|x|^\alpha} \chi_{\mathbb{R}^n \setminus B(0,1)}(x) dx \\ &= r^{n-\alpha} \int_{\mathbb{R}^n \setminus B(0,1)} \frac{1}{|x|^\alpha} dx < \infty, \quad \alpha > n, \end{aligned}$$

and, in a similar way,

$$\int_{B(0,r)} \frac{1}{|x|^\alpha} dx = r^{n-\alpha} \int_{B(0,1)} \frac{1}{|x|^\alpha} dx < \infty, \quad \alpha < n.$$

Observe, that here we formally make the change of variables $x = ry$.

On the other hand, the integrals can be computed directly by (3.37). This gives

$$\begin{aligned} \int_{\mathbb{R}^n \setminus B(0,r)} \frac{1}{|x|^\alpha} dx &= \omega_{n-1} \int_r^\infty \rho^{-\alpha} \rho^{n-1} d\rho \\ &= \frac{\omega_{n-1}}{-\alpha + n} \rho^{-\alpha+n} \Big|_r^\infty = \frac{\omega_{n-1}}{\alpha - n} r^{-\alpha+n} < \infty, \quad \alpha > n \end{aligned}$$

and

$$\begin{aligned} \int_{B(0,r)} \frac{1}{|x|^\alpha} dx &= \omega_{n-1} \int_0^r \rho^{-\alpha} \rho^{n-1} d\rho \\ &= \frac{\omega_{n-1}}{-\alpha + n} \rho^{-\alpha+n} \Big|_0^r = \frac{\omega_{n-1}}{n - \alpha} r^{n-\alpha} < \infty, \quad \alpha < n. \end{aligned}$$

Remark 3.39. Formula (3.37) (or Example 3.33) implies following claims:

(1) If $|f(x)| \leq c|x|^{-\alpha}$ in a ball $B(0, r)$, $r > 0$, for some $\alpha < n$, then $f \in L^1(B(0, r))$.

On the other hand, if $|f(x)| \geq c|x|^{-\alpha}$ in $B(0, r)$ for some $\alpha > n$, then $f \notin L^1(B(0, r))$.

(2) If $|f(x)| \leq c|x|^{-\alpha}$ in $\mathbb{R}^n \setminus B(0, r)$ for some $\alpha > n$, then $f \in L^1(\mathbb{R}^n \setminus B(0, r))$.

On the other hand, if $|f(x)| \geq c|x|^{-\alpha}$ in $\mathbb{R}^n \setminus B(0, r)$ for some $\alpha < n$, then $f \notin L^1(\mathbb{R}^n \setminus B(0, r))$.

Approximation by continuous functions

L^1 functions have the following approximation properties.

Theorem 3.40. Let $f \in L^1(\mathbb{R}^n)$ and $\varepsilon > 0$.

(1) There is a simple function $g \in L^1(\mathbb{R}^n)$ such that $\|f - g\|_{L^1(\mathbb{R}^n)} < \varepsilon$.

(2) There is a compactly supported continuous function $g \in C_0(\mathbb{R}^n)$ such that $\|f - g\|_{L^1(\mathbb{R}^n)} < \varepsilon$.

T H E M O R A L : Simple functions and compactly supported continuous functions are dense in $L^1(\mathbb{R}^n)$.

Proof. (1) Since $f = f^+ - f^-$, we may consider f^+ and f^- separately and assume that $f \geq 0$. By Theorem 2.33 there exists an increasing sequence of simple functions f_i , $i = 1, 2, \dots$, such that $f_i \rightarrow f$ everywhere in \mathbb{R}^n as $i \rightarrow \infty$. By the dominated convergence theorem (Theorem 3.26), we have

$$\lim_{i \rightarrow \infty} \|f_i - f\|_{L^1(\mathbb{R}^n)} = \lim_{i \rightarrow \infty} \int_{\mathbb{R}^n} |f_i - f| dx = \int_{\mathbb{R}^n} \underbrace{\lim_{i \rightarrow \infty} |f_i - f|}_{=0} dx = 0,$$

because $|f_i - f| \leq |f_i| + |f| \leq 2|f| \in L^1(\mathbb{R}^n)$ for every $i = 1, 2, \dots$, gives an integrable dominating function.

(2) **Step 1:** Since $f = f^+ - f^-$ we may assume that $f \geq 0$.

Step 2: The dominated convergence theorem (Theorem 3.26) gives

$$\begin{aligned} \lim_{i \rightarrow \infty} \|f \chi_{B(0,i)} - f\|_{L^1(\mathbb{R}^n)} &= \lim_{i \rightarrow \infty} \int_{\mathbb{R}^n} |f \chi_{B(0,i)} - f| dx \\ &= \int_{\mathbb{R}^n} \underbrace{\lim_{i \rightarrow \infty} |f \chi_{B(0,i)} - f|}_{=0} dx = 0, \end{aligned}$$

since $|f \chi_{B(0,i)} - f| \leq |f| \in L^1(\mathbb{R}^n)$ for every $i = 1, 2, \dots$. Thus compactly supported integrable functions are dense in $L^1(\mathbb{R}^n)$.

Step 3: By Theorem 2.33 there exists an increasing sequence of simple functions $f_i : \mathbb{R}^n \rightarrow [0, \infty)$, $i = 1, 2, \dots$, such that $f_i \rightarrow f$ everywhere in \mathbb{R}^n as $i \rightarrow \infty$. The dominated convergence theorem gives

$$\lim_{i \rightarrow \infty} \|f_i - f\|_{L^1(\mathbb{R}^n)} = \lim_{i \rightarrow \infty} \int_{\mathbb{R}^n} |f_i - f| dx = \int_{\mathbb{R}^n} \underbrace{\lim_{i \rightarrow \infty} |f_i - f|}_{=0} dx = 0.$$

since $|f_i - f| \leq |f_i| + |f| \leq 2|f| \in L^1(\mathbb{R}^n)$ for every $i = 1, 2, \dots$. Thus we can assume that we can approximate a nonnegative simple function which vanishes outside a bounded set.

Step 4: Such a function is of the form $\sum_{i=1}^k a_i \chi_{A_i}$, where A_i are bounded Lebesgue measurable set and $a_i \geq 0$, $i = 1, 2, \dots$. Thus if we can approximate each χ_{A_i} by a compactly supported continuous function, then the corresponding linear combination will approximate the simple function.

Step 5: Let A be a bounded Lebesgue measurable set and $\varepsilon > 0$. Since $m(A) < \infty$ by Theorem 1.73 there exist a compact set K and a open set G such that $K \subset A \subset G$ and $m(G \setminus K) < \varepsilon$.

Claim: There exists a continuous function $g : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

- (1) $0 \leq g(x) \leq 1$ for every $x \in \mathbb{R}^n$,
- (2) $g(x) = 1$ for every $x \in K$ and
- (3) the support of g is a compact subset of G .

Reason. Let

$$U = \{x \in \mathbb{R}^n : \text{dist}(x, K) < \frac{1}{2} \text{dist}(K, \mathbb{R}^n \setminus G)\}.$$

Then $K \subset U \subset \overline{U} \subset G$, U is open and \overline{U} is compact. The function $g : \mathbb{R}^n \rightarrow \mathbb{R}$,

$$g(x) = \frac{\text{dist}(x, \mathbb{R}^n \setminus U)}{\text{dist}(x, K) + \text{dist}(x, \mathbb{R}^n \setminus U)},$$

has the desired properties, see Remark 1.32. Moreover,

$$\text{supp } g = \overline{\{x \in \mathbb{R}^n : g(x) \neq 0\}} = \overline{U}$$

is compact. ■

Observe that

$$\begin{aligned} x \in K &\implies |\chi_A(x) - g(x)| = 1 - 1 = 0, \\ x \in \mathbb{R}^n \setminus G &\implies |\chi_A(x) - g(x)| = 0 - 0 = 0, \\ x \in A \setminus K &\implies |\chi_A(x) - g(x)| = 1 - g(x) < 1, \\ x \in G \setminus A &\implies |\chi_A(x) - g(x)| = g(x) < 1. \end{aligned}$$

Thus $|\chi_A - g| \leq 1$, $|\chi_A - g|$ vanishes in K and outside $G \setminus K$ and we have

$$\|\chi_A - g\|_{L^1(\mathbb{R}^n)} = \int_{\mathbb{R}^n} |\chi_A - g| dx \leq m(G \setminus K) < \varepsilon.$$

This completes the proof of the approximation property. □

Remark 3.41. The claim in the step 5 can also be proved using Lusin's theorem (Theorem 2.53) (exercise).

3.8 Cavalieri's principle

Recall, that every nonnegative measurable function satisfies Tchebyshev's inequality (Theorem 3.8 (5))

$$m(\{x \in \mathbb{R}^n : f(x) > t\}) \leq \frac{1}{t} \int_{\mathbb{R}^n} f \, dx, \quad t > 0.$$

In particular, if $f \in L^1(\mathbb{R}^n)$, then

$$m(\{x \in \mathbb{R}^n : f(x) > t\}) \leq \frac{c}{t}, \quad t > 0, \quad (3.42)$$

with $c = \|f\|_{L^1(\mathbb{R}^n)} < \infty$. The converse claim is not true, that is, if f satisfies an inequality of the form (3.42), it does not follow that $f \in L^1(\mathbb{R}^n)$.

Reason. Let $f : \mathbb{R}^n \rightarrow [0, \infty]$, $f(x) = |x|^{-n}$. Then f satisfies (3.42), but $f \notin L^1(\mathbb{R}^n)$. ■

The function $t \mapsto m(\{x \in \mathbb{R}^n : f(x) > t\})$ is called the distribution function of f . Observe that the distribution set $\{x \in \mathbb{R}^n : f(x) > t\}$ is Lebesgue measurable and the distribution function is a nonincreasing function of $t > 0$ and hence Lebesgue measurable. Let us consider the behaviour of $tm(\{x \in \mathbb{R}^n : f(x) > t\})$ as t increases. By Tchebyshev's inequality

$$tm(\{x \in \mathbb{R}^n : f(x) > t\}) \leq \|f\|_{L^1(\mathbb{R}^n)}$$

for every $t > 0$ if $f \in L^1(\mathbb{R}^n)$, but there is a stronger result as $t \rightarrow \infty$.

Lemma 3.43. If $f \in L^1(\mathbb{R}^n)$ is a nonnegative function, then

$$\lim_{t \rightarrow \infty} tm(\{x \in \mathbb{R}^n : f(x) > t\}) = 0.$$

Proof. Let $A = \{x \in \mathbb{R}^n : f(x) < \infty\}$. Since $f \in L^1(\mathbb{R}^n)$, by Lemma 3.9 (2), we have $m(\mathbb{R}^n \setminus A) = 0$. Let $A_t = \{x \in \mathbb{R}^n : f(x) > t\}$, $t > 0$. Then

$$A = \bigcup_{0 < t < \infty} (\mathbb{R}^n \setminus A_t) \quad \text{and} \quad \lim_{t \rightarrow \infty} \chi_{\mathbb{R}^n \setminus A_t}(x) = \chi_A(x) \quad \text{for every } x \in \mathbb{R}^n.$$

Clearly

$$\int_{\mathbb{R}^n} f \, dx = \int_{A_t} f \, dx + \int_{\mathbb{R}^n \setminus A_t} f \, dx.$$

By the dominated convergence theorem (Theorem 3.26)

$$\begin{aligned} \lim_{t \rightarrow \infty} \int_{\mathbb{R}^n \setminus A_t} f \, dx &= \lim_{t \rightarrow \infty} \int_{\mathbb{R}^n} \chi_{\mathbb{R}^n \setminus A_t} f \, dx \\ &= \int_{\mathbb{R}^n} \lim_{t \rightarrow \infty} \chi_{\mathbb{R}^n \setminus A_t} f \, dx = \int_{\mathbb{R}^n} \chi_A f \, dx \\ &= \int_A f \, dx = \int_A f \, dx + \underbrace{\int_{\mathbb{R}^n \setminus A} f \, dx}_{=0} = \int_{\mathbb{R}^n} f \, dx. \end{aligned}$$

Thus

$$\begin{aligned} \underbrace{\int_{\mathbb{R}^n} f dx}_{<\infty} &= \lim_{t \rightarrow \infty} \left(\int_{A_t} f dx + \int_{\mathbb{R}^n \setminus A_t} f dx \right) \\ &= \lim_{t \rightarrow \infty} \int_{A_t} f dx + \underbrace{\int_{\mathbb{R}^n} f dx}_{<\infty}, \end{aligned}$$

which implies that

$$\lim_{t \rightarrow \infty} \int_{A_t} f dx = 0.$$

By Tchebyshev's inequality

$$0 \leq tm(A_t) \leq \int_{A_t} f dx \xrightarrow{t \rightarrow \infty} 0,$$

which implies $tm(A_t) \rightarrow 0$ as $t \rightarrow \infty$. \square

The following representation of the integral is very useful.

Theorem 3.44 (Cavalieri's principle). Let $A \subset \mathbb{R}^n$ be a Lebesgue measurable set and let $f : A \rightarrow [0, \infty]$ be a Lebesgue measurable function. Then

$$\int_A f dx = \int_0^\infty m(\{x \in A : f(x) > t\}) dt.$$

T H E M O R A L : In order to estimate the integral of a function it is enough to estimate the distribution sets of the function.

Remarks 3.45:

(1) For signed Lebesgue measurable functions we have

$$\int_A |f| dx = \int_0^\infty m(\{x \in A : |f(x)| > t\}) dt.$$

(2) Without Cavalieri's principle it can be shown (exercise) that

$$\int_A |f|^p dx < \infty \iff \sum_{i=-\infty}^{\infty} 2^{ip} m(\{x \in A : |f(x)| > 2^i\}) < \infty, \quad 0 < p < \infty.$$

(3) If $g : A \rightarrow [0, \infty]$ is a rearrangement of f such that

$$m(\{x \in A : g(x) > t\}) = m(\{x \in A : f(x) > t\})$$

for every $t \geq 0$, then $\int_A g dx = \int_A f dx$. Thus the integral of a nonnegative measurable function is independent under rearrangements that preserve the distribution function.

(4) Cavalieri's principle can be taken as the definition of the Lebesgue integral but then we have to be able to define the right hand side of Cavalieri's principle without using the one-dimensional Lebesgue integral. If $\mu(A) <$




Figure 3.5: Cavalieri's principle.

∞ , then the distribution function is a bounded monotone function and thus continuous almost everywhere in $[0, \infty)$. This implies that the distribution function is Riemann integrable on any compact interval in $[0, \infty)$ and thus that the right-hand side of Cavalieri's principle can be interpreted as an improper Riemann integral, see Remark 3.49.

Proof. **Step 1:** First assume that f is a nonnegative simple function that vanishes outside a bounded set. Then $f = \sum_{i=0}^k a_i \chi_{A_i}$, where $A_i = f^{-1}(\{a_i\})$. We may assume

that $0 = a_0 < a_1 < \dots < a_k$. Then

$$\begin{aligned}
 \int_0^\infty m(\{x \in A : f(x) > t\}) dt &= \int_0^{a_k} m(\{x \in A : f(x) > t\}) dt \\
 &= \sum_{i=1}^k \int_{a_{i-1}}^{a_i} m(\{x \in A : f(x) > t\}) dt \\
 &= \sum_{i=1}^k (a_i - a_{i-1}) m\left(\bigcup_{j=i}^k A_j \cap A\right) \quad \left(f = \sum_{i=0}^k a_i \chi_{A_i}\right) \\
 &= \sum_{i=1}^k (a_i - a_{i-1}) \sum_{j=i}^k m(A_j \cap A) \quad (A_j \text{ measurable and disjoint}) \\
 &= \sum_{i=1}^k a_i \sum_{j=i}^k m(A_j \cap A) - \sum_{i=1}^k a_{i-1} \sum_{j=i}^k m(A_j \cap A) \\
 &= \sum_{j=1}^k m(A_j \cap A) \sum_{i=1}^j a_i - \sum_{j=1}^k m(A_j \cap A) \sum_{i=1}^j a_{i-1} \\
 &= \sum_{j=1}^k m(A_j \cap A) \sum_{i=1}^j (a_i - a_{i-1}) = \sum_{j=1}^k a_j m(A_j \cap A) = \int_A f dx.
 \end{aligned}$$

This proves the claim for nonnegative simple functions that vanish outside a bounded set.




Figure 3.6: Cavalieri's principle for a simple function.

Step 2: Assume then that f is a nonnegative measurable function. As in the proof of Theorem 3.40 there exists a sequence of nonnegative simple functions that vanish outside a bounded set f_i , $i = 1, 2, \dots$, such that $f_i \leq f_{i+1}$ and $f_i(x) \rightarrow f(x)$ as

$i \rightarrow \infty$ for every $x \in A$. Thus

$$\{x \in A : f_i(x) > t\} \subset \{x \in A : f_{i+1}(x) > t\}$$

and

$$\bigcup_{i=1}^{\infty} \{x \in A : f_i(x) > t\} = \{x \in A : f(x) > t\}.$$

Let

$$\varphi_i(t) = m(\{x \in A : f_i(x) > t\}) \quad \text{and} \quad \varphi(t) = m(\{x \in A : f(x) > t\}).$$

Then φ_j is an increasing sequence of functions and $\varphi_i(t) \rightarrow \varphi(t)$ for every $t \geq 0$ as $i \rightarrow \infty$. The monotone convergence theorem implies

$$\begin{aligned} \int_0^{\infty} m(\{x \in A : f(x) > t\}) dt &= \lim_{i \rightarrow \infty} \int_0^{\infty} m(\{x \in A : f_i(x) > t\}) dt \\ &= \lim_{i \rightarrow \infty} \int_A f_i dx = \int_A f dx. \end{aligned} \quad \square$$

Remarks 3.46:

(1) By a change of variables, we have

$$\int_A |f|^p dx = p \int_0^{\infty} t^{p-1} m(\{x \in A : |f(x)| > t\}) dt$$

for $0 < p < \infty$.

(2) More generally, if $\varphi : [0, \infty) \rightarrow [0, \infty)$ is a nondecreasing continuously differentiable function with $\varphi(0) = 0$, then

$$\int_A \varphi \circ |f| dx = \int_0^{\infty} \varphi'(t) m(\{x \in A : |f(x)| > t\}) dt.$$

(3) These results hold not only for the Lebesgue measure, but also for other measures.

We shall give another proof for Cavalieri's principle later, see Corollary 3.63.

Example 3.47. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $f(x) = |x|^{-\alpha}$, $0 < \alpha < n$. Then

$$\begin{aligned} \int_{B(0,1)} f(x) dx &= \int_{B(0,1)} |x|^{-\alpha} dx = \int_0^{\infty} m(\{x \in B(0,1) : |x|^{-\alpha} > t\}) dt \\ &= \int_0^1 m(B(0,1)) dt + \int_1^{\infty} m(\{x \in \mathbb{R}^n : |x| < t^{-1/\alpha}\}) dt \\ &= m(B(0,1)) + \int_1^{\infty} m(B(0, t^{-1/\alpha})) dt \\ &= m(B(0,1)) + \int_1^{\infty} t^{-n/\alpha} m(B(0,1)) dt \\ &= m(B(0,1)) \left(1 + \frac{\alpha}{n-\alpha}\right). \end{aligned}$$

3.9 Lebesgue and Riemann

T H E M O R A L : The main difference between the Lebesgue and Riemann integrals is that in the definition of Riemann integral with step functions we subdivide the domain of the function but in the definition of the Lebesgue integral with simple functions we subdivide the range of the function.

We shall briefly recall the definition of the one-dimensional Riemann integral. Let I_i , $i = 1, \dots, k$, be pairwise disjoint intervals in \mathbb{R} with $\bigcup_{i=1}^k I_i = [a, b]$ with $a, b \in \mathbb{R}$ and let a_i , $i = 1, \dots, k$, be real numbers. A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be a step function, if

$$f = \sum_{i=1}^k a_i \chi_{I_i}.$$

Observe, that a step function is just a special type of a simple function. Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Recall that the lower Riemann integral is

$$\underline{\int_a^b} f(x) dx = \sup \left\{ \int_{[a,b]} g dx : g \leq f \text{ on } [a, b] \text{ and } g \text{ is a step function} \right\}$$

and the upper Riemann integral is

$$\overline{\int_a^b} f(x) dx = \inf \left\{ \int_{[a,b]} h dx : f \leq h \text{ on } [a, b] \text{ and } h \text{ is a step function} \right\}.$$

Observe that we use the definition of integral for a simple function for the integral of the step function. The function f is said to be Riemann integrable, if its lower and upper integrals coincide. The common value of lower and upper integrals is the Riemann integral of f on $[a, b]$ and it is denoted by

$$\int_a^b f(x) dx.$$

If $f : [a, b] \rightarrow \mathbb{R}$ is a bounded Lebesgue measurable function, then by the definition of the Lebesgue integral,

$$\underline{\int_a^b} f(x) dx \leq \int_{[a,b]} f(x) dx \leq \overline{\int_a^b} f(x) dx.$$

This implies that if f is Riemann integrable, then the Riemann and Lebesgue integrals coincide provided f is Lebesgue measurable.

Lemma 3.48. Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded Riemann integrable function. Then f is Lebesgue integrable and

$$\int_a^b f(x) dx = \int_{[a,b]} f(x) dx.$$

T H E M O R A L : The Lebesgue integral is an extension of the Riemann integral.

Proof. By adding a constant we may assume that $f \geq 0$. By the definition of the Riemann integral, there are step functions g_i and h_i such that $g_i \leq f \leq h_i$,

$$\lim_{i \rightarrow \infty} \int_{[a,b]} g_i dx = \lim_{i \rightarrow \infty} \int_{[a,b]} h_i dx = \int_a^b f(x) dx.$$

By passing to sequences $\max\{g_1, \dots, g_j\}$ and $\min\{h_1, \dots, h_j\}$ we may assume that $g_i \leq g_{i+1}$ and $h_i \geq h_{i+1}$ for every $i = 1, 2, \dots$. These sequences are monotone and bounded and thus they converge pointwise. Denote

$$g(x) = \lim_{i \rightarrow \infty} g_i(x) \quad \text{and} \quad h(x) = \lim_{i \rightarrow \infty} h_i(x).$$

By the dominated convergence theorem

$$\int_a^b f(x) dx = \lim_{i \rightarrow \infty} \int_{[a,b]} g_i(x) dx = \int_{[a,b]} g(x) dx$$

and

$$\int_a^b f(x) dx = \lim_{i \rightarrow \infty} \int_{[a,b]} h_i(x) dx = \int_{[a,b]} h(x) dx.$$

Since $h - g \geq 0$ and

$$\begin{aligned} \int_{[a,b]} (h(x) - g(x)) dx &= \int_{[a,b]} h(x) dx - \int_{[a,b]} g(x) dx \\ &= \int_a^b f(x) dx - \int_a^b f(x) dx = 0, \end{aligned}$$

we have $h - g = 0$ almost everywhere in $[a, b]$. Since $g \leq f \leq h$ we have $h = g = f$ almost everywhere in $[a, b]$. Thus f is measurable and since it is also bounded it is integrable in $[a, b]$. \square

Remark 3.49. A necessary and sufficient condition for a function f to be Riemann integrable on an interval $[a, b]$ is that f is bounded and that its set of points of discontinuity in $[a, b]$ forms a set of Lebesgue measure zero, see [6] page 163.

Note that the definition of the Riemann integral only applies to bounded functions defined on bounded intervals. It is possible to relax these assumptions, but this becomes delicate. The definition of the Lebesgue integral applies directly to not necessarily bounded functions and sets. Note that the Lebesgue integral is defined not only over intervals but also over more general measurable sets. This is a very useful property. Moreover, the Lebesgue integral behaves better under limits compared to the Riemann integral. The following examples show differences between the Lebesgue and Riemann integrals.

Examples 3.50:

(1) $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \chi_{[0,1] \cap \mathbb{Q}}(x)$ is not Riemann integrable, but is a simple function for Lebesgue integral and

$$\int_{\mathbb{R}} \chi_{[0,1] \cap \mathbb{Q}}(x) dx = 0.$$

(2) Let q_i , $i = 1, 2, \dots$, be an enumeration of rational numbers in the interval $[0, 1]$ and define $f_i : \mathbb{R} \rightarrow \mathbb{R}$,

$$f_i(x) = \chi_{\{q_1, \dots, q_i\}}(x), \quad i = 1, 2, \dots$$

Then each f_i is Riemann integrable with zero integral, but the limit function

$$f(x) = \lim_{i \rightarrow \infty} f_i(x) = \chi_{[0,1] \cap \mathbb{Q}}(x)$$

is not Riemann integrable. This means that a pointwise limit of Riemann integrable functions may be not Riemann integrable.

(3) Define $f : [0, 1] \rightarrow \mathbb{R}$ by setting $f(0) = 0$ and

$$f(x) = \begin{cases} \frac{2^{i+1}}{i}, & \frac{1}{2^i} < x \leq \frac{3}{2^{i+1}}, \\ -\frac{2^{i+1}}{i}, & \frac{3}{2^{i+1}} < x \leq \frac{1}{2^{i-1}}, \end{cases}$$

for $x \in (0, 1]$. Note that $\frac{3}{2^{i+1}}$ is the midpoint of the interval $\left[\frac{1}{2^i}, \frac{1}{2^{i-1}}\right]$ and that the length of the interval is 2^{-i-1} . Then

$$\int_{[0,1]} f^+(x) dx = \sum_{i=1}^{\infty} \frac{2^{i+1}}{i} 2^{-i-1} = \sum_{i=1}^{\infty} \frac{1}{i} = \infty$$

and similarly

$$\int_{[0,1]} f^-(x) dx = \infty.$$

Thus f is not Lebesgue integrable in $[0, 1]$. However, the improper integral

$$\int_0^1 f(x) dx = \lim_{\epsilon \rightarrow 0} \int_{\epsilon}^1 f(x) dx = 0$$

exists because of the cancellation.

(4) Let $f : [0, \infty) \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} \frac{\sin x}{x}, & x > 0, \\ 1, & x = 0. \end{cases}$$

Observe that, since f is continuous, it is Lebesgue measurable.

Claim: $f \notin L^1([0, \infty))$.

Reason.

$$\begin{aligned} \int_{[0, \infty)} |f| dx &= \int_0^{\pi} |f(x)| dx + \sum_{i=1}^{\infty} \int_{i\pi}^{(i+1)\pi} \frac{|\sin x|}{|x|} dx \\ &\geq \int_0^{\pi} |f(x)| dx + \sum_{i=1}^{\infty} \frac{1}{i+1} = \infty. \end{aligned} \quad \blacksquare$$

Claim: The improper Riemann integral $\lim_{a \rightarrow \infty} \int_0^a \frac{\sin x}{x} dx$ exists.

Reason. Denote $I(a) = \int_0^a \frac{\sin x}{x} dx$, $a \geq \pi$. Then

$$I(k\pi) = \int_0^{k\pi} \frac{\sin x}{x} dx = \sum_{i=0}^{k-1} \int_{i\pi}^{(i+1)\pi} \frac{\sin x}{x} dx, \quad k = 1, 2, \dots,$$

where

$$\sum_{i=0}^{k-1} a_i \quad \text{with} \quad a_i = \int_{i\pi}^{(i+1)\pi} \frac{\sin x}{x} dx, \quad i = 1, 2, \dots,$$

is an alternating series with the properties

$$a_i a_{i+1} < 0, \quad |a_{i+1}| \leq |a_i| \quad \text{and} \quad \lim_{i \rightarrow \infty} a_i = 0.$$

Thus this series converges and

$$s = \sum_{i=0}^{\infty} \int_{i\pi}^{(i+1)\pi} \frac{\sin x}{x} dx = \lim_{k \rightarrow \infty} I(k\pi).$$

Since $a \geq \pi$, we have $a \in [k\pi, (k+1)\pi)$ for some $k = 1, 2, \dots$ and

$$|I(a) - I(k\pi)| = \left| \int_{k\pi}^a \frac{\sin x}{x} dx \right| \leq \int_{k\pi}^a \frac{1}{k\pi} dx \leq \frac{1}{k}.$$

This shows that $\lim_{a \rightarrow \infty} I(a) = s$. ■

Thus f is not Lebesgue integrable on $[0, \infty)$, but the improper integral $\int_0^{\infty} f dx$ exists (and equals to $\frac{\pi}{2}$ by complex analysis).

Remark 3.51. There exists an everywhere differentiable function such that its derivative is bounded but not Riemann integrable. Let $C \subset [0, 1]$ be a fat Cantor set with $m(C) = \frac{1}{2}$. Then

$$(0, 1) \setminus C = \bigcup_{i=1}^{\infty} I_i,$$

where I_i are pairwise disjoint open intervals and $\sum_{i=1}^{\infty} \text{vol}(I_i) = \frac{1}{2}$. For every $i = 1, 2, \dots$, choose a closed centered subinterval $J_i \subset I_i$ such that $\text{vol}(J_i) = \text{vol}(I_i)^2$. Define a continuous function $f : [0, 1] \rightarrow \mathbb{R}$ such that

$$f(x) = 0 \quad \text{for every } x \in [0, 1] \setminus \bigcup_{i=1}^{\infty} J_i,$$

$0 \leq f(x) \leq 1$ for every $x \in [0, 1]$ and $f(x) = 1$ at the center of every J_i . The set $\bigcup_{i=1}^{\infty} I_i$ is dense in $[0, 1]$, from which it follows that the upper Riemann integral is one and the lower Riemann integral is zero. Let

$$F(x) = \sum_{i=1}^{\infty} \int_{J_i \cap [0, x]} f(t) dt.$$

Then $F'(x) = f(x) = 0$ for every $x \in C$ (exercise) and $F'(x) = f(x)$ for every $x \in [0, 1] \setminus C$. Thus f is a derivative.

3.10 Fubini's theorem

We shall show that certain multiple integrals can be computed as iterated integrals. Moreover, under appropriate assumptions, the value of an iterated integral is independent of the order of integration.

Definition 3.52. Let μ be an outer measure on X and ν an outer measure on Y . We define the product outer measure $\mu \times \nu$ on $X \times Y$ as

$$(\mu \times \nu)(S) = \inf \left\{ \sum_{i=1}^{\infty} \mu(A_i) \nu(B_i) : S \subset \bigcup_{i=1}^{\infty} (A_i \times B_i) \right\},$$

where the infimum is taken over all collections of μ -measurable sets $A_i \subset X$ and ν -measurable sets $B_i \subset Y$, $i = 1, 2, \dots$.

THE MORAL : It is an exercise to show that $\mu \times \nu$ is an outer measure on $X \times Y$. The product outer measure is defined in such a way that product sets $A \times B$, where $A \subset X$ is a μ -measurable set and $B \subset Y$ is a ν -measurable set, inherit the natural measure $(\mu \times \nu)(A \times B) = \mu(A)\nu(B)$, see Fubini's theorem below.

Remark 3.53. Many texts develop the theory of product measures on products of measure spaces without outer measures, see [3], [8], [9] and [11].

Theorem 3.54 (Fubini's theorem). Let μ be an outer measure on X and ν an outer measure on Y .

- (1) Then $\mu \times \nu$ is a regular outer measure on $X \times Y$, even if μ and ν are not regular.
- (2) If $A \subset X$ is a μ -measurable set and $B \subset Y$ is a ν -measurable set, then $A \times B$ is $\mu \times \nu$ -measurable and $(\mu \times \nu)(A \times B) = \mu(A)\nu(B)$.
- (3) If $S \subset X \times Y$ is $\mu \times \nu$ -measurable and both measures μ and ν are σ -finite, then $S_y = \{x \in X : (x, y) \in S\}$ is μ -measurable for ν -almost every $y \in Y$ and $S_x = \{y \in Y : (x, y) \in S\}$ is ν -measurable for μ -almost every $x \in X$. Moreover,

$$(\mu \times \nu)(S) = \int_Y \mu(S_y) d\nu(y) = \int_X \nu(S_x) d\mu(x).$$

- (4) If f is $\mu \times \nu$ -measurable, both measures μ and ν are σ -finite and the integral of f is defined, that is, at least one of the functions f^+ and f^- has finite integral, then

$$y \mapsto \int_X f(x, y) d\mu(x)$$

is a ν -measurable function,

$$x \mapsto \int_X f(x, y) d\nu(y)$$

is a μ -measurable function and

$$\begin{aligned}\int_{X \times Y} f(x, y) d(\mu \times \nu) &= \int_Y \left(\int_X f(x, y) d\mu(x) \right) d\nu(y) \\ &= \int_X \left(\int_Y f(x, y) d\nu(y) \right) d\mu(x).\end{aligned}$$

T H E M O R A L : Claim (2) shows how to compute the product measure $(\mu \times \nu)(A \times B)$ of a product set by using the two measures $\mu(A)$ and $\nu(B)$. For a more general set S the product measure $(\mu \times \nu)(S)$ can be computed by integrating over the slices of the set S parallel to the coordinate axes by claim (3). Claim (4) shows that the integral of a function with respect to the product measure can be computed by iterated integrals.

Proof. **Step 1** Let \mathcal{F} denote the collection of all sets $S \subset X \times Y$ for which the integral

$$\int_X \chi_S(x, y) d\mu(x)$$

exists for ν -almost every $y \in Y$ and, in addition, such that

$$\rho(S) = \int_Y \left(\int_X \chi_S(x, y) d\mu(x) \right) d\nu(y)$$

exists. Note that $+\infty$ is allowed here.

Claim: If $S_i \in \mathcal{F}$, $i = 1, 2, \dots$, are pairwise disjoint, then $S = \bigcup_{i=1}^{\infty} S_i \in \mathcal{F}$.

Reason. Note that $\chi_S = \sum_{i=1}^{\infty} \chi_{S_i}$. By Corollary 3.16 we have $\rho(S) = \sum_{i=1}^{\infty} \rho(S_i)$. This shows that \mathcal{F} is closed under countable unions of pairwise disjoint sets. ■

Claim: If $S_i \in \mathcal{F}$, $i = 1, 2, \dots$, $S_1 \supset S_2 \supset \dots$, and $\rho(S_1) < \infty$, then $S = \bigcap_{i=1}^{\infty} S_i \in \mathcal{F}$.

Reason. Note that $\chi_S = \lim_{i \rightarrow \infty} \chi_{S_i}$. By the dominated convergence theorem we have

$$\rho(S) = \lim_{i \rightarrow \infty} \rho(S_i).$$

This shows that \mathcal{F} is closed under decreasing convergence of sets with a finiteness condition. ■

Let

$$\mathcal{P}_0 = \{A \times B : A \text{ is } \mu\text{-measurable and } B \text{ is } \nu\text{-measurable}\},$$

$$\mathcal{P}_1 = \left\{ \bigcup_{i=1}^{\infty} S_i : S_i \in \mathcal{P}_0 \right\} \quad \text{and} \quad \mathcal{P}_2 = \left\{ \bigcap_{i=1}^{\infty} S_i : S_i \in \mathcal{P}_1 \right\}.$$

The members of \mathcal{P}_0 are called measurable rectangles, the class \mathcal{P}_1 consists of countable unions of measurable rectangles and \mathcal{P}_2 of countable intersections of these. The latter sets constitute a class relative to which the product measure will be regular.

Note that $\mathcal{P}_0 \subset \mathcal{F}$ and

$$\rho(A \times B) = \mu(A)\nu(B)$$

whenever $A \times B \in \mathcal{P}_0$. If $A_1 \times B_1, A_2 \times B_2 \in \mathcal{P}_0$, then

$$(A_1 \times B_1) \cap (A_2 \times B_2) = (A_1 \cap A_2) \times (B_1 \cap B_2) \in \mathcal{P}_0$$

and

$$(A_1 \times B_1) \setminus (A_2 \times B_2) = ((A_1 \setminus A_2) \times B_1) \cup ((A_1 \cap A_2) \times (B_1 \setminus B_2)) \in \mathcal{P}_0$$

as a disjoint union of members of \mathcal{P}_0 . As in the proof of Theorem 1.11 it follows that every member of \mathcal{P}_1 is a countable union of pairwise disjoint members of \mathcal{P}_0 and hence $\mathcal{P}_1 \subset \mathcal{F}$.

Step 2 **Claim:** $(\mu \times \nu)(S) = \inf\{\rho(R) : S \subset R \in \mathcal{P}_1\}$ for every $S \subset X \times Y$.

Reason. Let $A_i \times B_i \in \mathcal{P}_0$, $i = 1, 2, \dots$ and $S \subset R = \bigcup_{i=1}^{\infty} (A_i \times B_i)$. Then $\chi_R \leq \sum_{i=1}^{\infty} \chi_{A_i \times B_i}$ and

$$\rho(R) \leq \sum_{i=1}^{\infty} \rho(A_i \times B_i) = \sum_{i=1}^{\infty} \mu(A_i)\nu(B_i).$$

Thus

$$\inf\{\rho(R) : S \subset R \in \mathcal{P}_1\} \leq (\mu \times \nu)(S).$$

Moreover, if $R = \bigcup_{i=1}^{\infty} (A_i \times B_i)$ is any such set, there exist pairwise disjoint sets $A'_i \times B'_i \in \mathcal{P}_0$ such that

$$R = \bigcup_{i=1}^{\infty} (A_i \times B_i) = \bigcup_{i=1}^{\infty} (A'_i \times B'_i).$$

Thus

$$\rho(R) = \sum_{i=1}^{\infty} \mu(A'_i)\nu(B'_i) \geq (\mu \times \nu)(S).$$

This shows that the equality holds. ■

Step 3 Fix $A \times B \in \mathcal{P}_0$. Then

$$(\mu \times \nu)(A \times B) \leq \mu(A)\nu(B) = \rho(A \times B) \leq \rho(R)$$

for every $R \in \mathcal{P}_1$ such that $A \times B \subset R$. Thus the claim above implies

$$(\mu \times \nu)(A \times B) = \mu(A)\nu(B).$$

We show that $A \times B$ is $\mu \times \nu$ -measurable. Let $R \in \mathcal{P}_1$ with $T \subset R$. Then $R \setminus (A \times B)$ and $R \cap (A \times B)$ are disjoint members of \mathcal{P}_1 . Thus

$$\begin{aligned} & (\mu \times \nu)(T \setminus (A \times B)) + (\mu \times \nu)(T \cap (A \times B)) \\ & \leq \rho(R \setminus (A \times B)) + \rho(R \cap (A \times B)) = \rho(R). \end{aligned}$$

The claim above implies

$$(\mu \times \nu)(T \setminus (A \times B)) + (\mu \times \nu)(T \cap (A \times B)) \leq (\mu \times \nu)(T).$$

Since this holds for all $T \subset X \times Y$, we have shown that $A \times B$ is $\mu \times \nu$ -measurable. Theorem 1.11 implies that $\mathcal{P}_0, \mathcal{P}_1$ and \mathcal{P}_2 consist of $\mu \times \nu$ -measurable sets. This proves claim (2) of the theorem.

Step 4 Next we show that $\mu \times \nu$ is a regular measure.

Claim: For every $S \subset X \times Y$ there exists $R \in \mathcal{P}_2$ such that $S \subset R$ and

$$(\mu \times \nu)(S) = (\mu \times \nu)(R) = \rho(R).$$

Reason. If $(\mu \times \nu)(S) = \infty$, set $R = X \times Y$. Thus we may assume that $(\mu \times \nu)(S) < \infty$. By the claim in (2), for every $i = 1, 2, \dots$ there exists a set $R_i \in \mathcal{P}_1$ such that $S \subset R_i$ and

$$\rho(R_i) < (\mu \times \nu)(S) + \frac{1}{i}.$$

Let $R = \bigcap_{i=1}^{\infty} R_i \in \mathcal{P}_2$. Since $R_i \in \mathcal{F}$ for every $i = 1, 2, \dots$, we conclude that $R \in \mathcal{F}$ and by the dominated convergence theorem

$$(\mu \times \nu)(S) \leq \rho(R) = \lim_{k \rightarrow \infty} \rho\left(\bigcap_{i=1}^k R_i\right) \leq (\mu \times \nu)(S).$$

This shows that $\mu \times \nu$ is \mathcal{P}_2 -regular. The claim follows from this, since every set in \mathcal{P}_2 is $\mu \times \nu$ -measurable by claim (2) of the theorem. \blacksquare

Step 5 If $S \subset X \times Y$ with $(\mu \times \nu)(S) = 0$, then there exists a set $R \in \mathcal{P}_2$ such that $S \subset R$ and $\rho(R) = 0$. Thus $S \in \mathcal{F}$ and $\rho(S) = 0$.

Assume that $S \subset X \times Y$ is $\mu \times \nu$ -measurable and $(\mu \times \nu)(S) < \infty$. Then there is $R \in \mathcal{P}_2$ such that $S \subset R$ and $(\mu \times \nu)(R \setminus S) = 0$. and, consequently, $\rho(R \setminus S) = 0$. It follows that

$$\mu(\{x \in X : (x, y) \in S\}) = \mu(\{x \in X : (x, y) \in R\})$$

for ν -almost every $y \in Y$ and

$$(\mu \times \nu)(S) = \rho(R) = \int_Y \mu(\{x \in X : (x, y) \in S\}) d\nu(y).$$

This proves claim (3) of the theorem, because the other formula is symmetric with X replaced by Y and μ by ν . The extension to σ -finite case can be done by exhausting the space by set of finite measure.

Step 6 Claim (4) reduces to (3) when $f = \chi_S$. If f is a nonnegative $\mu \times \nu$ -measurable function and is σ -finite with respect to $\mu \times \nu$, we use approximation by simple functions (Theorem 2.33) and the monotone convergence theorem (Theorem 3.12). Finally, for general f we consider $f = f^+ - f^-$. \square

As a corollary of Fubini's theorem we obtain Tonelli's theorem for nonnegative product measurable functions.

Theorem 3.55 (Tonelli's theorem). Let μ be an outer measure on X and ν an outer measure on Y and assume that both measures are σ -finite. Let $f : X \times Y \rightarrow [0, \infty]$ be a $\mu \times \nu$ -measurable function. Then

$$y \mapsto \int_X f(x, y) d\mu(x)$$

is a ν -measurable function,

$$x \mapsto \int_X f(x, y) d\nu(y)$$

is a μ -measurable function and

$$\begin{aligned} \int_{X \times Y} f(x, y) d(\mu \times \nu) &= \int_Y \left(\int_X f(x, y) d\mu(x) \right) d\nu(y) \\ &= \int_X \left(\int_Y f(x, y) d\nu(y) \right) d\mu(x). \end{aligned}$$

T H E M O R A L : The order of iterated integrals can be switched for all non-negative product measurable functions even in the case when the integrals are infinite.

Remarks 3.56:

(1) If μ and ν are counting measures, Tonelli's theorem reduces to a corresponding claim for series. Let $x_{i,j} \in [0, \infty]$, $i, j = 1, 2, \dots$. Then

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} x_{i,j} = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} x_{i,j}.$$

This means that we may rearrange the series without affecting the sum if the terms are nonnegative, compare with Corollary 3.16.

(2) Let μ be an outer measure on X and ν an outer measure on Y and suppose that both measures are σ -finite. Let $f : X \rightarrow [-\infty, \infty]$ be a $\mu \times \nu$ -measurable function. If any of the three integrals

$$\begin{aligned} &\int_{X \times Y} |f(x, y)| d(\mu \times \nu), \\ &\int_Y \left(\int_X |f(x, y)| d\mu(x) \right) d\nu(y), \\ &\int_X \left(\int_Y |f(x, y)| d\nu(y) \right) d\mu(x) \end{aligned}$$

is finite, then all of them are finite and the conclusion of Fubini's theorem holds (exercise). In particular, it follows that the function $y \mapsto f(x, y)$ is ν -integrable for μ -almost every $x \in X$ and that the function $x \mapsto f(x, y)$ is μ -integrable for ν -almost every $y \in Y$.

3.11 Fubini's theorem for Lebesgue measure

We discuss Tonelli's and Fubini's theorems for the Lebesgue measure, see [6, Chapter 8], [10, p. 75–86], [14, Chapter 6] and [9, Section 7.4]. By expressing $\mathbb{R}^{n+m} = \mathbb{R}^n \times \mathbb{R}^m$ it holds that

$$m^{n+m} = m^n \times m^m,$$

that is, the Lebesgue outer measure on \mathbb{R}^{n+m} is the product outer measure of the n -dimensional Lebesgue outer measure on \mathbb{R}^n and the m -dimensional Lebesgue outer measure on \mathbb{R}^m .

To see this, we observe that every interval I in \mathbb{R}^{n+m} can be represented as $I = J \times K$, where J is an interval in \mathbb{R}^n and K is an interval in \mathbb{R}^m . By Theorem 3.54 (2) and Lemma 1.60, we have

$$\begin{aligned} (m^n \times m^m)(I) &= (m^n \times m^m)(J \times K) = m^n(J)m^m(K) \\ &= \text{vol}_n(J)\text{vol}_m(K) = \text{vol}_{n+m}(J \times K) \\ &= \text{vol}_{n+m}(I) = m^{n+m}(I) \end{aligned}$$

for every interval I in \mathbb{R}^{n+m} . By Lemma 1.67 every open set $G \subset \mathbb{R}^{n+m}$ can be represented as a union of pairwise disjoint half open dyadic cubes, countable additivity on pairwise disjoint measurable sets implies that

$$(m^n \times m^m)(G) = m^{n+m}(G)$$

for all open sets $G \subset \mathbb{R}^{n+m}$. Since $m^n \times m^m$ and m^{n+m} are Radon measures on \mathbb{R}^{n+m} , by Corollary 1.51 (1) we have

$$\begin{aligned} (m^n \times m^m)(A) &= \inf\{(m^n \times m^m)(G) : A \subset G, G \subset \mathbb{R}^{n+m} \text{ open}\} \\ &= \inf\{m^{n+m}(G) : A \subset G, G \subset \mathbb{R}^{n+m} \text{ open}\} = m^{n+m}(A) \end{aligned}$$

for every $A \subset \mathbb{R}^{n+m}$.

In particular, we have $m^n = m^1 \times \dots \times m^1$ (n times), that is, the n -dimensional Lebesgue outer measure on \mathbb{R}^n is a product of the 1-dimensional Lebesgue outer measures, see Remark 1.76.

Theorem 3.57 (Tonelli's theorem). Let $f : \mathbb{R}^{n+m} \rightarrow [0, \infty]$ be a nonnegative m^{n+m} -measurable function. Then $y \mapsto f(x, y)$ is m^m -measurable for m^n -almost every $x \in \mathbb{R}^n$, $x \mapsto f(x, y)$ is m^n -measurable for m^m -almost every $y \in \mathbb{R}^m$,

$$y \mapsto \int_{\mathbb{R}^n} f(x, y) dm^n(x)$$

is a m^m -measurable function

$$x \mapsto \int_{\mathbb{R}^m} f(x, y) dm^m(y)$$

is a m^n -measurable function and

$$\begin{aligned} \int_{\mathbb{R}^{n+m}} f(x, y) dm^{n+m} &= \int_{\mathbb{R}^m} \left(\int_{\mathbb{R}^n} f(x, y) dm^n(x) \right) dm^m(y) \\ &= \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^m} f(x, y) dm^m(y) \right) dm^n(x). \end{aligned}$$

Remarks 3.58:

- (1) The function $x \mapsto f(x, y)$ is not necessary a m^n -measurable function for every $y \in \mathbb{R}^m$. Nor is the slice $A_y = \{x \in \mathbb{R}^n : (x, y) \in A\}$ a m^n -measurable set for every $y \in \mathbb{R}^m$. Let $E \subset \mathbb{R}$ be a set which is not m^1 -measurable and consider $A = E \times \{0\} = \{(x, y) \in \mathbb{R}^2 : x \in E, y = 0\}$. Then $m^2(A) = 0$ and thus A is m^2 -measurable, but A_y is not m^1 -measurable for $y = 0$. Note that measurability holds for almost every slice.
- (2) If A is m^{n+m} -measurable, then the slice $A_y = \{x \in \mathbb{R}^n : (x, y) \in A\}$ is m^n -measurable for m^m -almost every $y \in \mathbb{R}^m$. A corresponding statement holds with the roles of x and y interchanged. However, the reverse implication is not true in general. Let $E \subset \mathbb{R}$ be a set which is not m^1 -measurable and consider $A = [0, 1] \times E \subset \mathbb{R} \times \mathbb{R}$. Then

$$A_y = \begin{cases} [0, 1], & y \in E, \\ \emptyset, & y \notin E. \end{cases}$$

Thus A_y is m^1 -measurable for every $y \in \mathbb{R}$. However, if A were m^2 -measurable, then $A_x = \{y \in \mathbb{R} : (x, y) \in A\}$ were m^1 -measurable for almost every $x \in \mathbb{R}$. This is not true, since $A_x = E$ for every $x \in [0, 1]$. This implies that A is not m^2 -measurable. There exists a set $A \subset [0, 1] \times [0, 1]$, which is not m^2 -measurable with the property that A_y and A_x are m^1 -measurable for every $x, y \in [0, 1]$ with $m^1(A_y) = 0$ and $m^1(A_x) = 1$ for every $x, y \in [0, 1]$, see [10, p. 82-83].

Theorem 3.59 (Fubini's theorem). Let $f : \mathbb{R}^{n+m} \rightarrow [-\infty, \infty]$ be a m^{n+m} -measurable function and assume that at least one of the integrals

$$\begin{aligned} &\int_{\mathbb{R}^{n+m}} |f(x, y)| dm^{n+m}, \\ &\int_{\mathbb{R}^m} \left(\int_{\mathbb{R}^n} |f(x, y)| dm^n(x) \right) dm^m(y), \\ &\int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^m} |f(x, y)| dm^m(y) \right) dm^n(x), \end{aligned}$$

is finite. Then $y \mapsto f(x, y)$ is integrable in \mathbb{R}^m for m^n -almost every $x \in \mathbb{R}^n$, $x \mapsto f(x, y)$ is integrable in \mathbb{R}^n for m^m -almost every $y \in \mathbb{R}^m$,

$$y \mapsto \int_{\mathbb{R}^n} f(x, y) dm^n(x)$$

is integrable in \mathbb{R}^m ,

$$x \mapsto \int_{\mathbb{R}^m} f(x, y) dm^m(y)$$

is integrable in \mathbb{R}^n and

$$\begin{aligned} \int_{\mathbb{R}^{n+m}} f(x, y) dm^{n+m} &= \int_{\mathbb{R}^m} \left(\int_{\mathbb{R}^n} f(x, y) dm^n(x) \right) dm^m(y) \\ &= \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^m} f(x, y) dm^m(y) \right) dm^n(x). \end{aligned}$$

We consider few corollaries of the previous theorems.

Corollary 3.60. Assume that $f : \mathbb{R}^n \rightarrow [-\infty, \infty]$ is a m^n -measurable function. Then the function $\tilde{f} : \mathbb{R}^{n+m} \rightarrow [-\infty, \infty]$ defined by $\tilde{f}(x, y) = f(x)$ is a m^{n+m} -measurable function.

THEOREM: The trivial extension of a measurable function to higher dimensions is measurable.

Proof. We may assume that f is real valued. Since f is m^n -measurable, the set $A = \{x \in \mathbb{R}^n : f(x) < a\}$ is m^n -measurable for every $a \in \mathbb{R}$. Since

$$\{(x, y) \in \mathbb{R}^n \times \mathbb{R}^m : \tilde{f}(x, y) < a\} = A \times \mathbb{R}^m,$$

we conclude that the set is m^{n+m} -measurable for every $a \in \mathbb{R}$. Thus \tilde{f} is a m^{n+m} -measurable function. \square

Corollary 3.61. Assume that $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a m^n -measurable function and let

$$A = \{(x, y) \in \mathbb{R}^n \times \mathbb{R} : x \in \mathbb{R}^n, y = f(x)\}.$$

Then $m^{n+1}(A) = 0$ and thus $A \subset \mathbb{R}^{n+1}$ is m^{n+1} -measurable.

THEOREM: The graph of a measurable function is a set of measure zero.

Proof. Let $r > 0, \varepsilon > 0$ and

$$E_k = \{x \in B(0, r) : \varepsilon k \leq f(x) < (k+1)\varepsilon\}, \quad k \in \mathbb{Z}.$$

The sets E_k , $k \in \mathbb{Z}$, are m^n -measurable, pairwise disjoint and $B(0, r) = \bigcup_{k \in \mathbb{Z}} E_k$.

Let

$$A_k = \{(x, y) \in E_k \times \mathbb{R} : y = f(x)\}, \quad k \in \mathbb{Z}.$$

Then

$$\{(x, y) : x \in B(0, r), y = f(x)\} = \bigcup_{k \in \mathbb{Z}} A_k.$$

Since $A_k \subset E_k \times \{y \in \mathbb{R} : \varepsilon k \leq y < (k+1)\varepsilon\}$, $k \in \mathbb{Z}$, by Fubini's theorem, we have

$$\begin{aligned} m^{n+1}(A_k) &\leq m^{n+1}(E_k \times \{y \in \mathbb{R} : \varepsilon k \leq y < (k+1)\varepsilon\}) \\ &= m^n(E_k)m^1(\{y \in \mathbb{R} : \varepsilon k \leq y < (k+1)\varepsilon\}) = \varepsilon m^n(E_k), \quad k \in \mathbb{Z}. \end{aligned}$$

Thus we have

$$\begin{aligned} m^{n+1}(\{(x, y) : x \in B(0, r), y = f(x)\}) &= m^{n+1}\left(\bigcup_{k \in \mathbb{Z}} A_k\right) \leq \sum_{k \in \mathbb{Z}} m^{n+1}(A_k) \\ &\leq \sum_{k \in \mathbb{Z}} \varepsilon m^n(E_k) = \varepsilon m^n\left(\bigcup_{k \in \mathbb{Z}} E_k\right) = \varepsilon m^n(B(0, r)). \end{aligned}$$

Since this holds true for every $\varepsilon > 0$, we conclude that

$$m^{n+1}(\{(x, y) : x \in B(0, r), y = f(x)\}) = 0$$

for every $r > 0$. Finally, we note that $\mathbb{R}^n = \bigcup_{k=1}^{\infty} B(0, k)$ and thus

$$\begin{aligned} m^{n+1}(A) &= m^{n+1}\left(\bigcup_{k=1}^{\infty} \{(x, y) : x \in B(0, k), y = f(x)\}\right) \\ &\leq \sum_{k=1}^{\infty} m^{n+1}(\{(x, y) : x \in B(0, k), y = f(x)\}) = 0. \end{aligned} \quad \square$$

Corollary 3.62. Assume that $f : \mathbb{R}^n \rightarrow [0, \infty]$ is a nonnegative function and let

$$A = \{(x, y) \in \mathbb{R}^n \times \mathbb{R} : 0 \leq y \leq f(x)\}.$$

Then the following claims are true.

- (1) f is a m^n -measurable function if and only if A is a m^{n+1} -measurable set.
- (2) If the conditions in (1) hold, then

$$\int_{\mathbb{R}^n} f \, dm^n = m^{n+1}(A).$$

T H E M O R A L : The integral gives the area under the graph of a nonnegative measurable function.

Proof. Assume that f is a m^n -measurable function. By Corollary 3.60 the functions $(x, y) \mapsto -f(x)$ and $(x, y) \mapsto y$ are m^{n+1} -measurable and thus

$$F(x, y) = y - f(x)$$

is a m^{n+1} -measurable function. This implies that

$$\begin{aligned} A &= \{(x, y) \in \mathbb{R}^n \times \mathbb{R} : 0 \leq y \leq f(x)\} \\ &= \{(x, y) \in \mathbb{R}^n \times \mathbb{R} : y \geq 0\} \cap \{(x, y) \in \mathbb{R}^n \times \mathbb{R} : F(x, y) \leq 0\} \end{aligned}$$

is a m^{n+1} -measurable set.

Conversely, assume that A is m^{n+1} -measurable. For every $x \in \mathbb{R}^n$ the slice

$$A_x = \{y \in \mathbb{R} : (x, y) \in A\} = [0, f(x)]$$

is a closed one-dimensional interval. By Fubini's theorem $m^1(A_x) = f(x)$ is a measurable function for m^n -almost every x and

$$\begin{aligned} m^{n+1}(A) &= \int_{\mathbb{R}^{n+1}} \chi_A(x, y) dm^{n+1}(x, y) \\ &= \int_{\mathbb{R}^n} m(A_x) dm^n(x) \\ &= \int_{\mathbb{R}^n} f(x) dm^n(x). \end{aligned} \quad \square$$

We give an alternative proof for Cavalieri's principle by Fubini's theorem. Compare to Theorem 3.44. By Theorem 3.62, we have

$$m^{n+1}(A) = \int_{\mathbb{R}^n} m(A_x) dm^n(x) = \int_{\mathbb{R}^n} f(x) dm^n(x).$$

On the other hand, by Fubini's theorem we have

$$m^{n+1}(A) = \int_{\mathbb{R}} m(A_y) dm^1(y) = \int_0^\infty m(\{x \in \mathbb{R}^n : f(x) \geq y\}) dm^1(y),$$

where $A_y = \{x \in \mathbb{R}^n : (x, y) \in A\} = \{x \in \mathbb{R}^n : f(x) \geq y\}$, if $y \geq 0$, and $A_y = \emptyset$, if $y < 0$.

Thus

$$\int_{\mathbb{R}^n} f(x) dm^n(x) = \int_0^\infty m(\{x \in A : f(x) \geq y\}) dm^1(y).$$

We present a direct proof below.

Corollary 3.63. Let $A \subset \mathbb{R}^n$ be a Lebesgue measurable set and let $f : A \rightarrow [0, \infty]$ be a Lebesgue measurable function. Then

$$\int_A f dx = \int_0^\infty m(\{x \in A : f(x) > t\}) dt.$$

Proof.

$$\begin{aligned} \int_A f dx &= \int_{\mathbb{R}^n} \chi_A(x) f(x) dx \\ &= \int_{\mathbb{R}^n} \int_0^\infty \chi_A(x) \chi_{[0, f(x))}(t) dt dx \\ &= \int_0^\infty \int_{\mathbb{R}^n} \chi_A(x) \chi_{[0, f(x))}(t) dx dt \quad (\text{Fubini}) \\ &= \int_0^\infty \int_{\mathbb{R}^n} \chi_A(x) \chi_{\{x \in \mathbb{R}^n : f(x) > t\}}(x) dx dt \\ &= \int_0^\infty m(\{x \in A : f(x) > t\}) dt. \end{aligned} \quad \square$$

Remark 3.64. Note that

$$\begin{aligned} \int_A f dx &= \int_0^\infty m(\{x \in A : f(x) > t\}) dt \\ &= \int_0^\infty m(\{x \in A : f(x) \geq t\}) dt. \end{aligned}$$

Example 3.65. Let $A \subset \mathbb{R}^2$ be a Lebesgue measurable set with $m^2(A) = 0$. We claim that almost every horizontal line intersects A in a set whose one-dimensional Lebesgue measure is zero. The corresponding claim holds for vertical lines as well.

Reason. Let $A_1(y) = \{x \in \mathbb{R} : (x, y) \in A\}$ and $A_2(x) = \{y \in \mathbb{R} : (x, y) \in A\}$ with $x, y \in \mathbb{R}$. We shall show that $m^1(A_1(y)) = 0$ for almost every $y \in \mathbb{R}$ and, correspondingly, $m^1(A_2(x)) = 0$ for almost every $x \in \mathbb{R}$. Let $f = \chi_A$. Fubini's theorem implies

$$0 = m^2(A) = \int_{\mathbb{R}^2} \chi_A \, dm^2 = \int_{\mathbb{R}} \left(\int_{\mathbb{R}} f(x, y) \, dy \right) dx.$$

It follows that

$$m^1(A_2(x)) = \int_{\mathbb{R}} f(x, y) \, dy = 0$$

for almost every $x \in \mathbb{R}$. The proof for the claim $m^1(A_1(y)) = 0$ for almost every $y \in \mathbb{R}$ is analogous. \blacksquare

Conversely, if $A \subset \mathbb{R}^2$ is a Lebesgue measurable set such that $m^1(A_1(y)) = 0$ for almost every $y \in \mathbb{R}$ or $m^1(A_2(x)) = 0$ for almost every $x \in \mathbb{R}$, then $m^2(A) = 0$.

Reason. Fubini's theorem for the measurable function $f = \chi_A$. \blacksquare

W A R N I N G : The assumption that $A \subset \mathbb{R}^2$ is measurable is essential. Indeed, there exist a set $A \subset \mathbb{R}^2$ such that

- (1) A is not Lebesgue measurable and thus $m(A) > 0$,
- (2) every horizontal line intersects A at most one point and
- (3) every vertical line intersects A at most one point.

(Sierpinski: Fundamenta Mathematica 1 (1920), p. 114)

The examples below show how we may apply Fubini's theorem to evaluate certain integrals.

Examples 3.66:

- (1) We show that

$$I = \int_{-\infty}^{\infty} e^{-x^2} \, dx = \sqrt{\pi}$$

in two ways.

(1) Note that $e^{-x^2} > 0$ for every $x \in \mathbb{R}$ and

$$I \leq \int_{-\infty}^{-1} -xe^{-x^2} \, dx + \int_{-1}^1 e^{-x^2} \, dx + \int_1^{\infty} xe^{-x^2} \, dx < \infty.$$

Since $x \mapsto e^{-x^2}$ is even, we have

$$I = 2 \int_0^{\infty} e^{-x^2} \, dx.$$

It follows that

$$I^2 = \left(\int_{-\infty}^{\infty} e^{-x^2} dx \right) \left(\int_{-\infty}^{\infty} e^{-y^2} dy \right) = 4 \int_0^{\infty} \left(\int_0^{\infty} e^{-(x^2+y^2)} dy \right) dx.$$

Substitution $y = xs$ implies $dy = x ds$. By Fubini's theorem

$$\begin{aligned} \frac{I^2}{4} &= \int_0^{\infty} \left(\int_0^{\infty} e^{-(1+s^2)x^2} x ds \right) dx = \int_0^{\infty} \left(\int_0^{\infty} e^{-(1+s^2)x^2} x dx \right) ds \\ &= \frac{1}{2} \int_0^{\infty} \frac{1}{1+s^2} ds = \frac{1}{2} \arctan s \Big|_0^{\infty} = \frac{\pi}{4}. \end{aligned}$$

Thus $I = \sqrt{\pi}$.

(2) By the polar coordinates, we have

$$\begin{aligned} \int_{\mathbb{R}^2} e^{-(x^2+y^2)} dx dy &= \lim_{i \rightarrow \infty} \int_{B(0,i)} e^{-(x^2+y^2)} dx dy = \lim_{i \rightarrow \infty} \int_{B(0,i)} e^{-(x^2+y^2)} dx dy \\ &= \lim_{i \rightarrow \infty} \int_0^i \int_0^{2\pi} e^{-r^2} r dr d\theta = 2\pi \lim_{i \rightarrow \infty} \int_0^i e^{-r^2} r dr \\ &= \pi \lim_{i \rightarrow \infty} -e^{-r^2} \Big|_0^i = -\pi \lim_{i \rightarrow \infty} (e^{-i^2} - 1) = \pi, \end{aligned}$$

where all integrals are (possibly improper) Riemann integrals. By the Lebesgue monotone convergence theorem the Riemann and Lebesgue integrals

$$\int_{\mathbb{R}^2} e^{-(x^2+y^2)} dx dy$$

coincide. By Fubini's theorem

$$\begin{aligned} \pi &= \int_{\mathbb{R}^2} e^{-(x^2+y^2)} dx dy = \int_{\mathbb{R}} \left(\int_{\mathbb{R}} e^{-x^2} e^{-y^2} dy \right) dx \\ &= \int_{\mathbb{R}} e^{-x^2} \left(\int_{\mathbb{R}} e^{-y^2} dy \right) dx = \left(\int_{\mathbb{R}} e^{-x^2} dx \right)^2 \end{aligned}$$

and thus

$$\int_{\mathbb{R}} e^{-x^2} dx = \sqrt{\pi}.$$

(2) Consider

$$\int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} dx, \quad a, b > 0.$$

Since

$$\frac{e^{-ax} - e^{-bx}}{x} = \int_a^b e^{-xy} dy,$$

we have

$$\int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} dx = \int_0^{\infty} \int_a^b e^{-xy} dy dx.$$

The function $(x, y) \mapsto e^{-xy}$ is continuous and thus Lebesgue measurable.

Since $e^{-xy} > 0$, we have

$$\int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} dx = \int_a^b \int_0^{\infty} e^{-xy} dx dy = \int_a^b \frac{1}{y} dy = \log \frac{b}{a}$$

The following examples show that we cannot always switch the order of integration in iterated integrals.

Examples 3.67:

(1) Consider

$$\int_0^1 \int_0^1 \frac{x^2 - y^2}{(x^2 + y^2)^2} dy dx.$$

Note that

$$\begin{aligned} \int_0^1 \frac{x^2 - y^2}{(x^2 + y^2)^2} dy &= \int_0^1 \frac{x^2 + y^2}{(x^2 + y^2)^2} dy + \int_0^1 \frac{-2y^2}{(x^2 + y^2)^2} dy \\ &= \int_0^1 \frac{1}{x^2 + y^2} dy + \int_0^1 y \left(\frac{d}{dy} \frac{1}{x^2 + y^2} \right) dy. \end{aligned}$$

An integration by parts gives

$$\begin{aligned} \int_0^1 y \left(\frac{d}{dy} \frac{1}{x^2 + y^2} \right) dy &= \frac{y}{x^2 + y^2} \Big|_0^1 - \int_0^1 \frac{1}{x^2 + y^2} dy \\ &= \frac{1}{x^2 + 1} - \int_0^1 \frac{1}{x^2 + y^2} dy. \end{aligned}$$

Thus

$$\int_0^1 \frac{x^2 - y^2}{(x^2 + y^2)^2} dy = \frac{1}{x^2 + 1},$$

from which it follows that

$$\int_0^1 \int_0^1 \frac{x^2 - y^2}{(x^2 + y^2)^2} dy dx = \int_0^1 \frac{1}{x^2 + 1} dx = \arctan x \Big|_0^1 = \frac{\pi}{4}.$$

By symmetry

$$\int_0^1 \int_0^1 \frac{x^2 - y^2}{(x^2 + y^2)^2} dx dy = - \int_0^1 \int_0^1 \frac{x^2 - y^2}{(x^2 + y^2)^2} dy dx = -\frac{\pi}{4}.$$

Observe that in this case both iterated integrals exist and are finite, but they are not equal. This does not contradict Fubini's theorem, since

$$\begin{aligned} \int_0^1 \int_0^1 \left| \frac{x^2 - y^2}{(x^2 + y^2)^2} \right| dy dx &= \int_0^1 \left(\int_0^x \frac{x^2 - y^2}{(x^2 + y^2)^2} dy + \int_x^1 \frac{y^2 - x^2}{(x^2 + y^2)^2} dy \right) dx \\ &= \int_0^1 \frac{1}{x} dx - \int_0^1 \frac{1}{x^2 + 1} dx = \infty. \end{aligned}$$

The integral in the brackets can be evaluated by integration by parts.

(2) Let

$$A = \bigcup_{i=0}^{\infty} \chi_{[i, i+1] \times [i, i+1]} \quad \text{and} \quad B = \bigcup_{i=0}^{\infty} \chi_{[i+1, i+2] \times [i, i+1]}.$$

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, $f = \chi_A - \chi_B$. Then

$$\int_{\mathbb{R}} f(x, y) dx = 0$$

for every $y \in \mathbb{R}$ and thus

$$\int_{\mathbb{R}} \left(\int_{\mathbb{R}} f(x, y) dx \right) dy = 0.$$

On the other hand,

$$\int_{\mathbb{R}} f(x, y) dy = \chi_{[0,1]}(x)$$

for every $x \in \mathbb{R}$ and thus

$$\int_{\mathbb{R}} \left(\int_{\mathbb{R}} f(x, y) dy \right) dx = 1.$$

It follows that

$$\int_{\mathbb{R}} \left(\int_{\mathbb{R}} f(x, y) dx \right) dy = 0 \neq 1 = \int_{\mathbb{R}} \left(\int_{\mathbb{R}} f(x, y) dy \right) dx.$$

(3) Let $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} 2^{2i}, & 2^{-i} \leq x < 2^{-i+1}, 2^{-i} \leq y < 2^{-i+1}, \\ -2^{2i+1}, & 2^{-i-1} \leq x < 2^{-i}, 2^{-i} \leq y < 2^{-i+1}, \\ 0, & \text{otherwise.} \end{cases}$$

Then (exercise)

$$\int_{[0,1]} \left(\int_{[0,1]} f(x, y) dx \right) dy = 0 \neq 1 = \int_{[0,1]} \left(\int_{[0,1]} f(x, y) dy \right) dx.$$

(4) Let $0 = \delta_1 < \delta_2 < \dots < 1$ and $\delta_i \rightarrow 1$ as $i \rightarrow \infty$. Let g_i , $i = 1, 2, \dots$ be continuous functions such that

$$\text{supp } g_i \subset (\delta_i, \delta_{i+1}) \quad \text{and} \quad \int_0^1 g_i(t) dt = 1$$

for every $i = 1, 2, \dots$. Let

$$f(x, y) = \sum_{i=1}^{\infty} (g_i(x) - g_{i+1}(x)) g_i(y).$$

The function f is continuous except at the point $(1, 1)$, but

$$\int_0^1 \int_0^1 f(x, y) dy dx = 1 \neq 0 = \int_0^1 \int_0^1 f(x, y) dx dy.$$

THE END

Bibliography

- [1] V. I. Bogachev, Measure Theory 1 and 2, Springer, 2007.
- [2] A. M. Bruckner, J.B. Bruckner, and B.S. Thomson, Real Analysis, Prentice-Hall, 1997.
- [3] D. L. Cohn, Measure theory (2nd edition), Birkhäuser, 2013.
- [4] L. C. Evans and R. F. Gariepy, Measure Theory and Fine Properties of Functions, CRC Press, 1992.
- [5] G. B. Folland, Real Analysis. Modern Techniques and Their Applications (2nd edition), John Wiley & Sons, 1999.
- [6] F. Jones, Lebesgue Integration on Euclidean Space (revised edition), Jones and Bartlett Publishers, 2001.
- [7] K. L. Kuttler, Modern Real Analysis, CRC Press, 1998.
- [8] W. Rudin, Real and Complex Analysis, McGraw-Hill, 1986.
- [9] D. A. Salamon, Measure and Integration, EMS, 2016.
- [10] E. Stein and R. Sakarchi, Real Analysis: Measure Theory, Integration, and Hilbert Spaces, Princeton University Press, 2005.
- [11] T. Tao, Introduction to Measure Theory, American Mathematical Society, 2011.
- [12] M.E. Taylor, Measure theory and integration, American Mathematical Society, 2006.
- [13] S. Wagon, The Banach-Tarski Paradox, Encyclopedia of Mathematics and its Applications, Vol. 24, Cambridge University Press, 1985.
- [14] R. L. Wheeden and A. Zygmund, Measure and Integral: An introduction to Real Analysis, Marcel Dekker, 1977.
- [15] W. P. Ziemer, Modern Real Analysis, PWS Publishing Company, 1995.
- [16] J. Yeh, Real Analysis, Theory of Measure and Integration (2nd edition), World Scientific, 2006.

Index

- σ -algebra, 9
 - Borel, 22
 - generated, 22
- σ -finite measure, 15
 - almost everywhere, 83
 - almost everywhere convergence, 89
 - almost uniform convergence, 90
- Borel
 - function, 70
 - outer measure, 23
 - regular, 23
 - set, 22
- Cantor set, 61, 143
- Cantor-Lebesgue function, 72
- Carathéodory's construction, 6
- Cavalieri's principle, 136, 153
- complete measure, 15, 82, 84
- completion of a measure space, 15
- convergence
 - almost everywhere, 89
 - almost uniform, 90
 - in measure, 90
 - locally uniform, 97
 - monotone, 108
 - pointwise, 88
 - uniform, 88
- convergence in measure, 90
- countable subadditivity, 1
- counting measure, 3
- cube, 39
 - dyadic, 46
- de Morgan's law, 9
- diameter, 19
- Dirac measure, 2
- Discrete measure, 2
- distance function, 19
 - cutoff function, 21
- exhaustion of an open set, 20
- Lipschitz continuity, 20
- distribution function, 135
- dominated convergence theorem, 116
- dyadic, 46
 - collection, 46
 - covering, 46
 - cube, 46
 - half-open, 46
 - nesting property, 46
- Egoroff's theorem, 94
- extension of a measure, 16
- Fatou's lemma, 112
- Fubini's theorem, 144
 - Lebesgue measure, 150
- Hahn-Kolmogorov theorem, 16
- Hausdorff measure, 5
- integrable function, 104, 113
- integral
 - nonnegative measurable function, 104
 - nonnegative simple function, 101
 - Riemann, 140
 - signed function, 113
- interval, 3
 - almost disjoint, 39
 - closed, 3
 - degenerate, 3, 4, 49
 - Lebesgue measure, 41
 - open, 41
- Jordan measure, 38
- Lebesgue integral, 123
 - L^1 convergence, 127
 - L^1 space, 126
 - change of variables, 128, 129
 - invariance properties, 128

polar coordinates, 130
Lebesgue measurable set, 54, 56
Lebesgue measure, 3, 56
 σ -finite, 16
 Borel measure, 36
 Borel regular, 24
 finite, 3
 infinite, 3
 linear change of variables, 52
 locally finite, 24
 metric outer measure, 36
 nonmeasurable set, 58
 of an interval, 41
 of the entire space, 49
 outer measure, 3, 4
 product structure, 149
 Radon measure, 36
 reflection invariance, 51
 rotation invariance, 52
 scaling property, 51
 translation invariance, 51
 zero, 3, 48
Littlewood's principles, 97
locally uniform convergence, 97
Lusin's theorem, 97

measurable function, 66, 67, 68, 69, 71, 71, 80
 approximation by simple functions, 85, 87
measurable set, 7
 σ -algebra, 10
 Lebesgue measure, 54, 56
 monotone converge theorem, 16
 Radon measure, 28, 30
measure, 13
 σ -finite, 15
 complete, 15
 completion, 15
 extension, 16, 16
 induced by an outer measure, 16
measure space, 13

metric outer measure, 34
monotone convergence theorem, 108
negative part of a function, 71, 113

outer measure, 1
 σ -finite, 15
 Borel, 23
 Borel regular, 23
 complete, 15
 countable additivity, 13
 countable subadditivity, 1
 induced by a measure, 16
 Lebesgue, 3
 locally finite, 24
 metric, 34
 monotonicity, 1
 Radon, 23
 regular, 24
 restriction, 27

pointwise convergence, 88
polar coordinates, 130
positive part of a function, 71, 113
product outer measure, 144

Radon outer measure, 23
restriction of an outer measure, 27
Riemann integral, 140

simple function, 84
 approximation of a measurable function, 85, 87
 integral, 101

Tchebyshev's inequality, 135
 simple functions, 105
Tieze's extension theorem, 100, 100
Tonelli's theorem, 148
 Lebesgue measure, 149

uniform convergence, 88